

## RSV ComNet: Data collection progress seasons 2021/22 & 2022/23

### Aim

The aim of the RSV ComNet study is to measure the clinical burden and socio-economic impact of a respiratory syncytial virus (RSV) infection in young children (<5 years old) in primary care.<sup>1,2</sup>

### Background

In this bi-weekly update, we provide an update on developments regarding data collection. Due to low RSV circulation in the 2021/22 season, data collection is extended during the 2022/23 winter. Countries have restarted data collection around week 40 2022.

This update includes detailed information on the total number of swabs collected and the total number of RSV+ cases identified over both seasons. In addition the response rate on the parental Day 14 and Day 30 questionnaires for the 2022/23 season is shown. Data for earlier weeks can slightly change because of reporting delays.

The purpose of this update is to allow for the inter-country comparison of progress towards data collection objectives and to discover trends in data collection across countries. In later stages, these trends will be elaborated upon, keeping in mind differences in recruitment procedures between countries.

### Key points:

- Since the start of data collection, in all countries together:
  - **2637 swabs** have been collected
  - **1010** tested **RSV positive**
  - **682** parents have completed **at least one questionnaire\***
- Italy has collected 770 swabs, the United Kingdom 757, Belgium 425, Spain 351, the Netherlands 319 and France 15†.
- Number of participants with at least one completed questionnaire: Italy: 215, the United Kingdom: 83, Belgium: 139, Spain: 152, the Netherlands: 89 and France: 4† (see page 6).

\*The first questionnaire is reported 14 days after an RSV+ test result

†Data collection in France started significantly later than in the other countries, in week 50 in 2022.

- Although swabs are still collected, the number of RSV positive cases has considerably decreased in all countries over the last weeks.

### References

1 van Summeren et al. BMC Infect Dis. 2021 Jul 26;21(1):705

2 <https://www.nivel.nl/en/RSVComNet>


### More information?

Go to [www.nivel.nl/en/RSVComNet](http://www.nivel.nl/en/RSVComNet)

Mail: [j.vansummeren@nivel](mailto:j.vansummeren@nivel.nl)

## Total number of swabs as a fraction of the planned number of swabs


This figure shows per country the number of swabs that were collected as a fraction of the planned number of swabs. The planned number of swabs varies between countries and is indicated in the legend.



\* For the Netherlands, data collection is slightly different compared to the other countries: 1) data collection is stretched out over 2 years, 2) only children under the age of 2 years are included, and 3) data collection ends when 150 RSV positive cases are included, instead of aiming for a specific number of swabs.

## Cumulative number of RSV+ children


This figure shows the cumulative number of RSV+ cases that were identified in each country. Not all parents of RSV+ children have completed the questionnaires.




## Response on parental questionnaires season 2022/23

In the following figures, the weekly number of RSV+, Day 14 and Day 30 questionnaires are presented (left y-axis) from the week that data collection has been started.

**Note:** The receipt of the Day 14 and Day 30 questionnaires is aligned with the week the swab was taken. This implies that for the Day 14 and Day 30 questionnaires respectively the last two weeks and last four weeks are not up to date, as the questionnaires for those weeks have yet to be received.




- In total **154 children tested RSV+**, **83 parents** completed the **Day-14** questionnaire and **40 parents** the **Day-30** questionnaire.
- In the UK, 25 GP practices participating in the sentinel surveillance are also participating in the RSV ComNet study. These GPs will invite parents of children that tested RSV+ to complete the RSV ComNet follow-up questionnaires. This means that it is possible that not all parents of RSV+ children give informed consent to complete follow-up questionnaires.




- In total **160 children tested RSV+**, **152 parents** completed the **Day-14** questionnaire and **138 parents** the **Day-30** questionnaire.
- In Spain 17 pediatrician practices are participating in the RSV ComNet study in 9 regions.

## Response on parental questionnaires season 2022/23




- In total **179 children tested RSV+**, **139 parents** completed the **Day-14** questionnaire and **118 parents** the **Day-30** questionnaire.
- In Belgium, 15 pediatricians are participating in the RSV ComNet study.




- In total **115 children tested RSV+**, **89 parents** completed the **Day-14** questionnaire and **74 parents** the **Day-30** questionnaire.
- In the NL, 44 GP practices are participating in the RSV ComNet study.

## Response on parental questionnaires season 2022/23



- In total **398 children tested RSV+**, **215 parents** completed the **Day-14** questionnaire and **105 parents** the **Day-30** questionnaire.
- In Italy, pediatricians from four regions are participating in the RSV ComNet study.




- In total **4 children tested RSV+**, **4 parents** completed the **Day-14** questionnaire and **3 parents** the **Day-30** questionnaire.
- In France, data collection started in week 50 2022, after the RSV epidemic passed the peak for the winter of 2022/2023.


## RSV positivity rate since start data collection

In this figure the RSV positivity rate is presented for each country. The RSV positivity rate is calculated by dividing the total number of RSV+ cases by the total number of swabs collected since the first week of data collection.

Data collected via the RSV ComNet study cannot be considered as surveillance data. This is because only a small selection of GPs and/or pediatricians are participating in the RSV ComNet study, which is not a representative reflection for the country as a whole. In addition, opportunistic sampling is applied which means that swabbing is increased in weeks with a suspected increase in RSV circulation.



## Total number of patients with at least one completed parental questionnaire



### More information?

Go to [www.nivel.nl/en/RSVComNet](http://www.nivel.nl/en/RSVComNet)

Mail: [j.vansummeren@nivel.nl](mailto:j.vansummeren@nivel.nl)