

NIV 1.4(2)

DUTCH NATIONAL SURVEY OF GENERAL PRACTICE

MORBIDITY IN GENERAL PRACTICE

J. van der Velden
D.H. de Bakker
A.A.M.C. Claessens
F.G. Schellevis

August 1992

Netherlands Institute of Primary Health Care
(NIVEL)
P.O. Box 1568 - 3500 BN Utrecht - tel. nr. ...-3130319946



bibliotheek
drieharingstraat 6
postbus 1568
3500 bn utrecht
tel 030 2729 614/615
fax 030 2729729

CIP-GEGEVENS KONINKLIJKE BIBLIOTHEEK, DEN HAAG

Morbidity

Morbidity in general practice: Dutch National Survey of Morbidity
and Interventions in General Practice / J. van der Velden (et al.).

Utrecht: Netherlands Institute of Primary Health Care (NIVEL)

Met lit. opg.

ISBN 90-6905-189-3

Trefw.: medische statistiek; huisartsgeneeskunde.

Lay-out: Mieke Cornelius

Tekstprocessing: Christine Ruiter

PREFACE

This report of the Dutch National Survey of (diseases and interventions in) general practice bears the title "Morbidity in General Practice".

The report is the third in a series of four reports. The first report contains a detailed description of measuring instruments and procedures followed in data collection. The second report is called "Health status and health behaviour in the practice population and the fourth "Interventions in general practice".

This report gives an overview of the morbidity, presented to general practice. The quantitative information is representative for Dutch general practice. The description of the methods of data-collection has been restricted, because of its comprehensive description in the first report. However, extensive attention has been given to specific calculations carried out for this report (calculation of incidence, prevalence and weighting factors).

Besides a total rate, there is a splitting up in characteristics of the patient and characteristics of general practice. In this way morbidity is represented by subpopulations distinguished to demographic (like age, sex, marital status), socio-economic (like occupational class, educational level, insurance-type) and geographic characteristics of general practice (like region and degree of urbanization).

The data, represented in this report, are the basis to further investigation of the work of the general practitioner and the position of the general practitioner in the health care system of the Netherlands.

Utrecht, August 1992

ACKNOWLEDGEMENTS

Many people contributed to the realization of this report. In the first place this concerns the 161 general practitioners, the 193 practice-assistants, the 90 fieldworkers/datatypists and the fieldwork-organization, who were successful with each other in the collection and the processing of more than 300.000 episodes of illness of 335.000 patients.

In the quality control of data and production of the final tables of morbidity many people contributed too. This applies especially to Harald Abrahamse, Marleen Foets, Jan Gravestein, Han van Snellenberg, Loek Stokx, Patricia Swords and last but not least Jan van der Steen.

TABLES

- 2.1 Registering general practitioners and other physicians by stratification variables
- 2.2 Registering general practitioners and other physicians by season
- 2.3 Response to patient registration by age and sex, stratification variables and season
- 2.4 Significance level of differences in average number of episodes per 1000 patients of each ICPC-chapter between general practices broken down by stratification variables
- 2.5 Practice population National Survey and Dutch population by age and sex
- 2.6 Weighting factors step 1
- 2.7 Calculation of the weighting factors distance from practice address to the hospital by degree of urbanization and region
- 2.8 Calculation of correction factors per registration period
- 2.9 Age/sex distribution of practice population before and after weighting compared with the Netherlands
- 2.10 The investigated practice population by marital status
- 2.11 The investigated practice population by highest level of education attained
- 2.12 The investigated practice population by form of insurance
- 2.13 The investigated practice population by country of birth
- 2.14 The investigated practice population by occupational class
- 2.15 The investigated practice population by current employment status
- 2.16 The investigated practice population by living situation
- 2.17 Practice population by urbanization degree, region and distance from hospital and season
- 2.18 Percentages of agreement by fieldworkers in the classification of diagnoses
- 2.19 Incompatibility of diagnosis/age and of diagnosis/sex in percentages of the contacts and episodes respectively

- 3.1 Incidence and prevalence per ICPC chapter after weighting per 3 months absolute and per 1000 persons
- 3.2 Average incidence/prevalence with including variation coefficient per ICPC-chapter
- 3.3 Incidence per ICPC-chapter by degree of urbanization after weighting per 3 months per 1000 persons
- 3.4 Prevalence per ICPC-chapter by degree of urbanization after weighting per 3 months per 1000 persons
- 3.5 Incidence per ICPC-chapter by region after weighting per 3 months per 1000 persons
- 3.6 Prevalence per ICPC-chapter by region after weighting per 3 months per 1000 persons
- 3.7 Incidence per ICPC-chapter by the distance to hospital after weighting per 3 months per 1000 persons
- 3.8 Prevalence per ICPC-chapter by the distance to hospital after weighting per 3 months per 1000 persons
- 3.9 Incidence per ICPC-chapter by season after weighting per 3 months per 1000 persons
- 3.10 Prevalence per ICPC-chapter by season after weighting per 3 months per 1000 persons
- 3.11 Incidence and prevalence per ICPC chapter by sex after weighting per 3 months per 1000 persons
- 3.12 Incidence and prevalence per ICPC-chapter by age categories after weighting per 3 months per 1000 persons
- 3.13 Incidence and prevalence per ICPC-chapter by marital status after weighting per 3 months per 1000 persons
- 3.14 The 20 highest incident and prevalent diagnoses per 3 months per 1000 persons
- 3.15 The 20 highest incident diagnosis among males and females per 3 months per 1000 persons
- 3.16 The 20 highest prevalent diagnoses among males and females per 3 months per 1000 persons

FIGURES

- 1.1 The professional care process
- 2.1 Contact registration form of the National Survey
- 2.2 Percentage of agreement by general practitioners on paper vignettes
- 3.1 Incidence and prevalence of ICPC-chapters by age and sex after weighting per 3 months per 1000 persons
- 3.2 Incidence and prevalence per ICPC-chapter by form of insurance after weighting per 3 months per 1000 persons
- 3.3 Incidence and prevalence per ICPC-chapter by family status after weighting per 3 months per 1000 persons
- 3.4 Incidence and prevalence per ICPC-chapter by birth country after weighting per 3 months per 1000 persons
- 3.5 Incidence and prevalence per ICPC-chapter by educational level after weighting per 3 months per 1000 persons
- 3.6 Incidence and prevalence per ICPC-chapter by current-employment status after weighting per 3 months per 1000 persons
- 3.7 Incidence and prevalence per ICPC chapter by occupational class after weighting per 3 months per 1000 persons

CONTENTS

page

PREFACE

ACKNOWLEDGEMENTS

1. INTRODUCTION	1
1.1 Introduction	1
1.2 Backgrounds and objectives	2
1.3 Outline of the report	4
2. DATA AND METHODS	5
2.1 Introduction	5
2.2 Sample of general practitioners	5
2.3 The epidemiological numerator	7
2.3.1 Encounter registration	7
2.3.2 From problem to episode	10
2.3.3 Classification of morbidity	10
2.3.4 Summary	12
2.4 The epidemiological denominator	12
2.4.1 Patient registration	12
2.4.2 Weighting	14
2.4.3 Description of denominator per patient characteristic	19
2.4.4 Description of denominator per practice characteristic	23
2.4.5 Summary	24
2.5 Incidence and prevalence	25
2.6 Validity and reliability of data	26
2.6.1 Validity	26
2.6.2 Reliability	28
3. RESULTS	31
3.1 Introduction	31
3.2 Morbidity per ICPC chapter by practice characteristics and season	31
3.3 Morbidity per ICPC chapter by patients characteristics	41

LITERATURE

- Appendix A Basic tables of morbidity in general practice of the Netherlands
- A.1 Clarification
 - A.2 Abbreviations, used in the basic tables
 - A.3 Basic table A: morbidity in general practice by age and sex
 - A.4 Basic table B: morbidity in general practice by urbanization degree, region, season and insurance mode
- Appendix B Diagnoses (numbers) corresponding with ICPC-codes of NIVEL-edition (1989) and Oxford-edition (1987)

1. INTRODUCTION

1.1 Introduction

Statistical information on the health of the population has long been used as an instrument for health policy and planning of health care and evaluation thereof. The presentation of this in the "Nota 2000" may be described as the most recent and telling example of this. The working paper "Epidemiologie en Informatica", a part of the "Kerndocument Gezondheidsbeleid" (1989), elaborates on this tradition and is directed towards the following goals:

- insight into the health status of the population;
- insight into effects and quality of the care;
- systematic evaluation of health policy;
- improvement of the provision of information in health care.

Recently extensive surveys have become available on what is present in the way of health statistics in the Netherlands (Bonte, 1989; Hoogendoorn, 1989; Pieters, 1989; Hofman, 1989).

Broadly speaking, with regard to public health the following data are present structurally and routinely:

- cause-of-death and stillbirth statistics for which there is a legal notification requirement by physicians;
- statistics concerning specific disorders (cancer registration, surveillance of infectious diseases, industrial accidents);
- statistics from medical examinations and checks (absenteeism, incapacity for work, conscripts);
- statistics from specific care situations (diagnosis statistics of hospitals, sentinel stations and morbidity registration by general practitioners, obstetric assistance and place of birth);
- statistics from the population (health surveys, living situation surveys).

This variety of registrations, based on and organized around specific objectives, gives a good but not yet complete insight into public health in the Netherlands. Moreover, various methodological problems play a part in the interpretation of data derived from these registrations on behalf of health policy.

The following gaps have been noted (Bonte, 1989):

1. As part of the Health Survey as performed by the Central Statistical Office (CBS): the absence as yet of data on the population living in institutions; observations on health-affecting aspects of life such as eating, drinking and sport; data directed towards certain sections of the population such as migrants.
2. Data on the health status of health of the population with regard to the 'objective component' are only fragmentary: surveys of congenital defects and data representative of the population on diseases, disturbances and limitations, as established by physicians, are missing.

3. Statistics on incapacity for work are incomplete and not fully geared to each other, while data on occupational diseases are due for revision.
4. Differentiation of a number of statistics by socio-economic characteristics is desirable.
5. As part of the services supplied the Health Survey to a certain extent fills the gaps regarding consumption in the extramural sector (in the intramural sector registrations by among others the SIG usually supply the necessary data. Special mention should be made of the absence of (diagnostic) data on mental public health and extramural and semi-mural care, such as morbidity data in general practice, outpatient department and day and night nursing.

The importance of data representative of the population on disease, disturbances and limitations, as established by physicians, is emphasized by an increased life expectancy with a simultaneous increased burden of disease. However important registration of disease is in itself, for health policy insight into and evaluation of the existing care are also required, which on the one hand makes it necessary to link disease registration to registration of items of service and on the other hand tries to place disease registration in the perspective of the diseases occurring among the population. The transition from epidemiology to health (services) research is taking shape here and it is in this field that the morbidity registration of the National Survey of Morbidity and Interventions in general practice takes place.


1.2 Backgrounds and objectives

Within (primary) health care in the Netherlands the general practice occupies a key position. On the one hand, via that general practice the greatest number of contacts between population and health care takes place, on the other the general practitioner has the possibilities of making selective use or having selective use made of care that is not directly accessible in the Netherlands. However, at national level there is little good and relevant information available on general practice. As the starting point of study the epidemiological basis of general-practice care in the Netherlands has therefore been taken (van der Velden, 1986; Foets, van der Velden, 1990). The following objectives have been pursued in this (Foets et al., 1986):

1. Obtaining insight on a national scale into the presentation of diseases, complaints and problems in general practice and into the actions taken by the general practitioners as a result of these presented problems.
2. Obtaining insight into the factors that influence the presentation of health problems in general practice and also the varying reactions by the general practitioners to the problems submitted to them.

This basic report is directed towards the first part of the first objective: obtaining insight on a national scale into the presentation of diseases, complaints and problems in general practice. This can be explained by means of Figure 1.1, showing the chain of the professional aid process. In this report attention is directed towards the demand for professional help and the diagnosis that the general practitioner makes thereby.

Figure 1.1 Chain of the professional aid process



The National Survey of morbidity and interventions in general practice (to be referred to further as the National Survey) must be regarded as a study in which a certain sector of health care, in this case the medicine of general practice, is described and analysed. It is thus a typical 'health services research' project, intended above all to acquire information on health care facilities and the use of these; the registration of morbidity is not a goal in itself but an aid to studying the caring process. In this an important position is occupied by the description and explanation of differences between general practitioners, as regards both the morbidity presented to them and interventions.

Morbidity in the National Survey

Morbidity has been registered in general practice, by the general practitioner and his or her practice assistant, by means of a contact registration for three months, whereby reasons for encounter, (differential) diagnosis/working hypothesis and underlying diseases have been recorded. In addition, by means of a health interview survey and a health diary the subjective state of health has been measured in a random sample of the population. Finally, the admission and discharge diagnoses are known of the patients that were admitted and discharged from hospital during the registration period. Because all these data have been collected in the same population and in the same period, this information can be linked at the level of the individual patient.

A morbidity registration as described here has not been performed before in the Netherlands, and it enables us to acquire new insights with regard to the epidemiology of diseases in the Netherlands. However, that does not mean that the most precise incidence/prevalence rates of all disorders in the population can be given. This is due to the fact that problems presented in general practice form a fraction of problems occurring among the population and moreover do not always properly reflect these. For this reason among others, in the health interview survey of the National Survey detailed questions are asked about the state of health, so that a good insight can be obtained into the relation between the health problems in the population and the presentation of problems in general practice: the so-called iceberg phenomenon (van Es, 1983, van de Lisdonk, 1985; Goldberg, Huxley, 1980). This morbidity recorded in the patient survey relates to problems reported by the patient, and not to any diagnoses made by the general practitioner.

This report will pass over this subjective health status of health of the population. Attention is directed towards the morbidity presented to the general practitioner, with as central question:

What is the epidemiology of diseases in general practice in the Netherlands?

On the basis of the episode-oriented morbidity registration of the National Survey a calculation of incidence and prevalence rate of disease in general practice has been performed. The goals of this basic report may be described as follows in this regard:

- presentation of nationally representative data with regard to morbidity in Dutch general practice;
- description of differences both between sections of the population, distinguished by a number of basic characteristics, and between geographical units in morbidity presented to the general practitioner.

The registration of morbidity in the National Survey is episode-oriented, in emulation of the National Morbidity Surveys in the United Kingdom and Dutch morbidity registration projects (Office of Population Censuses and Surveys, 1974a, 1974b and 1984; van Weel et al., 1986; Lamberts, 1986, Lamberts et al., 1987a). An episode-oriented registration has been opted for in the first place on account of the possibility of recording the trend of morbidity over time - whether or not under the influence of diagnostics and therapy. In the second place it is possible by placing episodes side by side to increase insight into the diagnostic and therapeutic action of general practitioners.

The concept disease episode has been further operationalized on behalf of the National Survey, a distinction being made between the concepts 'episode of need' and 'episode of care' (Foets, van der Velden, 1990). On behalf of the morbidity registration in general practice only the concept 'episode of care' is used. This 'episode of care' forms the numerator of the epidemiological fraction. For a description of the way in which morbidity data from the contact registration have been processed into disease episodes see Section 2.3.2. The population data required for calculation of the denominator of the epidemiological fraction have been obtained from the well-described practice populations of the participating practices.

The investigation has been performed among a random non-proportionally stratified sample of 161 established Dutch general practitioners, from 103 general practices, serving a joint practice population of 335,000 patients. They form a 2.5% sample of the Dutch population of general practitioners.

Data collection took place in four registration periods, each of three months, between 01-04-1987 and 31-03-1988. The three months form a balance between required numbers of data per unit of time and the possibilities with respect to the workload of the registering general practitioner and practice assistant and are at the same time a representation of seasons. Computerized processing of morbidity data took place at NIVEL. A 'field worker' who visited the registering general practice once a week formed the important link between the general practice and NIVEL. These field workers helped the GPs with the organization of data collection and checking the completed registration forms. At NIVEL they coded the completed forms and entered them into the computer.

1.3 Outline of the report

In Chapter 1 the background to the morbidity study is presented. In Chapter 2 a justification is given of the way in which the data have been obtained, the processing performed is accounted for and the value of the data is discussed. Successively the realization of the sample (including the weighting), the epidemiological numerator and the epidemiological denominator is explained, the operationalization of the incidence and prevalence is described and the quality of the data is discussed. In Chapter 3 the results, largely in tabular form, are presented.

2. DATA AND METHODS

2.1 Introduction

In this chapter a justification is given of the performance of the National Survey and the processing applied to the data. In Section 2.2 it is explained how a sample of general practitioners was taken so as to obtain data representative of the Netherlands. In Section 2.3 justification of the morbidity registration, or the epidemiological numerator, takes place. The contact registration performed in the participating general practices and the classification system used for the morbidity are discussed. In Section 2.4 determination of the denominator by means of the patient registration performed is described. The denominator has been determined on the basis of data on the patients of all participating general practices. Also discussed are the data collection, the non-response, the representativeness of the data and the applied weighting to the Dutch population. Section 2.5 describes how the incidence and prevalence rates were ultimately calculated. In the last section of this chapter attention is devoted to the quality of the material. A full description of the methodology, the procedure and the measuring instruments of the National Survey is given in the basic report 'Meetinstrumenten en procedures' (Foets and van der Velden, 1990).

2.2 Sample of general practitioners

For the National Survey a random non-proportionally stratified sample of general practitioners was taken. The sample framework was formed by the general practitioner file of NIVEL, which contains all officially registered independently established general practitioners as at 01-01-1985. The sample is meant to give a representative picture of the Dutch general practitioner and his or her practice and should display sufficient variation in GP characteristics, so that differences between general practitioners can be examined with sufficiently large numbers. Use is made of *stratification variables* three stratification variables that are known to be connected with morbidity (Lako et al., 1978, van Sonsbeek, 1987) or aspects of GP action (see for instance Wijkel, 1986). The variables are region, distance from the practice address to the nearest hospital and degree of urbanization.

The region variable is divided into;

- north (Groningen, Friesland and Drenthe);
- south (North Brabant and Limburg);
- centre (remaining provinces).

The distance from practice address to nearest hospital variable has the following breakdown:

- hospital in same place as the practice address;
- hospital outside the place and up to 15 km from the practice;
- hospital outside the place and more than 15 km from the practice.

The degree of urbanization variable has the following four (CBS) categories:

- rural (fewer than 30.000 inhabitants);
- suburban (30 000 to 50.000 inhabitants);
- urban (more than 50.000 inhabitants);
- city (Amsterdam, Rotterdam and The Hague).

The non-proportional nature of the sample is intended to guarantee that all sample cells are equally filled. The intention was to recruit 6-7 registering physicians per sample cell.

The recruitment resulted in a participation of 161 established general practitioners from 103 GP practices. Of these 161 general practitioners, 98 come from the sample; the other established general practitioners are colleagues of sample physicians in the same practice (n=57) and volunteers (n=6). The non-established physicians (other physicians) consist of practice assistant GPs (n=10), locums (n=9) and general practitioners in training (n=13). In total 193 general practitioners and other physicians collaborated in the investigation.

Table 2.1 surveys all the registering physicians, broken down by the stratification variables region, distance from practice address to hospital and degree of urbanization.

Theoretically there are $3 \times 3 \times 4 = 36$ cells. As the table shows, not all cells are applicable. The cell filling is reasonable, with the exception that one cell in the south region has only one participating GP. In respect of reality GPs and other physicians from the north region, rural areas and GPs and other physicians working at a practice address that is more than 15 km from the nearest hospital are deliberately over-represented.

No morbidity has been registered of the contacts with colleagues' patients. Telephone encounters were registered only if these resulted in a referral or prescription. Encounters consisting only of writing out a repeat prescription were also recorded. The feasibility study that preceded the investigation (Foets et al., 1986) had shown that three months' registration for the short-term episodes is in any case enough. With the long-term episodes (in particular chronic diseases) an under-estimation of the prevalence may occur, because in a number of the patient in question will not have been in the GP's practice during those three months. Since telephone encounters (including requests for repeat prescriptions) have also been registered, the under-estimation will be limited.

On behalf of this report use has been made of the following data:

Patient data:

- age;
- sex.

Encounter data:

- date of the encounter;
- nature of the encounter (first encounter, relapse, repeat encounter, periodical check).

Reasons for encounter and diagnosis/working hypothesis:

- reason(s) for encounter;
- probability diagnosis/working hypothesis at the highest possible diagnostic level (Meyboom, 1979);
- underlying disease, provided that this was relevant in the framework of this encounter.

The total registration period amounted to one calendar year (01-04-1987 to 31-03-1988). In the distribution of the general practitioners over the four registration periods maximum allowance has been made for the distribution among the sample cells (see Table 2.2). In this way seasonal influences have been equally divided among the registering general practitioners.

Table 2.2 Registering GPs and other physicians by season

season	registration period	number of registering GPs and other physicians (n = 193)
spring	01-04-1987 - 30-06-1987	45
summer	01-07-1987 - 30-09-1987	49
autumn	01-10-1987 - 31-12-1987	50
winter	01-01-1988 - 31-03-1988	49

Of the 161 established GPs and the remaining 32 physicians, one GP performed the registration for only six weeks. This leads to an under-estimation of the incidence and prevalence. Correcting for the duration of the registration is not reliably possible. This practice has therefore not been included in the analyses of the basic report.

General practitioners and practice assistants have registered data per encounter with a patient. As within a encounter more than one problem can be presented, the number of problems exceeds the number of encounters. Per problem a diagnosis has been made by the general practitioner. This diagnosis is the starting point for calculation of the morbidity.

In the National Survey a total of 387,250 encounters (in which 418,750 problems were presented) have been registered. Of these, 780 encounters (in which problems were presented) are encounters of the general practitioner who registered for only six weeks, while 26,000 encounters with colleagues' patients have been recorded. In this report the results of 360,470 encounters (in which 417,850 problems were presented) have ultimately been included.

2.3.2 From problem to episode

The registration of morbidity in the National Survey is episode-oriented in emulation of the National Morbidity Surveys in the United Kingdom and Dutch morbidity registration projects (Office of Population Censuses and Surveys, 1974a, 1974b and 1984; Van Weel et al., 1986; Lamberts et al., 1987a). According to the WONCA (Classification Committee 1981), an episode is a problem or illness in a patient over the entire period of time from its onset to its resolution. Following this definition in an operational sense, the concept 'episode' is confined here to the phase in which professional help is sought (the 'episode of care'). Within one episode one or more than one encounter can take place. The goal of the episode-oriented registration is to give a more accurate picture of the incidence and the prevalence than is possible on the basis of the individual encounters. Registering at episode level results in a reduction in the number of false-positive and false-negative diagnoses because many 'probability diagnoses' crystallize out into definitive diagnoses. The data registered at encounter level make comparisons of the trend within episodes of the same disease among different patients possible.

Arranging the problems of one patient that are presented in one or more encounters into episodes took place afterwards by combining a number of data: patient code, date of the encounter, nature of the encounter (first encounter, encounter in connection with relapse, repeat encounter, encounter in connection with periodical check), reason for encounter, diagnosis, the underlying disease and the degree of certainty of the diagnosis (measured on 5-point scales). The diagnosis of the last encounter in an episode is adhered to as the diagnosis of the whole episode. These episodes form the starting point for determination of the epidemiological numerator. All constructed episodes have been checked and in the case of doubt about their correctness submitted to the general practitioner concerned. The morbidity data in this report are based on 309,125 episodes (see Section 2.3.3).

2.3.3 Morbidity classification

Within medicine, particularly that of general practice, a number of morbidity classifications are available: namely the ICD-9-CM (International Classification of Diseases, 9th revision, Clinical Modification 1980), the E list of the British Royal College of General Practitioners, compiled by Eimerl (College of General Practitioners, 1959), the ICHPPC-2-Defined (International Classification of Health Problems in Primary Care, 1983), the RFEC or Reason for Encounter Classification (Lamberts et al., 1982 and 1984) and the ICPC (International Classification of Primary Care, Lamberts, Wood, 1987b). All the classification systems

mentioned, except the ICD-9-CM, have been specially designed for use in general practice. The ICD-9-CM is strongly clinically oriented and comprises almost exclusively diagnoses and barely complaints and problems that occur often in general practice. Moreover, the coding in the ICD-9-CM is extensive and complicated for semi-trained coders. The recently developed ICPC, on the other hand, is clearly laid out in components and chapters and offers the possibility of coding diagnoses and complaints. The coding-friendly nature and the comparability with other research were the decisive criteria in determination of a choice of a classification system. (Crombie, van der Zee, 1984; Schellevis et al., 1984a; Schellevis, 1984b; Severien, Van der Zee, 1984). This ultimately led to the choice of the ICPC. A shortcoming in 1987 of the then available version of the ICPC was the limited number of diagnosis-categories. The ICPC was therefore adapted to a version of our own with a finer classification that is compatible with the international version (van der Velden et al., 1989). The refinement that has been applied by NIVEL consists of a third figure that has been added in a number of cases. It is important to remark that with the ICPC not only diagnoses can be classified but also reasons for encounter, preventive activities, diagnostic and therapeutical interventions.

ICPC system

An ICPC code consists of a letter followed by two figures (the alfa-numeric code). The 17 letter codes, which refer to an equal number of ICPC chapters, are based on localization (for instance A = general, B = blood etc.). The figure codes follow a corresponding system per chapter as far as possible. For instance, D26 is the code for fear of cancer of the gastrointestinal tract and R26 the code for fear of cancer of the respiratory organs. Further, the component involved can be derived from the figures:

component	codes
1. symptoms/complaints	1-29
2. diagnostic/preventive services	30-49
3. medication/treatment/therapeutical services	50-59
4. results	60-61
5. administrative services	62
6. referrals and other reasons for encounter	63-69
7. diagnoses/diseases	70-99

The basis used is 316,947 episodes of care. Of these, in 309,125 cases a symptom diagnosis or diagnosis could be made, coded in accordance with component one and seven respectively. Further, in 7822 episodes of care it was not possible to code a diagnosis or a symptom, either because the problem entered was not convertible into a diagnosis, e.g. the code for sport (A98) and localization of face (L22) or because there was no question of a disease or a complaint but of a procedure (e.g. preventive activity: codes terminating in 44, 45 and 49). The morbidity data are therefore based on 309.125 episodes of care coded in accordance with components 1 and 7.

On behalf of the presentation of figures in this basic report a number of ICPC codes have been clustered. The clustering of ICPC codes concerns only the symptom diagnoses (ICPC codes from component one). Criteria used in clustering diagnoses are: the prevalence of the disorder, the frequency of encounters, clinical homogeneity and etiological grounds. Component seven has been left intact. The codes are numbered from 1 to 684. In the basic tables the morbidity figures are given per diagnosis thus distinguished. The diagnoses are arranged

per chapter. In Appendix B a description is given per diagnosis number and both the corresponding NIVEL-ICPC code and the corresponding Oxford ICPC code are listed.

2.3.4 Summary

This section has discussed the epidemiological numerator. On the basis of a random non-proportional stratified sample 193 GPs and other physicians, of whom 161 established GPs, have been recruited. During four three-month periods all encounters of patients with these physicians and practice assistants were registered. All problems presented during these encounters have been classified through the ICPC. The problems presented have been arranged in episodes. The episodes form the basis for determination of the numerator of the incidence and the prevalence per 3 months.

2.4 The epidemiological denominator

2.4.1 Patient registration

On behalf of determination of the epidemiological denominator patient registration has been performed. This patient registration consists of the collection of basic data on all patients of the participating physicians. This makes it possible to determine denominators not only for the total population but also for characteristics like age, sex, highest level of education attained, marital status, form of insurance, country of birth, occupational class, current working situation and living situation. The starting point for the patient registration was the GP's patient records. This registration provided for the majority of the patients the characteristics age, sex and form of insurance. This information has been supplemented with the aid of forms, on which in addition to age and sex a number of socio-demographic data could be entered (Foets et al., 1990). The forms were completed as much as possible during an encounter of the patient with the general practitioner or the practice assistant. Patients who during the registration period had no encounter with the GP or for other reasons had not completed a registration form were sent a form upon completion of the registration period. In the final phase the data were further supplemented by means of reminders, telephone encounter with the patient and health insurance fund lists. The ultimate response for the complete set was 91.2%.

For all patients from the practice population age and sex are known via the GP's records. Supplementary patient data were collected by means of the patient registration. The non-response to the patient registration (having completed the form or not) is 8.7%. Table 2.3 shows the (non-)response by age, sex, stratification variables and season. The differences prove to be small in general. The greatest are the differences in the age variable. A higher non-response is seen in the group below the age of five and above the age of 75. A lower non-response is observed in the 5-14 age group. In addition a relatively high non-response has to be reported in the three big cities. A relatively low non-response may be noted in the north region and in the spring (the first registration period). In the other categories the deviation from the average non-response is less than 10%.

Table 2.3 Response to patient registration (form completed or not) by age and sex, stratification variables and season

	response		non-response	
	n	%	n	%
total	303.248	91,3	29.055	8,7
sex				
male	147.040	92,0	12.691	8,0
female	155.895	90,6	16.190	9,4
age				
0-4 years	19.694	89,4	2.332	10,6
5-14 years	39.530	93,6	2.693	6,4
15-24 years	51.705	90,8	5.261	9,2
25-44 years	97.617	91,5	9.059	8,5
45-64 years	58.941	91,8	5.290	8,2
65-74 years	20.674	90,5	2.162	9,5
75 years and over	15.042	87,1	2.228	12,9
degree of urbanization				
rural	116.718	91,0	11.547	9,0
suburban	121.634	91,6	11.147	8,4
urban	50.281	91,7	4.577	8,3
city	14.615	89,1	1.784	10,9
region				
north	82.355	92,3	6.880	7,7
centre	111.233	91,2	10.684	8,8
south	109.660	90,5	11.491	9,5
distance from hospital				
same place	119.781	91,0	11.891	9,0
1-15 km	112.432	91,9	9.975	8,1
> 15 km	71.035	90,8	7.189	9,2
season				
spring	83.766	93,2	6.142	6,8
summer	69.587	90,2	7.599	9,1
autumn	79.987	90,6	8.332	9,4
winter	69.908	90,9	6.982	9,1

In this report the morbidity is distinguished by the characteristics age, sex, marital status, living situation, form of insurance and country of birth, level of education, current employment status and occupational class. In the selection of these characteristics two criteria played a part:

1. It must be known from the literature that the characteristic in question is connected with morbidity.
2. The categories that are distinguished on the basis of the characteristics in question must be of sufficient size to be able to give reasonable estimates of incidence and prevalence.

In Section 2.4.3 the denominators per characteristic used in the calculation of the incidence and prevalence rates will be accounted for. Before this can be done the weighting factors used will be further explained, since the weighting influences the denominators.

2.4.2 Weighting

This report aims at nationally representative figures on morbidity presented in general practice. In principle that is possible because a national sample of GPs collaborated in the National Survey. Since the sample is stratified a weighting may be necessary to arrive at accurate estimates of incidence and prevalence. May be, because if it proves that there is no connection between morbidity and the stratification characteristics used, weighting is not required. In this section, after an analysis has shown that weighting is necessary, the weighting procedure is explained.

To investigate the latter an analysis has been performed. Per practice the number of episodes per ICPC chapter has been determined and related to the number of patients (number of episodes per 1000 patients). Then variance analyses were performed at practice level with as dependent variables these ratios per ICPC chapter and as independent variables region, degree of urbanization and distance from the practice to the hospital (Table 2.4). First region, degree of urbanization and distance from the practice to the hospital were separately related to the number of episodes per 1000 patients. It becomes clear that for each of the three variables there are significant correlations with the number of episodes per 1000 patients. This need not necessarily lead to the conclusion that therefore weighting has to be applied for all three of the variables. For it may be that certain connections cease to apply if the other stratification variables are checked for.

Table 2.4 Significance level of differences in average number of episodes per 1000 patients per ICPC chapter between practices broken down by stratification variables (n=102)

	degree of urbanization		region		distance from hospital	
A. General and unspecified	.89	.66	.18	.28	.11	.06
B. Blood	.29	.54	<u>.01</u>	<u>.02</u>	.09	.18
D. Digestive	.53	.54	.20	.22	.07	.05
F. Eye	<u>.02</u>	<u>.03</u>	.30	.77	.86	.65
H. Ear	.24	.32	.38	.45	.73	.85
K. Circulatory	.09	.47	.05	.13	<u>.04</u>	.17
L. Musculoskeletal	<u>.03</u>	<u>.02</u>	<u>.00</u>	<u>.01</u>	.06	<u>.01</u>
N. Neurological	<u>.01</u>	.21	<u>.05</u>	.26	<u>.01</u>	.10
P. Psychological	.72	.63	.38	.31	<u>.00</u>	<u>.00</u>
R. Respiratory	.53	.57	<u>.00</u>	<u>.01</u>	.26	.12
S. Skin	.60	.63	<u>.03</u>	<u>.04</u>	.45	.35
T. Endocrine/metabolic/nutritional	.34	.56	.25	.30	.09	.17
U. Urology	.96	.94	.19	.27	<u>.04</u>	<u>.05</u>
W. Pregnancy/family planning	.10	.21	<u>.03</u>	<u>.03</u>	.31	.47
X. Female genital system	<u>.04</u>	.25	.77	.89	<u>.02</u>	.18
Y. Male genital system	.21	.20	.57	.35	.11	.19
Z. Social problems	<u>.04</u>	.29	.17	.10	<u>.01</u>	.10

p1 = significance of difference between averages

p2 = significance of difference between averages after check for other stratification variables

Consequently analyses have also been performed in which the three stratification variables together have been related to the prevalence rates. In this way it is investigated to what extent a connection exists between a stratification variable and the prevalence rates if a check is made for the other two variables. The table shows that in that case too statistically significant differences in prevalence between practices distinguished by the three stratification variables continue to exist. The conclusion that must be drawn from this is that weighting has to be applied to degree of urbanization, region and distance from the practice to the hospital.

To obtain nationally representative data on morbidity in general practice a second correction is necessary. For the question is to what extent the practice populations of the participating general practitioners reflect the practice populations of all Dutch general practitioners together (assuming that the practice populations of all Dutch GPs together are equivalent to the Dutch population). Through non-response and sample fluctuation it is possible that the population investigated differs in composition from the Dutch population. If the age/sex distribution of the population investigated is compared with the Dutch population the differences are admittedly slight (Table 2.5), but if within a given stratification cell the age/sex distribution of the practice populations of the collaborating physicians is compared with the actual distribution in the population there are considerable differences (this will be made evident later). To arrive at nationally representative figures a weighting by age and sex will therefore have to be applied per stratification cell.

Table 2.5 Practice population National Survey and Dutch population (01-01-1987) by age and sex (absolute and in percentages)

age category and sex		National Survey		Netherlands	
0-4 years	M	11.234	3,4	449.475	3,1
	F	10.712	3,2	430.920	2,9
5-14 years	M	21.322	6,4	953.309	6,5
	F	20.735	6,2	911.655	6,2
15-24 years	M	27.486	8,3	1.274.319	8,7
	F	29.362	8,8	1.221.351	8,4
25-44 years	M	51.960	15,6	2.353.807	16,1
	F	54.643	16,4	2.245.867	15,4
45-64 years	M	31.519	9,5	1.470.552	10,1
	V	32.691	9,8	1.498.584	10,3
65-74 years	M	9.885	3,0	459.306	3,1
	V	12.943	3,9	590.408	4,0
75 years and over	M	6.291	1,9	262.619	1,8
	F	10.965	3,3	491.585	3,4
unknown		555	0,2	-	-
total		332.303	100,0	14.613.757	100,0

One final correction that possibly has to be made is concerned with the fact that registration has taken place in four successive three-month periods. This was done to incorporate the seasonal factor. If, however, the populations investigated are not evenly divided among the seasons, there is a chance that a seasonal effect will nevertheless occur. Now the collaborating GPs are spread as evenly as possible over the four registration periods, but nevertheless differences may occur in the ultimate size of the practice populations per period. Differences in practice size may be at the bottom of this, but also differences in effects of the other weightings between the registration periods. This should be corrected for too.

The weighting procedure consists of three steps. First weighting factors are calculated at patient level on the basis of the stratification characteristics degree of urbanization and region and the variables age and sex. Then weighting factors are calculated at the level of general practices for the stratification variable distance from the practice address to the hospital. Finally, weighting factors per season are determined on the level of the total population.

The first step starts from the following variables:

- region (north, centre, south);
- degree of urbanization (rural, suburban, urban, three big cities);
- sex (male, female);
- age (up to 4 years inclusive, 5 to 14 years inclusive, 15 to 24 years inclusive, 25 to 44 years inclusive, 45 to 64 years inclusive, 65 to 74 years inclusive, 75 years and older).

The variable distance from practice address to the hospital has been left out of consideration, because no population data are available for this to which back-weighting can be applied. For this variable weighting takes place in the second step at the level of GPs. On the basis of the four variables mentioned a 3 x 4 x 2 x 7 table has been constructed. Per cell the number of patients occurring in the patient registration has been related to the number in the Dutch population (based on CBS data of 1 January 1987). The following formula has been used for this:

$$\text{weighting factor in cell } x = \frac{\text{number in population in cell } x}{\text{number in patient registration in cell } x}$$

The weighting factors calculated in this way are included in Table 2.6. The size of the weighting factors varies from about 10 in the suburban areas in the north of the Netherlands to over 100 in the big cities. The variation among the age-sex categories is much less great, although this differs per age category. In some sample cells the practice populations from the study form a good reflection of the age-sex structure (rural north and suburban north and south). In other sample cells that is much less the case (urban centre and city centre are the peaks). In general the practice populations in urban areas and in the centre of the Netherlands form a less accurate reflection of the total population in the various sample cells than in other areas. By use of the weighting factors given in the table these deviations are corrected for.

Table 2.6 Weighting factors step 1

	rural			suburban			urban			big city
	north	centre	south	north	centre	south	north	centre	south	centre
males 0-4 jr	26,6	56,5	21,3	9,4	83,4	22,8	27,1	58,8	58,4	96,4
females 0-4 jr	27,1	57,7	21,2	9,4	83,3	23,4	25,2	53,9	55,4	103,1
males 5-14 jr	30,0	62,7	24,2	10,7	104,6	27,8	31,4	60,4	69,1	140,8
females 5-14 jr	29,2	64,0	23,2	10,4	100,8	28,1	32,9	57,8	66,6	118,7
males 15-24 jr	28,3	59,1	24,4	10,3	100,8	28,6	31,5	79,1	75,2	127,6
females 15-24 jr	26,6	56,9	22,2	9,2	93,2	25,2	21,7	68,7	63,0	92,8
males 25-44 jr	28,1	64,5	24,0	10,1	91,1	28,3	23,3	70,0	72,0	97,2
females 25-44 jr	26,5	60,8	21,7	9,4	84,5	27,0	17,9	59,7	61,0	80,8
males 45-64 jr	28,0	59,2	24,2	10,4	100,4	30,9	24,5	78,6	72,6	125,6
females 45-64 jr	27,2	58,0	23,2	10,3	97,5	30,7	21,8	78,7	70,7	116,7
males 65-74 jr	27,5	54,0	24,9	10,1	92,5	29,4	20,3	98,4	68,2	112,6
females 65-74 jr	26,0	50,2	22,4	10,6	90,6	29,7	18,4	98,1	61,9	101,7
males 75 jr e.o.	25,8	46,0	27,6	10,1	79,2	26,8	19,4	97,5	54,7	112,6
females 75 jr e.o.	27,7	48,2	24,3	12,2	79,0	26,4	19,1	109,5	56,1	111,9

The second step is the weighting by distance from practice address to the hospital. Because the national distribution of the population among the various categories by distance from practice address to the hospital is unknown, the basis taken here is the national distribution of general practitioners among the various categories. Per stratification cell the number of physicians that collaborated in the National Survey has been related to the number in the Dutch GP population. The ratio thus obtained has been related to the same ratio in cells defined by the variables degree of urbanization and region. In formula:

$$W_{ijk} = \frac{N_{ijk}}{n_{ijk}} / \frac{\sum N_{ij}}{\sum n_{ij}}$$

- Here: W_{ijk} : weighting factor per stratification cell ijk
 N_{ijk} : number of GPs in the Netherlands per cell ijk
 n_{ijk} : numbers of GPs in Study per cell ijk
 $\sum N_{ij}$: number of GPs in cell ij in the Netherlands
 $\sum n_{ij}$: number of GPs in cell ij in Study
 ijk : subscript of degree of urbanization, region and distance from practice address to the hospital

The results of the calculation appear in Table 2.7. A sample calculation may make matters clear. The weighting factor for the north/rural/hospital cell in GP's location is equal to the reciprocal of the sample fraction in this cell (71/6) divided by the reciprocal of the sample fraction in the cell defined by only degree of urbanization and region (369/18) and thus equal to 0.577. In other words back-weighting to the situation in a degree of urbanization/region cell in the GP population takes place.

Table 2.7 Calculation of the weighting factors distance from practice address to the hospital by degree of urbanization and region

region	urbanization	total	practice address	< 15 km	> 15 km
north	rural pop.	369	71	204	94
	N.S.	18	6	6	6
	factor		0,577	1,659	0,764
north	suburban pop.	185	71	99	15
	N.S.	23	6	11	6
	factor		1,471	1,119	0,311
north	urban pop.	101	101	-	-
	N.S.	7	7	-	-
	factor		1,000	-	-
centre	rural pop.	1003	254	564	185
	N.S.	19	3	8	8
	factor		1,604	1,335	0,438
centre	suburban pop.	1244	247	885	112
	N.S.	16	3	10	3
	factor		1,059	1,138	0,480
centre	urban pop.	903	830	73	-
	N.S.	73	7	4	-
	factor		1,444	0,222	-
centre	3 cities' pop.	742	742	-	-
	N.S.	10	10	-	-
	factor		1,000	-	-
south	rural pop.	390	58	275	57
	N.S.	23	8	8	7
	factor		0,428	2,027	0,480
south	suburban pop.	548	203	300	45
	N.S.	23	5	12	6
	factor		1,704	1,049	0,315
south	urban pop.	341	315	26	-
	N.S.	5	4	1	-
	factor		1,155	0,381	-

The last step is a correction for seasonal influences. After the two preceding steps had been performed it was investigated what the distribution of the patients over the four registration periods was. Then a correction was applied in such a way that the distribution over the four periods became the same (Table 2.8).

Table 2.8 Calculation of correction factors per registration period

	frequency after performance of steps 1 and 2 in %	desired frequency in %	correction factor
period 1	28,0	25,0	0,894
period 2	22,7	25,0	1,100
period 3	31,1	25,0	0,804
period 4	18,2	25,0	1,373

2.4.3 Description of denominators per patient characteristic

In this report the morbidity rates are broken down by patient characteristics such as age, sex, marital status, level of education, form of insurance etc. The subdivisions made are accounted for in this section. At the same time both the unweighted and the weighted denominator is given per population category for which morbidity rates are presented. The size of the unweighted denominator is directly connected with the reliability of the figure presented. According as the denominator is lower the unreliability margins are higher. With the aid of the weighted denominators the morbidity rates in the report can be converted into estimates of the absolute number of prevalent and incident cases as are presented in the Netherlands in general practice.

The age/sex distribution before and after weighting is shown in Table 2.9. By way of comparison the age/sex distribution of the Dutch population at the time of the National Survey is placed alongside it. We see that the age distribution after weighting corresponds almost perfectly to that of the Dutch population, which is logical, since after all the former has been weighted back to the latter. We see the lowest unweighted denominator among the males of 75 years and over. Having regard to this low absolute number wide reliability intervals must be adhered to for this category of the population.

Table 2.9 Age/sex distribution of practice population before and after weighting compared with the Netherlands (CBS, 1-1-1987)

age/sex category		before weighting		after weighting		Netherlands
		absolute	%	absolute	%	%
0 - 4 yr.	M	11.243	3,4	452.080	3,1	3,1
0 - 4 yr.	F	10.712	3,2	436.193	3,0	2,9
5 - 14 yr.	M	21.322	6,4	951.358	6,5	6,5
5 - 14 yr.	F	20.735	6,2	921.702	6,3	6,2
15 - 24 yr.	M	27.486	8,3	1.264.573	8,6	8,7
15 - 24 yr.	F	29.362	8,8	1.230.203	8,4	8,4
25 - 44 yr.	M	51.960	15,6	2.366.499	16,1	16,1
25 - 44 yr.	F	54.643	16,4	2.266.671	15,5	15,4
45 - 64 yr.	M	31.519	9,5	1.463.485	10,0	10,1
45 - 64 yr.	F	32.691	9,8	1.495.778	10,2	10,3
65 - 74 yr.	M	9.885	3,0	459.549	3,1	3,1
65 - 74 yr.	F	12.943	3,9	596.071	4,1	4,0
75 jr. e.o.	M	6.291	1,9	267.445	1,8	1,8
75 jr. e.o.	F	10.965	3,3	503.428	3,4	3,4
unknown		555	0,2	-	-	-

A second breakdown that has been made is by the formal marital status as indicated by the patients on the patient registration form (Table 2.10). This breakdown must be carefully distinguished from the living situation (whether or not living together), which is discussed further on. Morbidity rates are given for the single, the married, the divorced and the widowed. The categories 'marital status unknown' and 'marital status child' are left out of consideration. By 'child' males to 17 years inclusive and females to 15 years inclusive are designated. The latter has been done so as not to distort the figures for the single group too strongly.

Table 2.10 The investigated practice population by marital status (before and after weighting)

marital status	before weighting		after weighting	
	abs.	%	abs.	%
child	74.751	22,5	3.235.323	22,0
single	59.863	18,0	2.722.266	18,6
married	141.286	42,5	6.039.627	41,2
divorced	7.898	2,4	415.709	2,8
widowed	15.449	4,6	711.666	4,8
onbekend	33.056	9,9	1.550.444	10,6
totaal	332.303	100,0	14.657.035	100,0

The respondents were asked to indicate on the patient registration form the highest level of education attained. Because persons up to 17 years inclusive in general have not yet attained their highest level of education, this category has been left out of consideration, as also the category unknown (Table 2.11).

Table 2.11 The investigated practice population by highest level of education attained (before and after weighting)

level of education before weighting	after weighting		abs.	%
	abs.	%		
age < 18 years	80.668	24,3	3.464.359	23,6
no/primary education	56.432	17,0	2.413.693	16,4
secondary education	120.735	36,3	5.242.901	35,7
higher professional/university	22.564	6,8	1.128.248	7,7
unknown	51.904	15,6	2.425.834	16,5
total	332.303	100,0	14.675.035	100,0

Morbidity rates will be given for publicly insured versus privately insured persons (Table 2.12). The group that stated that they were not insured or unknown are left out of consideration in this report on account of the small size.

Table 2.12 The investigated practice population by type of insurance (before and after weighting)

form of insurance	before weighting		after weighting	
	abs.	%	abs.	%
public	204.245	61,5	8.715.637	59,4
private	104.415	31,4	4.904.773	33,4
no insurance	592	0,2	25.158	0,2
unknown	23.051	6,9	1.029.467	7,0
total	332.303	100,0	14.675.035	100,0

Origin has been determined on the basis of the country of birth stated by the respondent (Table 2.1.3). A distinction has been made between the Netherlands, Turkey/Morocco, other Western and other non-Western. (Non-)Western country of birth has not been broken down further by country of origin, although this was asked for, because the numbers of respondents per country were too small.

Table 2.13 The investigated practice population by country of birth (before and after weighting)

origin	before weighting		after weighting	
	abs.	%	abs.	%
Netherlands	259.633	78,1	11.121.963	75,8
Turkey/Morocco	2.684	0,8	174.376	1,2
other Western	4.143	1,2	201.026	1,4
other non-Western	6.350	1,9	358.556	2,4
unknown	59.463	17,9	2.819.114	19,2
total	332.303	100,0	14.675.035	100,0

The respondents were asked to indicate their (last practised) occupation. If they did not have one the occupation of the parents (in the case of children living at home) or the partner was included. Use was made of the Occupational Classification 1984 of the CBS. By means of a recoding key this was given an EGP code (class scheme of Erikson, Goldthorpe and Portocarero, 1979, 1982, 1983), running from one to ten. The ten-class distribution has been converted into a three-class distribution. Class 1 represents headwork, class 2 self-employed and farmers and class 3 manual work. It must be noted that the occupation remains unknown for a considerable percentage. In that case the question was looked at that relates to the situation if no occupation is performed (table 2.14).

Table 2.14 The investigated practice population by occupational class (before and after weighting)

	before weighting		after weighting	
	abs.	%	abs.	%
occupational class				
headwork: high + medium	60.623	18,2	2.832.566	19,3
headwork: low	57.922	17,4	2.635.720	18,0
self-employed + farmers	15.980	4,8	617.961	4,2
manual work: high + medium	57.502	17,3	2.328.244	15,9
manual work: low	58.343	17,6	2.416.768	16,5
non-classifiable	1.583	0,5	66.678	0,5
other				
age < 16 years	14.899	4,5	727.443	5,0
child/student	4.496	1,4	232.574	1,6
national service	315	0,1	13.497	0,1
housewife/husband	6.413	1,9	298.262	2,0
unemployed	1.894	0,6	98.017	0,7
recipient of disablement benefit	1.105	0,3	51.400	0,4
early retirement/pension	6.512	2,0	294936	2,0
not working/no reason	41	0,0	1.622	0,0
other unknown	44.675	13,4	2.059.347	14,0
total	332.303	100,0	14.675.035	100,0

Morbidity rates are given for working versus non-working persons (broken down into jobless and recipients of disablement benefit). This item has been derived from the question about the occupation and a question concerning the situation if one does not practise that occupation (table 2.15).

Table 2.15 The investigated practice population by current employment status (before and after weighting)

current work situation	before weighting		after weighting	
	abs.	%	abs.	%
working	102.988	31,0	4.627.885	31,5
unemployed	7.008	2,1	329.856	2,2
recipient of disablement benefit	8.141	2,4	355.686	2,4
< 16 years	69.421	20,9	2.981.654	20,3
child/student	22.573	6,8	994.139	6,8
national service	878	0,3	38.144	0,3
housewife/husband	44.678	13,4	1.820.216	12,4
early retirement/pension	26.152	7,9	1.209.605	8,2
not working/no reason	159	0,0	6.946	0,0
unknown	50.305	15,1	2.310.903	15,7
total	332.303	100,0	14.675.035	100,0

This refers to the form of cohabitation, that is to say whether people live alone, with a partner and children or with a partner without children. Persons living in institutions and communes have been left out of consideration on account of their small numbers. In the tables single persons (including one-parent families) are distinguished from the others (family, (married) couples). To render possible a meaningful comparison of single persons with those living in a family relationship, children below the age of 18 have been left out of consideration (table 2.16).

Table 2.16 The investigated practice population by living situation (before and after weighting)

living situation	before weighting		after weighting	
	abs.	%	abs.	%
age < 18 years	80.668	24,3	3.464.359	23,6
single	29.290	8,8	1.512.680	10,3
(married) couple/family	175.860	52,9	7.540.813	51,4
one-parent family	9.445	2,8	444.693	3,0
unknown	37.040	11,1	1.712.491	11,7
total	332.303	100,0	14.675.035	100,0

2.4.4 Description of denominators per practice characteristic

In this section it is indicated how the practice population of the collaborating general practitioners before and after weighting is distributed by region, degree of urbanization and distance from the practice address to the nearest hospital. Because the location of the general practice and not that of the patient has been taken here as point of departure, these data are always known. The denominators used are shown in Table 2.17.

Table 2.17 Practice population by degree of urbanization, region and distance from the hospital and season (before and after weighting)

	before weighting		after weighting	
	abs.	%	abs.	%
degree of urbanization				
rural	128.265	38,6	4.747.055	32,4
suburban	132.781	40,0	5.031.531	34,3
urban	54.858	16,5	3.248.938	22,1
city	16.399	4,9	1.647.511	11,2
region				
north	89.235	26,9	1.569.723	10,7
centre	121.917	36,7	9.490.706	64,7
south	121.151	36,5	3.614.606	24,6
distance from the hospital				
same place	131.672	39,6	7.226.622	49,2
1-15 km	122.407	36,8	6.127.396	41,8
> 15 km	78.224	23,5	1.321.017	9,0
season				
spring	89.908	27,1	3.668.758	25,0
summer	77.186	23,2	3.668.759	25,0
autumn	88.319	26,6	3.668.759	25,0
winter	76.890	23,1	3.668.759	25,0
total	332.303	100,0	14.675.035	100,0

2.4.5 Summary

The age and sex are known of practically all patients. Supplementary data are known on 91.2% of the patients through the patient registration. The differences between responding and non-responding patients regarding age, sex, stratification variables and season are slight.

On behalf of the pursuit of nationally representative figures on morbidity in general practice it has been investigated whether weighting is necessary as regards the stratification variables age and sex and season. As significant differences exist in the prevalence between practices in distinguishing the three stratification variables, weighting should be performed for degree of urbanization, region and distance from practice address to hospital. In view of the fact that within a stratification cell the age/sex distribution in the practice population differs from the Dutch population, a correction has also been made for age and sex. In order to eliminate the seasonal factor (differences in population investigated and differences in effects of other weightings), as the final step a correction has been performed in such a way that the distribution of patients over the four registration periods became the same.

In the description of denominators per patient characteristic it proves that among the responding patients the non-response is the highest with the socio-economic variables (level of education, occupational class and current employment status), with the exception of the form of insurance variable. As regards distribution by age and sex category the practice population of the National Survey may be described as representative of the Dutch population.

The total of the known practice populations of the participating physicians forms the epidemiological denominator. For breakdowns by characteristics of the patient such as age, sex and other characteristics the denominator is formed by the number of patients for whom the characteristic in question is known. For breakdowns by geographic characteristics the location of the general practice has been used as a starting point. These data are always known.

2.5 Incidence and prevalence

In this report both the cumulative incidence and the prevalence rate per disorder are indicated. The cumulative incidence and the prevalence rate may be defined as follows (Mausner, 1974):

Cumulative incidence: the number of new cases of disease that occur in the course of a period of time, divided by the number of persons present in the population at the beginning of the period of time considered. In the National Survey this period is three months.

Period-prevalence rate: the number of existing cases of disease in a given period divided by the average population. The period prevalence is built up from the prevalence at a given point in time, plus the number of new cases and relapses in a given period. In the National Survey this period is three months.

The operationalization of the incidence and prevalence in the National Survey may be described as follows. The incidence has been determined by counting those episodes in which the general practitioner indicated in the first encounter of the episode in question that a first encounter for the problem in question or a encounter in connection with a relapse for the problem was involved. The prevalence has been determined by counting the total number of episodes in which identical episodes were also counted once only per patient. As the length of registration was three months, the *incidence and prevalence* have been determined *per three months*.

Conversion of the numbers into figures on an annual basis is possible by multiplying the incidence by a factor of four, by which a cumulative incidence per year is obtained. Conversion of the prevalence of in particular chronic disorders into figures on an annual basis is particularly complicated and can be performed reliably only with further information. As a result of this, the prevalence cannot be roughly multiplied by four. In the case of an acute disorder with a high incidence and a high prevalence the prevalence per 3 months can be multiplied by four to obtain the prevalence per year. As regards a chronic disorder with a low incidence and a high prevalence, the prevalence per 3 months is equal to the prevalence per year. For the other disorders the multiplication factor is in between these.

An important fact is that calculation of perinatal mortality and mortality has been based on the same denominators as for the other disorders. The numbers are therefore not directly comparable with the perinatal mortality and mortality determined in other investigations.

The figures relate to problems presented to the general practitioners in three months. This implies that the figures are not a reflection of the occurrence of diseases and problems among the population, but only insofar as these have been presented to the general practiti-

oner. This has consequences notably for the prevalences of those disorders with which the patient is acquainted but which form a limited reason to consult the general practitioner (within the registration period). As an example the problem of deafness: this is presented to the general practitioner only if there is a '(re-)activation' of the problem of deafness (deterioration, replacement of hearing aid).

In the event of a differential diagnosis, only the diagnosis first reported has been included in the calculation.

Part of the morbidity is presented solely by a request for a repeat prescription. On account of the extent of this service, the repeat prescription was registered by the practice practice assistant only in the first registration period during the whole period and in the other periods only during half of the period. Introduction of a correction factor is therefore necessary. Without correction considerable differences occur in the amount of morbidity between on the one hand period one and on the other periods two, three and four. In period one there is then more morbidity. This relates to disorders for which repeat prescriptions are regularly written out, such as sleep disorders, depressive feelings, hypertension and oral anticonception. Therefore, a correction has been applied to the morbidity data from the second, third and fourth registration periods insofar as these have been generated solely by repeat prescriptions registered by the practice practice assistant. Per practice a correction factor has been determined by taking as a basis the repeat prescriptions recorded in the half period. The number of episodes that consisted exclusively of encounters in which solely repeat prescriptions were registered has been doubled. This has been done after it had emerged from the material of the first period that there was no difference in number of episodes with repeat prescriptions between the first and the second half of the registration period.

2.6 Validity and reliability of the data

2.6.1 Validity

The validity of the morbidity data means the degree in which a diagnosis made by the general practitioner (which is included in the data base coded in accordance with the ICPC) corresponds to the patient's actual disease. This is also referred to as the correctness of the diagnosis.

The validity of GP diagnoses must be seen in the light of the GP's working methods. It is inherent in these methods that many diagnoses are made as working hypotheses (Van der Velden, 1983) and are formulated at a low diagnostic level. According as the episode continues, a more differentiated picture forms and it is possible to make a medical diagnosis that can be fitted into a medico-nosological system.


In the episodic representation of morbidity the 'effect' of the course of time on the validity of the diagnosis is implicitly incorporated. Basing morbidity data on disease episodes increases the validity of those data.

In order to gain insight into the validity of the diagnoses made, 30 'paper vignettes' were submitted to all participating GPs with the request to make a diagnosis of each case. In

addition to an indication of the reliability, the uniformity in the naming of syndromes by the GPs is also to be regarded as an indication of the validity of the diagnoses, despite the absence of a 'gold standard'. Per paper vignette the response varied from 85% to 100%. On account of interpretation problems one paper vignette was not involved in the analysis.

In the following figures the percentage of agreement per case is shown.

Figure 2.2 Percentage of agreement by GPs on paper vignettes



In the case of 27 of the 29 patients the percentage of agreement is 75 or higher; for 21 of the 29 this is 90% or higher. The percentage of agreement is low for the case with dizziness or iridocyclitis as diagnosis (41% and 15% respectively).

In the coding of the morbidity data by the field worker interpretation of data was inevitable, which threatens the validity. In the discussion of the results of the uniformity tests among the field workers (see 2.6.2) attention was devoted to this in the sense of training. The field workers were moreover instructed to consult the registering general practitioner if matters were not clear.

The validity of the morbidity data can be increased by the application of diagnostic criteria by the registering GPs. For this training and verification meetings ought to take place. The scale on which the National Survey was held made this unfeasible.

2.6.2 Reliability

The reliability of the morbidity data means the reproducibility and the accuracy. Various procedures and measurements have been performed to track down systematic errors.

1. The diagnoses were coded 'centrally' by a (para)-medically trained field worker. After training in the use of the ICPC uniformity tests were regularly performed with paper vignettes. Through these tests insight was obtained into the uniformity of the coding. Further instruction was directed towards those points where uniformity left something to be desired. On account of the training aspect deliberate use was made of 'problem cases'. Table 2.18 contains the percentages of agreement of the field workers in coding of diagnoses. The average percentage of agreement is 80 for 'problem' codes of diagnoses at the level of encounters. Through the episode construction used this percentage - viewed theoretically - becomes higher.

Table 2.18 Percentages of agreement of field workers in the classification of diagnoses

checkpoint	number of field workers	average agreement (%)
1	38	69
2	32	81
3	35	77
4	38	83
5	34	80
6	32	88

In a qualitative sense acute infectious diseases and the ICPC chapter W (Pregnancy, delivery and lying-in) proved to contain the highest percentages of agreement. The agreement was the lowest for diagnoses from Component one (Symptoms and complaints) of the ICPC chapters A (General and unspecified), P (Psychological) and Z (Social problems).

2. The construction of disease episodes was performed by the field worker. In problem cases the GP in question was consulted. The construction of encounter diagnoses into disease episodes was fully checked by four physicians; if necessary the episode diagnosis was changed.
3. Data entry was checked by comparing the data input of a sample of 2% of the encounter registration forms with the data on the forms. In 1.1% of the checked forms an error was found that usually concerned one variable.

Incompatibility concerning the combinations diagnosis - age and diagnosis - sex was detected by a computer program. (van Weel, 1986). The results appear in Table 2.19.

Table 2.19 Incompatibility concerning the combinations diagnosis -age and diagnosis - sex in percentages of the encounters and episodes respectively

	% of encounters	% of episodes
diagnosis - age	0,05	
diagnosis - male sex		0,69
diagnosis - female sex		1,30

3. RESULTS

3.1. Introduction

In this chapter a description is given of the morbidity in Dutch general practice in a predominantly statistical sense. In Sections 3.2 and 3.3 the incidence and the prevalence, weighted to the Dutch population, are presented of the morbidity per ICPC chapter with breakdowns to practice and patient characteristics respectively. In Section 3.4 an overview is given of the twenty most frequent incident and prevalent episodes among men and women respectively. The most detailed incidence and prevalence rates are given in tables in the appendices. In these tables the morbidity has been broken down into 684 diagnoses.

3.2 Morbidity per ICPC chapter by practice characteristics and season

In this section differences between practices in presented morbidity per ICPC chapter are given. After a general overview the size of the variation in incidence and prevalence between practices is described per chapter. Next variations between practices distinguished by geographical characteristics of the location of the practice address are discussed. These geographical characteristics, degree of urbanization, region and distance from the hospital, at the same time form the stratification variables in taking the sample of GPs (see Section 2.2). Finally, the morbidity has been broken down by season. The characteristic season is in the National Survey in fact a practice characteristic, because in the four successive registration periods different practices registered each time.

In Table 3.1 a survey is given of the incidence and prevalence of the episodes per 1000 persons ICPC chapter, weighted to the Dutch population.

Table 3.1 Incidence and prevalence per ICPC chapter after weighting per 3 months absolute and per 1000 persons

ICPC chapter	estimate of incidence absolute	estimate of incidence per 1000 patients	estimate of prevalence absolute	estimate of prevalence per 1000 patients
A. general and unspecified	493.414	33,6	912.700	62,2
B. blood	59.102	4,0	186.168	12,7
D. digestive	541.975	36,9	984.354	67,1
F. eye	229.665	15,7	402.880	27,5
H. ear	373.997	25,5	469.108	32,0
K. circulatory	247.018	16,8	1.684.603	114,8
L. musculoskeletal	1.189.847	81,1	1.837.213	125,2
N. neurological	147.371	10,0	437.327	29,8
P. psychological	331.254	22,6	1.339.119	91,3
R. respiratory	1.240.696	84,5	1.910.728	130,2
S. skin	951.210	64,8	1.465.894	99,9
T. endocrine/metabolic/nutritional	71.511	4,9	435.777	29,7
U. urology	211.993	14,4	335.348	22,9
W. pregnancy & family planning	187.693	12,8	743.911	50,7
X. female genital system	237.435	16,2	438.982	29,9
Y. male genital system	51.256	3,5	80.373	5,5
Z. social problems	103.445	7,0	288.896	19,7
total	6.668.882	454,3	13.953.381	950,8

The incidence per three months is 454.3 per 1000 persons. Low incidences concern the ICPC chapters male genital system, blood, endocrine/metabolic/nutritional and social problems. High incidences, on the other hand, relate to the ICPC chapters respiratory, musculoskeletal and skin. The total prevalence is 950.8 per 1000 persons, despite the nature of the disorder. At ICPC chapter level the prevalence per 3 months varies from 5.5 per 1000 (disorders of male genitals) to 130.2 per 1000 (respiratory). Chapters with high prevalences are respiratory, musculoskeletal, circulatory, skin and psychological. Chapters with low prevalences are male genital system, blood, social problems, urology and eye.

There are differences in incidences and prevalences of ICPC chapters between the practices. This may be the result of actual differences in morbidity presented in general practice. However, differences between GPs in the designation of health problems may also play a part. This can be illustrated by means of the following table, in which the average incidence and prevalence per practice and the corresponding variation coefficient are stated. The variation coefficient is a measure of the size of the standard deviation expressed as a percentage of the average. Unlike the standard deviations, the variation coefficients are comparable with each other.

Table 3.2 Average incidence and variation coefficient and average prevalence and variation coefficient per ICPC chapter (n=102 practices)

ICPC chapter	average incidence	variation coefficient of incidence	average prevalence	variation coefficient of prevalence
A. general and unspecified	34,3	31,1	64,7	32,5
B. blood	3,9	44,6	13,4	41,3
D. digestive	35,9	25,4	66,9	22,0
F. eye	15,9	27,7	28,4	27,4
H. ear	25,9	26,6	32,4	24,4
K. circulatory	16,7	37,3	123,3	36,3
L. musculoskeletal	81,0	22,8	125,9	19,6
N. neurological	10,2	37,6	32,0	36,0
M. psychological	23,0	43,2	95,6	32,0
R. respiratory	85,2	31,6	133,1	27,5
S. skin	65,5	23,9	101,5	20,9
T. endocrine/metabolic/nutritional	4,8	41,9	31,2	34,1
U. urology	14,4	34,7	23,5	31,5
W. pregnancy & family planning	13,0	48,6	49,9	27,3
X. female genital system	16,2	35,1	30,1	29,1
Y. male genital system	3,6	37,5	5,7	28,6
Z. social problems	7,1	58,9	19,3	62,4

The highest variation coefficient of the incidence and the prevalence may be found in the ICPC chapter social problems. This is pre-eminently a chapter in which designation effects may occur. In general the variation coefficients of the incidence are higher than those of the prevalence. The high variation in the pregnancy chapter is connected with the fact whether the GP is obstetrically active or not. The variation coefficients are further high for the ICPC chapters blood, psychological and endocrine/metabolic/nutritional. For the prevalence the variation coefficients are high for the ICPC chapters circulatory, neurological, endocrine/metabolic/nutritional and general and unspecified. The lowest variation coefficients of both incidence and prevalence relate to the ICPC chapters musculoskeletal, skin and digestive.

In the following table the incidence per ICPC chapter is shown by the degree of urbanization of the municipality in which the registering general practice is located.

Table 3.3 Incidence per ICPC chapter by degree of urbanization after weighting per 3 months per 1000 persons

ICPC chapter	degree of urbanization			
	rural	suburban	urban	city
A. general and unspecified	30,9	35,9	34,0	33,7
B. blood	3,8	3,5	5,1	4,1
D. digestive	32,8	37,3	43,0	35,8
F. eye	15,8	16,0	16,7	12,3
H. ear	23,7	25,8	28,6	23,4
K. circulatory	15,0	15,4	20,9	18,5
L. musculoskeletal	75,4	85,4	89,9	67,0
N. neurological	10,3	10,2	10,0	8,8
M. psychological	16,7	24,3	26,9	25,5
R. respiratory	78,5	84,5	95,4	80,9
S. skin	61,3	67,4	67,9	61,2
T. endocrine/metabolic/nutritional	4,3	4,6	5,4	6,2
U. urology	13,1	13,9	16,9	15,2
W. pregnancy/family planning	15,8	10,8	12,6	10,5
X. female genital system	14,2	16,8	18,3	15,8
Y. male genital system	3,1	3,3	3,8	4,6
Z. social problems	5,8	7,0	7,9	9,2
total	420,5	462,1	503,3	432,7

The total incidence is the highest in towns with more than 50 000 inhabitants, excluding the three cities ('urban'), followed by urbanized rural municipalities ('suburban'), the three cities Amsterdam, Rotterdam and The Hague ('city') and finally the rural municipalities ('rural'). For the ICPC chapters endocrine/metabolic/nutritional, male genital system and social problems the incidence increases with the increase in the degree of urbanization. Further, the relatively low incidences of the ICPC chapters general and unspecified, psychological and respiratory in rural areas and of the ICPC chapters eye and musculoskeletal in the city are striking. Conversely, in urban areas the incidence is high of the ICPC chapters digestive, circulatory, respiratory and blood.

In Table 3.4 the prevalence per ICPC chapter is shown by degree of urbanization.

Table 3.4 Prevalence per ICPC chapter by degree of urbanization after weighting per 3 months per 1000 persons

ICPC chapter	degree of urbanization			
	rural	suburban	urban	city
A. general and unspecified	59,8	63,3	61,0	68,0
B. blood	15,2	10,1	13,4	12,0
D. digestive	65,7	64,8	74,3	63,7
F. eye	29,0	27,0	28,5	22,4
H. ear	30,2	32,0	35,4	30,1
K. circulatory	129,7	102,9	116,5	105,0
L. musculoskeletal	120,3	126,4	137,3	111,9
N. neurological	35,5	27,5	28,7	22,6
M. psychological	89,7	87,8	97,4	94,2
R. respiratory	129,1	125,9	141,6	124,1
S. skin	99,9	101,4	101,7	91,7
T. endocrine/metabolic/nutritional	32,7	25,8	30,4	31,4
U. urology	23,3	21,8	24,3	21,9
W. pregnancy/family planning	54,8	47,9	51,8	45,3
X. female genital system	27,8	29,4	33,4	30,5
Y. male genital system	5,0	5,2	6,0	6,5
Z. social problems	14,7	19,5	23,8	26,3
total	962,4	918,7	1.005,5	907,5

It is striking that the prevalence is the highest in urban areas, followed by rural areas, suburban areas and the cities (Table 3.4). As regards the prevalence of the ICPC chapters male genital system and social problems, there is an increase in the prevalence with an increase in the degree of urbanization. In rural areas the high prevalences of blood, circulatory, neurological and pregnancy/family planning are striking. Relatively high prevalences in urban areas relate to the ICPC chapters digestive, ear, musculoskeletal, psychological and respiratory. In the big cities the prevalence of eye disorders is relatively low.

In Tables 3.5 and 3.6 incidence and prevalence respectively by region are presented per ICPC chapter.

Table 3.5 Incidence per ICPC chapter by region after weighting per 3 months per 1000 persons

ICPC chapter	region north	centre	south
A. general and unspecified	34,1	32,5	36,3
B. blood	4,1	4,0	4,0
D. digestive	31,9	36,1	41,2
F. eye	15,7	15,4	16,4
H. ear	25,2	25,3	26,1
K. circulatory	15,9	16,9	17,2
L. musculoskeletal	75,4	77,5	93,0
N. neurological	9,7	9,8	10,9
M. psychological	20,4	23,1	22,2
R. respiratory	77,3	80,1	99,4
S. skin	61,1	63,1	70,8
T. endocrine/metabolic/nutritional	4,1	5,0	4,8
U. urology	14,7	14,4	14,3
W. pregnancy/family planning	12,6	13,4	11,3
X. female genital system	16,0	16,2	16,3
Y. male genital system	3,6	3,4	3,5
Z. social problems	6,7	6,8	7,8
total	428,5	443,0	495,5

The total incidence increases going from the north region to the centre and is the highest in south. In south region the incidence is above all relatively high with the ICPC chapters musculoskeletal, respiratory, skin and social problems.

Table 3.6 Prevalence per ICPC chapter by region after weighting per 3 months per 1000 persons

ICPC chapter	region north	centre	south
A. general and unspecified	68,6	59,6	66,2
B. blood	15,6	12,6	11,7
D. digestive	64,8	65,9	71,1
F. eye	28,1	26,9	28,6
H. ear	31,4	31,6	33,1
K. circulatory	139,0	109,4	118,5
L. musculoskeletal	118,5	120,2	141,3
N. neurological	34,7	28,6	30,7
M. psychological	97,8	92,8	84,5
R. respiratory	129,0	123,9	147,3
S. skin	95,9	97,4	108,2
T. endocrine/metabolic/nutritional	36,0	29,4	27,7
U. urology	26,1	22,3	22,9
W. pregnancy/family planning	47,8	50,9	51,5
X. female genital system	30,3	30,3	28,6
Y. male genital system	5,9	5,4	5,6
Z. social problems	16,7	19,1	22,6
total	986,2	926,2	1.002,2

The total prevalence is the highest in south region, followed by north region and centre region. North region is striking with a high prevalence of the chapters circulatory and psychological. In south region the prevalences of the chapters musculoskeletal, digestive and respiratory are relatively high.

In Tables 3.7 and 3.8 the incidence and prevalence respectively by distance from the practice address to the nearest hospital are presented per ICPC chapter.

Table 3.7 Incidence per ICPC chapter by the distance from the practice address to the nearest hospital after weighting per 3 months per 1000 persons

ICPC chapter	distance from practice address to hospital		
	same place	1-15 km	> 15 km
A. general and unspecified	36,8	30,5	30,8
B. blood	4,7	3,4	3,6
D. digestive	39,7	34,5	33,3
F. eye	15,5	16,0	14,9
H. ear	26,3	24,7	24,6
K. circulatory	19,4	14,4	13,8
L. musculoskeletal	83,1	79,3	78,3
N. neurological	10,1	10,1	9,6
P. psychological	25,5	19,7	20,2
R. respiratory	89,1	80,8	76,8
S. skin	66,2	63,5	63,6
T. endocrine/metabolic/nutritional	5,5	4,1	5,0
U. urology	16,0	13,0	12,8
W. pregnancy	12,7	12,6	14,4
X. female genital system	17,0	15,5	15,1
Y. male genital system	3,8	3,2	3,3
Z. social problems	8,5	5,6	5,9
total	479,9	430,9	426,0

If incidences are compared on distance from the practice address to the hospital it is striking that if the practice address is situated in the same place as the hospital the incidences are higher than when the hospital is located outside the place and further away from the practice address. This is in particular noticeable for the ICPC chapters general and unspecified, blood, digestive, circulatory, psychological, respiratory and social problems.

In Table 3.8 the prevalence per ICPC chapter by distance from the practice address to the nearest hospital is presented.

Table 3.8 Prevalence per ICPC chapter by distance from practice address to nearest hospital after weighting per 3 months per 1000 persons

ICPC chapter	distance from practice address to hospital		
	same place	1-15 km	> 15 km
A. general and unspecified	67,2	55,1	67,8
B. blood	13,7	11,1	14,5
D. digestive	71,7	61,3	68,5
F. eye	27,4	26,9	30,2
H. ear	32,6	31,1	32,7
K. circulatory	117,5	106,5	138,7
L. musculoskeletal	132,1	116,9	125,9
N. neurological	28,8	29,6	36,0
P. psychological	99,3	79,0	104,0
R. respiratory	137,1	123,0	125,8
S. skin	100,4	98,3	104,3
T. endocrine/metabolic/nutritional	31,2	26,7	34,9
U. urology	24,4	21,0	23,0
W. pregnancy/family planning	50,4	50,3	54,0
X. female genital system	31,7	27,6	30,8
Y. male genital system	6,0	4,9	5,5
Z. social problems	24,4	14,9	15,9
total	995,9	884,2	1012,5

In practices located within a distance of 15 kilometres from the nearest hospital (but not in the same place) the total prevalence is lower than further from the hospital and in practices where the hospital is located in the same place. Far from the hospital the prevalence of disorders from the circulatory chapter is high. Social problems are relatively common in practices with a hospital in the same place.

In the following tables the incidence and prevalence by season are presented per ICPC chapter. As already mentioned, in the National Survey season is a practice characteristic because per season registration has been performed by different practices.

Table 3.9 Incidence per ICPC chapter by season after weighting per 3 months per 1000 persons

ICPC chapter	season			
	spring	summer	autumn	winter
A. general and unspecified	36,0	31,1	34,6	32,8
B. blood	4,1	4,0	4,1	3,9
D. digestive	38,7	34,0	37,3	37,8
F. eye	16,4	14,5	16,6	15,1
H. ear	26,1	23,9	25,6	26,3
K. circulatory	15,7	20,9	16,4	14,3
L. musculoskeletal	77,8	74,9	86,1	85,5
N. neurological	9,2	9,1	11,5	10,4
P. psychological	21,4	20,8	25,4	22,7
R. respiratory	81,6	60,3	101,3	95,0
S. skin	66,2	64,2	69,3	59,5
T. endocrine/metabolic/nutritional	4,1	5,4	5,0	5,0
U. urology	13,3	13,4	16,8	14,3
W. pregnancy/family planning	12,2	14,1	12,5	12,4
X. female genital system	16,4	15,7	17,6	15,0
Y. male genital system	3,7	3,0	3,9	3,4
Z. social problems	6,4	5,6	7,8	8,4
total	449,3	414,9	491,8	461,8

The total incidence (Table 3.9) is the highest in the autumn and the lowest in the summer. In the summer the incidences are high of the ICPC chapters circulatory and pregnancy/family planning; in the autumn of psychological, respiratory and urology; and in the winter likewise of respiratory and of social problems. The ICPC chapters digestive, ear, musculoskeletal, respiratory and social problems have low incidences in the summer.

Table 3.10 Prevalence per ICPC chapter by season after weighting per 3 months per 1000 persons

ICPC chapter	season			
	spring	summer	autumn	winter
A. general and unspecified	62,6	63,3	61,3	61,6
B. blood	12,4	14,7	11,8	11,8
D. digestive	65,8	64,1	68,9	69,6
F. eye	26,8	26,4	29,1	27,5
H. ear	32,7	30,2	31,6	33,4
K. circulatory	97,4	130,1	122,9	108,8
L. musculoskeletal	121,5	117,0	132,1	130,1
N. neurological	26,9	30,0	31,8	30,6
P. psychological	82,2	92,4	97,0	93,4
R. respiratory	123,8	99,3	150,6	147,1
S. skin	101,2	98,1	103,1	97,2
T. endocrine/metabolic/nutritional	25,0	31,0	29,3	33,5
U. urology	19,5	22,9	25,5	23,4
W. pregnancy/family planning	52,2	47,1	50,6	52,9
X. female genital system	29,6	29,5	31,6	29,0
Y. male genital system	5,8	4,7	6,3	5,1
Z. social problems	18,4	15,8	22,5	22,1
total	903,8	916,6	1006,0	977,1

In the autumn and the winter the prevalences for the chapters respiratory and social problems are considerably higher than in the other seasons (Table 3.10). Psychological disorders have the lowest prevalence in the spring. In the summer and autumn the prevalence of disorders of the circulatory are higher than in the spring and winter.

3.3 Morbidity per ICPC chapter by patient characteristics

In this section a presentation follows of the morbidity per ICPC chapter by characteristics of the patient. The patient characteristics are age, sex, form of insurance, marital status, living situation, country of birth, level of education, current employment status and occupational class. The breakdowns by age, sex and form of insurance may also be found in the appendices, but then for the detailed diagnosis clusters.

Table 3.11 Incidence and prevalence per ICPC chapter by sex after weighting per 3 months per 1000 persons

ICPC chapter	incidence		prevalence	
	male	female	male	female
A. general and unspecified	27,2	39,8	47,6	76,3
B. blood	2,8	5,2	7,7	17,6
D. digestive	32,8	40,9	58,3	75,5
F. eye	15,4	15,9	24,9	29,9
H. ear	25,5	25,5	31,4	32,5
K. circulatory	13,5	20,1	95,8	133,2
L. musculoskeletal	80,1	82,0	114,6	135,5
N. neurological	8,3	11,8	23,0	36,4
P. psychological	17,3	27,7	63,7	118,0
R. respiratory	80,8	88,1	129,6	130,8
S. skin	62,5	67,0	93,2	106,4
T. endocrine/metabolic/nutritional	3,6	6,1	20,5	38,6
U. urology	6,6	22,1	12,4	33,0
W. pregnancy/family planning	-	25,0	-	98,9
X. female genital system	-	31,7	-	58,6
Y. male genital system	7,1	-	11,1	-
Z. social problems	5,6	8,5	14,0	25,2
total	389,1	517,4	747,8	1146,4

Table 3.11 gives incidence and prevalence rates by sex. In general both the incidences and the prevalences are higher for women than for men. This applies to a stronger extent to the ICPC chapters general and unspecified, blood, digestive, circulatory, psychological, endocrine/metabolic/nutritional and urology. In part this will be connected with the fact that women become older on average and therefore the elderly group is over-represented among women compared to men.

In Table 3.12 it can be seen that after childhood the incidences and the prevalences strongly increase with age. In this respect there are marked differences between the chapters. Disorders of the ear and of the respiratory occur very frequently, relatively speaking, among young children. The disorders that increase the most strongly in incidence and prevalence with increasing age are circulatory disorders.

Table 3.12 Incidence and prevalence per ICPC chapter by age categories after weighting per 3 months per 1000 persons

ICPC chapter	incidence						
	0-4	5-14	15-24	25-44	45-64	65-74	75 and over
A. general and unspecified	87,9	31,8	32,4	28,5	26,3	30,7	42,7
B. blood	4,6	4,7	4,7	4,4	2,5	2,9	5,1
D. digestive	63,5	37,5	31,8	32,9	35,6	39,1	47,7
F. eye	18,6	13,1	16,7	15,5	15,1	16,4	17,1
H. ear	74,2	32,2	15,0	20,6	22,2	26,6	27,4
K. circulatory	1,2	1,1	4,4	10,4	26,2	48,5	72,6
L. musculoskeletal	18,3	46,7	79,2	90,6	102,8	88,7	91,7
N. neurological	5,4	4,7	7,8	10,5	12,8	14,7	15,6
P. psychological	5,1	7,1	21,0	28,4	27,9	24,6	27,0
R. respiratory	220,8	89,6	79,6	77,2	68,2	67,2	61,4
S. skin	87,2	77,2	78,9	61,2	51,8	49,2	56,6
T. endocrine/metabolic/nutritional	4,0	2,7	3,8	4,5	5,9	8,2	8,7
U. urology	9,3	8,1	11,6	12,6	16,7	25,0	32,6
W. pregnancy/family planning	-	0,6	24,4	26,6	0,6	-	-
X. female genital system	1,3	3,6	23,0	24,6	16,1	5,8	6,1
Y. male genital system	3,4	3,4	2,9	4,2	3,0	3,3	3,7
Z. social problems	2,6	2,3	5,2	7,7	10,1	8,8	11,6
total	607,4	366,4	442,4	460,4	443,8	459,7	527,6

ICPC chapter	prevalence						
	0-4	5-14	15-24	25-44	45-64	65-74	75 and over
A. general and unspecified	112,4	43,4	48,4	50,0	60,9	87,7	137,8
B. blood	9,6	8,0	8,6	11,4	11,7	19,6	42,6
D. digestive	77,8	44,9	41,7	52,6	80,9	111,9	163,6
F. eye	25,0	22,2	22,6	22,1	28,4	44,4	64,3
H. ear	84,1	40,7	17,7	25,0	29,2	36,1	43,1
K. circulatory	3,5	3,0	9,9	37,2	191,0	429,4	596,9
L. musculoskeletal	26,7	57,0	99,3	124,1	171,6	184,1	236,0
N. neurological	9,7	9,7	15,8	26,3	43,4	55,7	80,2
P. psychological	11,8	14,9	39,9	82,2	142,6	194,5	250,7
R. respiratory	268,5	127,2	108,2	107,9	121,4	166,7	167,1
S. skin	116,4	100,5	120,1	91,0	88,3	94,0	120,0
T. endocrine/metabolic/nutritional	8,5	6,8	10,4	18,1	45,3	86,5	104,3
U. urology	11,4	9,7	14,2	16,3	26,0	48,4	88,5
W. pregnancy/family planning	-	1,6	118,7	90,2	8,5	-	-
X. female genital system	2,0	4,8	33,4	39,1	38,6	24,1	30,8
Y. male genital system	4,8	4,3	3,6	5,3	5,1	9,5	11,7
Z. social problems	6,3	4,9	12,7	21,0	27,4	31,5	40,2
total	778,5	503,6	725,2	819,8	1120,3	1624,1	2177,8

The combination age/sex in relation to the morbidity presented in general practice is shown in Fig. 3.1. Each chapter will be briefly considered.

general and unspecified

Disorders from the chapter 'General and unspecified' occur (both incidence and prevalence) relatively often among 0-4-year-olds. Further we see an increase in prevalence with increasing age. Per age category disorders from this chapter occur more often among women than among men (with the exception of the 0-4 age group). Disorders that are decisive for the level of the prevalence in this chapter are childhood diseases, unspecified viral diseases, allergies and above all the working hypothesis 'no disease'.

blood, blood-forming organs, lymphatics and spleen

Disorders from the chapter 'blood, blood-forming organs, lymphatics and spleen' (both incidence and prevalence) are relatively infrequent. The prevalences in this chapter increase with age and are somewhat higher for women than for men. Decisive diagnoses are anemias (iron-deficiency anemia, pernicious/other deficiency anemia and not further defined anemias) and to a much smaller extent enlarged lymph glands.


digestive

Digestive disorders occur relatively often among young children (notably incidence) and among the elderly. Among the elderly disorders from this chapter occur more frequently among women than among men. The most common diagnoses are in decreasing order of prevalence other gastrointestinal infections, irritable bowel syndrome, constipation, disorders of stomach function/gastritis, stomach ache, worms/parasites and generalized abdominal pain/cramps.

eye

Disorders of the eye occur equally in all age categories. This applies above all to the incidence; prevalence occurs somewhat more often with increasing age. Women over the age of 65 visit the GP with these disorders more frequently than men from this age category. The most common diagnosis in this chapter is allergic conjunctivitis/not classifiable elsewhere, followed by refractive errors, complaints about vision, other symptoms/complaints of eye.

Figure 3.1 Incidence and prevalence of ICPC chapters by age and sex after weighting per 3 months per 1000 persons



ear

Disorders of the ear often occur above all with children. Beyond childhood there is not much difference between the age groups and between the sexes. Incidence and prevalence rates are not far apart. Common disorders of the ear are ear wax, acute otitis media/myringitis and otitis externa.

circulatory

Incidence and prevalence rates of the circulatory are far apart. Moreover, they are by far the most age-specific disorders. Above the age of 45 the prevalences of the disorders from this chapter rise sharply. For women the prevalences are even somewhat higher than for the men, above all for the group of 75 years and over. The decisive diagnosis is above all uncomplicated hypertension, which takes up more than a third of the episodes from this chapter. Other common (but much less frequent) disorders are angina pectoris, heart failure, old infarction and edema of the extremities.









musculoskeletal

While the incidence rate falls with age, disorders of the musculoskeletal system are more frequently seen in general practice according as age increases. There is a clear difference between men and women here. Whereas the incidence for men rises before the 65th year and then falls, the prevalences increase up to the 45-64 age group and then remain fairly constant above this age group. For women the incidence rises strongly up to 65 years and then falls somewhat, but the prevalences still clearly continue to increase above the age of 65. The most common individual disorder from this chapter is myalgia/fibrositis. Further, sprains and strains often occur (ankle, knee, shoulder/arm, wrist/hand, foot/toes). and to a smaller extent lumbago, tendinitis/synovitis, shoulder syndrome (incl. bursitis) and symptoms/complaints relating to the back.

neurological

Disorders from the 'neurological' chapter are of relatively infrequent occurrence. This may be noted most strongly with the incidence: this increases slightly with age. The prevalence is higher with older people than with young ones and higher with women than with men. Headache/facial pain, migraine, other diseases of the peripheral neurological and epilepsy are the most common disorders.

Figure 3.1 Continued



psychological

In the 'psychological problems' chapter too incidence and prevalence differ strongly. The incidence occurs above all among adult females and remains constant, whereas with adult males a decline sets in at a greater age. As regards the prevalences, these become higher with increasing age and, as in the case of the musculoskeletal, the increase is stronger among women than among men. The symptom diagnoses 'feeling nervous' and 'disturbances of sleep/insomnia' are the most decisive for this chapter. Other very common diagnoses are depressive disorder and neurasthenia/surmenage.

respiratory

Disorders of the respiratory are relatively common among small children and old males. Because infections determine this chapter, incidence and prevalence rates are relatively close together. Upper respiratory tract infection and acute bronchitis have a high incidence. Less but nevertheless also frequent in occurrence are acute tonsillitis/adenoiditis, influenza, allergic rhinitis and cough. Acute bronchi(oli)tis, sinusitis, chronic bronchitis and asthma have high prevalences. Lung cancer is the most common cancer among males.


skin

Disorders from the skin chapter are very uniformly distributed among the age/sex groups. Both the incidence and the prevalence are somewhat higher among young children, young adults (15 to 24 years) and the elderly. The most common diagnosis in this chapter is eczema (contact dermatitis and above all other eczema). Dermatophytosis/mycosis, warts, acne vulgaris and lacerations/cuts also occur frequently.

endocrine, metabolic/nutritional

Low incidence and high prevalence among the elderly characterize disorders from this chapter, which occur more among women than among men. By far the most common diagnosis is diabetes mellitus. The only other common occurrence is obesity.

Figure 3.1 Continued



urology

Disorders from this chapter occur as regards both incidence and prevalence more among the elderly than among the young and more among women than among men. It is above all the diagnosis cystitis/other urinary infection, non-venereal that is decisive, and to a lesser degree incontinence, urine.

pregnancy, childbearing, family planning

Diagnoses from the pregnancy chapter occurred above all, as was to be expected, among women in the 15-44 age group. The most common episodes relate to the (continuous) use of the contraceptive pill, as a result of which the prevalence rates are so high, followed at a distance by the diagnosis 'pregnancy confirmed', which strongly determines the incidence rates.









genital system

The male and female genital system chapters are shown in one figure. Disorders of the female genital system occur above all in the middle age groups (15 to 64 years), whereby the chronic nature drives incidence and prevalence apart. Disorders of the male genital system occur very infrequently, and then somewhat more often among older men. The acute nature of these disorders brings incidence and prevalence together. Among women urogenital candidiasis and symptoms/complaints relating to menopause are the most common diagnoses. Among men there are no clearly outstanding diagnoses.

social problems


Social problems are relatively little diagnosed in general practice. Women present these problems somewhat more often than men, and a slight rise may be noted with increasing age. There are no clearly outstanding diagnoses. Problems as the result of loss of a partner and problems between partners are the most common occurrences.

Figure 3.1 Continued



In Figure 3.2 the incidence and prevalence per ICPC chapter are shown by form of insurance.

Figure 3.2 Incidence and prevalence per ICPC chapter by form of insurance after weighting per 3 months per 1000 persons



With regard to the breakdown by form of insurance it can be seen that the incidence and prevalence among those who are publicly insured is in all cases higher than with private patients, with the exception of disorders of the male genitals, while the incidence and prevalence of disorders of the ear is the same for both groups.

Morbidity by marital status

In Table 3.13 the incidence and prevalence are broken down by marital status.

The fact that the marital status is strongly related to age finds expression in the difference in the size of in particular the prevalences and the distribution among the chapters.

Table 3.13 Incidence and prevalence per ICPC chapter by marital status after weighting per 3 months per 1000 persons


ICPC chapter	incidence			
	single	married	divorced	widowed
A. general and unspecified	31,0	28,3	33,3	40,9
B. blood	4,5	3,6	5,6	4,3
D. digestive	32,2	35,1	43,3	46,4
F. eye	14,8	15,5	13,8	18,9
H. ear	18,8	22,7	20,6	24,1
K. circulatory	11,1	21,5	27,5	53,8
L. musculoskeletal	79,2	95,7	109,6	105,0
N. neurological	9,3	12,1	12,2	16,1
P. psychological	25,7	25,2	52,1	33,4
R. respiratory	73,0	73,7	83,4	71,8
S. skin	72,6	55,0	68,2	58,6
T. endocrine/metabolic/nutritional	4,1	5,3	8,7	8,8
U. urology	12,9	15,8	17,9	33,2
W. pregnancy/family planning	15,0	18,5	7,5	0,4
X. female genital system	22,0	18,8	35,1	9,5
Y. male genital system	3,4	3,7	3,7	1,0
Z. social problems	6,4	7,7	15,7	15,4
total	436,0	458,2	558,2	541,6

ICPC chapter	prevalence			
	single	married	divorced	widowed
A. general and unspecified	53,6	59,1	74,1	124,0
B. blood	10,1	12,9	16,1	34,8
D. digestive	53,0	70,6	85,8	142,9
F. eye	23,5	27,1	29,4	57,7
H. ear	22,9	29,0	27,3	35,9
K. circulatory	53,6	150,4	136,4	498,7
L. musculoskeletal	110,0	149,7	173,4	244,0
N. neurological	24,9	36,0	37,5	71,0
P. psychological	75,9	106,3	200,5	256,2
R. respiratory	105,0	122,9	138,0	157,3
S. skin	111,3	88,9	107,3	112,2
T. endocrine/metabolic/nutritional	18,6	34,9	42,2	111,6
U. urology	18,2	24,5	24,7	72,8
W. pregnancy/family planning	93,1	56,3	46,6	4,3
X. female genital system	34,3	35,7	64,9	39,4
Y. male genital system	4,4	6,3	6,8	3,6
Z. social problems	15,6	20,0	51,0	62,3
total	828,0	1030,6	1262,0	2028,7

Morbidity by living situation

In Figure 3.3 the incidence and prevalence are presented per ICPC chapter by the living situation.

Figure 3.3 Incidence and prevalence per ICPC chapter by living situation after weighting per 3 months per 1000 persons




Whereas the incidences for many chapters display no great differences between single persons and (married) couples-families, the prevalences for all chapters with the exception of the pregnancy chapter are higher for single persons. With regard to the psychological, circulatory and endocrine/metabolic/nutritional chapters the differences are the greatest. The age structure (single persons are on average older) is particularly responsible for this.

Morbidity by country of birth

In Fig. 3.4 the incidence and prevalence by country of birth are given.

Figure 3.4 Incidence and prevalence per ICPC chapter by country of birth after weighting per 3 months per 1000 persons




The incidences and the prevalences vary for people born in the Netherlands and outside the Netherlands. Disorders from the circulatory and psychological chapters occur more frequently (for the first time) among persons born in the Netherlands. The digestive, respiratory and social problems chapters occur more frequently (for the first time) among persons born outside the Netherlands.

Morbidity by level of education

In Figure 3.5 the incidence and prevalence by highest level of education attained are presented.

Figure 3.5 Incidence and prevalence per ICPC chapter by level of education, after weighting per 3 months per 1000 persons



With the exception of the pregnancy/family planning and male and female genital system chapters both the incidence and the prevalence decline and the level of education rises. The difference between those with a secondary and a higher education is less striking than the difference between those with a primary and a secondary education. The circulatory and psychological chapters display the greatest differences.

3.4 The twenty most common incident and prevalent diagnoses

Table 3.14 The twenty highest incident and prevalent diagnoses after weighting per 3 months per 1000 persons

incidence diagnosis		prevalence diagnosis		
1	upper respiratory tract infection	30,8	1 uncomplicated hypertension	44,9
2	myalgia/fibrositis	14,4	2 contraceptive pill	35,2
3	cystitis/other urinary infection non-venereal	11,6	3 upper respiratory tract infection	33,5
4	no disease	10,8	4 feeling anxious/nervous	21,0
5	acute bronchi(oli)tis	9,7	5 disturbances of sleep	19,7
6	cough	9,1	6 myalgia/fibrositis	19,2
7	sinusitis	8,7	7 no disease	18,4
8	ear wax	8,3	8 other eczema	15,8
9	other eczema	7,4	9 acute bronchi(oli)tis	13,0
10	acute otitis media/myringitis	6,7	10 cystitis/other urinary infection non-venereal	12,9
11	other infection of digestive	6,6	11 diabetes mellitus	12,0
12	lumbago	6,6	12 sinusitis	10,7
13	influenza	6,2	13 chronic bronchitis	10,5
14	acute tonsillitis/adenoiditis	6,0	14 asthma	10,2
15	other viral diseases NOS/NEC	5,9	15 cough	9,4
16	dermatophytosis/-mycosis	5,8	16 ear wax	9,3
17	allergic conjunctivitis NEC	5,4	17 lumbago	8,8
18	warts	5,0	18 dermatophytosis/-mycosis	8,4
19	tendinitis/synovitis	4,3	19 depressive disorder	8,3
20	irritable bowel syndrome	4,0	20 angina pectoris	8,0

Table 3.14 presents the twenty highest incident and prevalent diagnoses.

Upper respiratory tract infection is in first place in the incidence, but in the prevalence is preceded by episodes in connection with the use of contraception and uncomplicated hypertension. In the interpretation of this table it must be borne in mind that the classification of diagnoses used in the basic tables in Appendix B was used for it. In this classification, for instance, all sprains/strains are broken down by localization. However, as one group they would certainly have ended up in the top ten.

Table 3.15 The twenty highest incident diagnoses among males and females after weighting per 3 months per 1000 persons

males diagnosis		females diagnosis	
1 upper respiratory tract infection	29,2	1 upper respiratory tract infection	32,3
2 myalgia/fibrositis	13,3	2 cystitis/other urinary infection non-venereal	19,3
3 acute bronchi(oli)tis	9,8	3 myalgia/fibrositis	15,5
4 ear wax	8,4	4 no disease	13,2
5 no disease	8,3	5 cough	10,9
6 lumbago	7,7	6 sinusitis	9,9
7 sinusitis	7,5	7 acute bronchi(oli)tis	9,6
8 cough	7,3	8 other eczema	8,3
9 other infection of the digestive	6,6	9 ear wax	8,2
10 acute otitis media/myringitis	6,6	10 acute otitis media/myringitis	6,7
11 other eczema	6,5	11 other infection of the digestive	6,6
12 dermatophytosis/-mycosis	6,4	12 viral infection unspecified NEC	6,3
13 influenza	6,0	13 influenza	6,3
14 acute tonsillitis/adenoiditis	5,8	14 pregnancy confirmed	6,2
15 viral infection unspecified/NEC	5,4	15 acute tonsillitis/adenoiditis	6,2
16 laceration/cut	5,1	16 allergic conjunctivitis NEC	6,0
17 allergic conjunctivitis NEC	4,8	17 irritable bowel syndrome	5,8
18 warts	4,5	18 contraceptive pill	5,7
19 tendinitis/synovitis	4,5	19 warts	5,6
20 sprains/strains/injuries musculoskeletal	4,1	20 lumbago	5,4

Fifteen of the twenty diagnoses that appear in the top twenty for men also appear in the top twenty of most incident diagnoses for women (Table 3.15). Among the women dermatophytosis/mycosis, lacerations/cuts, tendinitis/synovitis and other sprains/strains/injuries with regard to the musculoskeletal do not occur. Among men cystitis/other urinary infections, pregnancy confirmed, irritable bowel syndrome, contraceptive pill and feeling anxious/nervous do not occur. For both men and women infection of the upper respiratory tract is the most common occurrence.

Table 3.16 The twenty highest prevalent diagnoses for males and females after weighting per 3 months per 1000 persons

males diagnosis		females diagnosis			
1	uncomplicated hypertension	32,9	1	contraceptive pill	68,7
2	upper respiratory tract infection	31,3	2	uncomplicated hypertension	56,5
3	myalgia/fibrositis	16,8	3	upper respiratory tract infection	35,6
4	other eczema	13,6	4	feeling anxious/nervous	28,4
5	no disease	13,6	5	disturbances of sleep	27,8
6	acute bronchi(oli)tis	13,3	6	no disease	23,1
7	feeling anxious/nervous	13,3	7	myalgia/fibrositis	21,5
8	chronic bronchitis	12,8	8	cystitis/other urinary infection NEC	21,3
9	disturbances of sleep	11,4	9	other eczema	17,9
10	asthma	10,6	10	diabetes mellitus	14,6
11	lumbago	10,0	11	acute bronchi(oli)tis	12,8
12	diabetes mellitus	9,3	12	sinusitis	12,1
13	sinusitis	9,3	13	depressive disorder	11,9
14	ear wax	9,2	14	cough	11,1
15	dermatophytosis/-mycosis	9,2	15	asthma	9,7
16	angina pectoris	7,8	16	ear wax	9,3
17	cough	7,6	17	pregnancy confirmed	9,3
18	old myocardial infarction	6,9	18	irritable bowel syndrome	8,9
19	other infection of digestive	6,9	19	obesity	8,8
20	acute otitis media/myringitis	6,8	20	chronic bronchitis	8,2

In the top twenty of prevalent disorders among men (Table 3.16) the diagnosis uncomplicated hypertension has by far the highest prevalence (32.9 per 1000), followed by an infection of the upper respiratory tract (31.3 per 1000). Psychological disorders (feeling anxious/nervous and disturbances of sleep), disorders of the respiratory (acute bronchi(oli)tis, chronic bronchitis, asthma, sinusitis), disorders of the musculoskeletal (myalgia/fibrositis and lumbago), diabetes mellitus and eczema all have a prevalence around 10 per 1000. The appearance of 'no disease' in fourth place of prevalent disorders is striking.

The top twenty of prevalent disorders among women is headed by the contraceptive pill (in particular repeat prescriptions). Uncomplicated hypertension scores high (and higher than among the men) with a prevalence of 56.5 per 1000. Psychological disorders (feeling anxious/nervous, disturbance of sleep) and no disease have prevalences around 15 per 1000. It is striking that in the top ten among women only one disorder of the respiratory (infection of the upper respiratory tract) occurs, while with men four disorders of the respiratory occur. For women too 'no disease' is represented in the top ten.

LITERATURE

- Bonte JTP. Het CBS als gegevensbron. In: Epidemiologie en Gezondheidsbeleid). Maas PJ van der, Hofman A, Dekker E. (red.) Samson Stafleu, Alphen aan den Rijn, 1989. (The Central Bureau of Statistics as a source of data).
- CBS. Beroepenclassificatie 1984. Voorburg: Centraal Bureau voor de Statistiek, 1985. (Occupational classification 1984. Voorburg: Central Bureau of Statistics).
- Classification Committee WONCA. An international glossary for primary care. Huisarts Wet. 1981, 24:342-6.
- College of General Practitioners. A classification of disease. J Coll Gen Pract 1959; 2: 140 - 159.
- Crombie DL, Zee J van der. The feasibility of a national general-practice morbidity and intervention survey in the Netherlands. Report of a symposium. Utrecht: Institute of the Dutch General Practitioners, 1984).
- Epidemiologie en informatica. Discussienota. Ontwerp-Kerndocument Gezondheidsbeleid. Ministerie van WVC Rijswijk, 1989. (Epidemiology and Informatics. Working paper. Department of welfare, Public Health and Culture, Rijswijk, 1989).
- Erikson R, Goldthorpe JH, Portocarero L. Intergenerational class mobility in three west european countries: England, France & Sweden. British Journal of Sociology 1979; 10: 415-41.
- Erikson R, Goldthorpe JH, Portocarero L. Social fluidity in industrial nations: England, France & Sweden. British Journal of Sociology 1982; 33: 1-34.
- Erikson R, Goldthorpe JH, Portocarero L. Intergenerational class mobility and the convergence thesis: England, France & Sweden. British Journal of Sociology 1983; 34: 303-43.
- Es JC van. Patiënt en huisarts. Een leerboek huisartsgeneeskunde. Utrecht: Bohn, Scheltema & Bolkema, 1983. (Patient and General Practitioner. A textbook of medical science of general practice).
- Foets M, van der Velden J, van der Zee J. Morbidity and interventions in general practice. A cross-national survey. Study-design. NIVEL. Utrecht, 1986.
- Foets M, van der Velden J. Basisrapport: Meetinstrumenten en Procedures. Nationale Studie naar ziekten en verrichtingen in de huisartspraktijk). Utrecht, NIVEL, 1990. (Basic report: Measuring instruments and procedures. Dutch National Survey of (Morbidity and Interventions in) general practice).
- Goldberg DP, Huxley P. Mental illness in the community: the pathway to psychiatric care. London: Tavistock Publications Andover Hants, 1980.
- Hofman A. De epidemiologie als basis voor gezondheidsbeleid. In: Epidemiologie en Gezondheidsbeleid. Maas PJ van der, Hofman A, Dekker E. (red.) Samson Stafleu, Alphen aan den Rijn, 1989. (Epidemiology as a base for health policy).
- Hoogendoorn D. Stichting Informatiecentrum voor de Gezondheidszorg (SIG). In: Epidemiologie en Gezondheidsbeleid. Maas PJ van der, Hofman A, Dekker E. (red.) Samson Stafleu, Alphen aan den Rijn, 1989. (Institute Centre for data-handling in health care [SIG]).
- International Classification of Health Problems in Primary Care - Defined. Oxford: Oxford University Press, 3rd ed 1983.

- International Classification of Diseases (9th revision). Clinical Modification. Washington DC: United States Department of Health and Human Services, 1980.
- Lako CJ, Zonneveld RJ van de, Cluitmans R, Fredrix L, Vasbinder I. Subjectieve gezondheid en medische consumptie van ouderen in grote steden en op het platteland. *Tijd Soc Gezondheidsz*, 1987, 65, 131-135. (Subjective health and medical consumption of elder people in big cities and rural).
- Lamberts M, Meads S, Wood M. The reason for encounters classification for use in primary care. Field trial version July, 1982a.
- Lamberts M, Meads S, Wood M. Waarom gaat iemand naar de huisarts? Een internationale studie met de Reason for Encounter Classification. *Huisarts Wet.*, 1984; 27: 234-44. (Why goes someone to the general practitioner? An international study with Reason for Encounter Classification).
- Lamberts H. Aan de diagnose gebonden informatie uit de huisartspraktijk; van een op de prevalentie naar een op de episode georiënteerde epidemiologie. *Ned Tijdschr Geneesk* 1986; 130: 292-6. (Data attached to diagnosis from general practice. From prevalence-oriented to episode-oriented epidemiology).
- Lamberts H, Brouwer H, Groen ASM, Huisman H. Het transitie-model in de huisartspraktijk. *Huisarts Wet* 1987a; 30: 105-13. (The transition model in general practice).
- Lamberts H, Wood M. International Classification of Primary Care. Oxford: Oxford University Press, 1987b.
- Lisdonk EH van de. Ervaren en aangeboden morbiditeit in de huisartspraktijk. Een onderzoek met dagboeken. [Dissertatie]. Nijmegen: Katholieke Universiteit Nijmegen, 1985. (Experienced and offered morbidity in general practice. An investigation in general practice. An investigation with diaries [dissertation]. Nijmegen: Catholic University Nijmegen, 1985).
- Mackenbach JP, Kunst AE, Looman CWN, Habbema JDF, Maas PJ van der. Regional differences in mortality from conditions amenable to medical intervention in the Netherlands: a comparison of four time periods. *Journal of Epidemiology and Community Health*; 1988, 42, 325-332.
- Mackenbach JP, Looman CWN, Kunst AE. Regionale doodsoorzaakprofielen. *Soc Gezondheidsz.*; 1990, 68, 57-66. (Profiles of regional causes of death).
- Mausner JS, Bahn AK. In: *Epidemiology, An Introductory Text. Measures of morbidity and mortality.* WB Saunders Company Philadelphia / London / Toronto, 1974
- Meyboom WA. Registreren op: "a true level of understanding". *Huisarts Wet.*; 1979, 22, 22-4. (Registering on "a true level of understanding").
- Office of Population Censuses and Surveys. Morbidity Statistics from General Practice, Second National Study 1970-71. London: HMSO, 1974a.
- Office of Population Censuses and Surveys. Morbidity Statistics from General Practice, Second National Study 1970-71. Manual of Definitions and Procedures. London: OPCS/RCGP, 1974b.
- Office of Population Censuses and Surveys. Morbidity Statistics from General Practice, Third National Study 1981. Manual of Definitions and Procedures. London: OPCS/RCGP, 1984.
- Pieters JJJ. Andere gegevensbronnen. In: *Epidemiologie en Gezondheidsbeleid.* Maas PJ van der, Hofman A, Dekker E. (red.) Samson Stafleu, Alphen aan den Rijn, 1989. (Other sources of data).

- Royal College of General Practitioners, Office of Population Censuses and Surveys and Department of Health and Social Security. Morbidity Statistics from General Practice 1970-1971. Studies on Medical and Population Subjects No. 26 London: HMSO, 1974.
- Schellevis F, Severien R, Zee J van der. Registratie/classificatie in de huisartspraktijk. Deel V: eindrapport. Utrecht: Nederlands Huisartsen Instituut 1984a. (Registering/classification in general practice. Part V: final report).
- Schellevis F. Registratie/classificatie in de huisartspraktijk. Een vergelijking van de E-lijst met de ICHPPC-2. Utrecht/Nijmegen: Nederlands Huisartsen Instituut/Nijmeegs Universitair Huisartsen Instituut, 1984b. (Registering/classification in general practice. A comparison of the E-list with ICHPPC-2).
- Severien R, Zee J van der. Registratie/classificatie in de huisartspraktijk. Deel III: De haalbaarheid van het centraal coderen van klachten, ziekten en problemen in de huisartspraktijk. Utrecht/Nijmegen: Nederlands Huisartsen Instituut/Nijmeegs Universitair Huisartsen Instituut, 1984. (Registering/classification in general practice. Part III: The feasibility of centralized coding of complaints, diseases and problems in general practice).
- Sonsbeek J van. Gezondheidsverschillen in regionaal perspectief. Maandbericht Gezondheidsstatistiek 1987; no. 1, 5-11. (Differences of health in regional perspective).
- Velden HGM van der. Diagnose of prognose. De betekenis van de epidemiologie voor het handelen van de huisarts. Huisarts en Wet 1983; 26: 125-8. (Diagnosis or prognosis. The significance of the epidemiology for the acting of the general practitioner).
- Velden J van der. Ziekten en Verrichtingen in de huisartspraktijk. Een Nationale Studie. Medisch Contact 1986; 41: 603-6. (Diseases and interventions in general practice. A national study).
- Velden J van der, Schellevis FG, Steen J van der. ICPC (International Classification of Primary Care), tabulaire lijst. NIVEL Utrecht, 1989. (ICPC tabular list).
- Weel C van, van den Bosch WJHM, van den Hoogen HJM. De continue morbiditeitsregistratie Nijmegen. Huisarts Wet 1986; 29: 373-7. (The continuous registration of morbidity Nijmegen).
- Wijkkel D. Samenwerken en verwijzen. Dissertatie Rijksuniversiteit Groningen. Utrecht, NIVEL, 1986. (To cooperate and to refer. Dissertation State University of Groningen).

APPENDIX A Basic tables of Morbidity in Dutch general practice

A1 Explanation

This paragraph includes mainly tables. Table A shows the morbidity in Dutch general practice according to age and sex. In table B the morbidity in the Dutch general practice is distinguished to degree of urbanization, region, season and type of insurance. In this table B no standardization to age and sex is applied. Incidence and prevalence per diagnose are defined as follows.

Cumulative incidence (abbreviation i): the number of new cases of disease in a specific period, divided by the number of persons present in the population at the beginning of the considered period. In the Dutch National Survey this period amounts 3 months.

Prevalence-figure of the period (abbreviation p): the number of cases of disease in a specific period divided by the average population. The prevalence of the period is built up from the prevalence of a specific moment increased with the number of new cases and recidives in a specific period. In the Dutch National Survey this period amounts 3 months.

Conversion of the numbers to figures on an annual basis is possible by multiplying the cumulative incidence with factor 4, by which a years cumulative incidence will be obtained. Conversion of the numbers of the prevalence of particularly chronic diseases to figures on an annual basis is very complicated and only reliably feasible with more detailed information. This is because many, and particularly chronic, diseases are going over the period of registration. That is why the prevalence cannot simply be multiplied with 4. An illustration will be given with the following 4 cases.

Case 1: Cystitis

The figure of the incidence represents the total number of new cases of cystitis that came to the knowledge of the general practitioner in a 3-month period. The number of 'new cases' regards cystitis, that appears at a patient for the first time and a relapse of cystitis.

Considering the averaged short duration of the disease and the chance of relapsing, it is possible, that from one patient more than one episode of cystitis has been counted. The figure of the incidence per year can be obtained by multiplying this number by 4. The prevalence represents the total number of cases of cystitis that came within 3 months to the knowledge of the general practitioners.

In this figure of prevalence the known cases of cystitis at the beginning of registering are included. Considering the average short duration of the disease it is understood that prevalence is nearly identical to incidence. So the prevalence a year of cystitis can be obtained by multiplying the figure of prevalence by 4.

Case 2: Chronic bronchitis

The incidence of chronic bronchitis includes the total number of newly diagnosed cases of chronic bronchitis within 3 months, when by the general practitioner himself diagnosed or came to his knowledge. Exacerbations of chronic bronchitis are not considered as 'new cases'. The incidence per year is obtained by multiplying by 4.

The prevalence of chronic bronchitis represents the total number of the known cases of chronic bronchitis measured in a period of 3 months. This concerns the patients at the starting moment of registering known as such to the general practitioner and the new cases of chronic bronchitis in the 3 subsequent months.

By the nature of the registration only that patients with a known chronic bronchitis are taken along in the calculation of the prevalence, who during the registering period:

- visited the general practice because of this disease (anomaly) (for example for control or repeated prescription) or
- consulted the general practitioner, because of another disease, to which the chronic bronchitis as "underlying disease" was esteemed relevant to the general practitioner (for example: a bacterial respiratory tract infection and chronic bronchitis as underlying disease).

The prevalence of chronic bronchitis cannot simply be multiplied by 4 to calculate the prevalence per year; only with the premiss that all patients with chronic bronchitis have a contact with general practice at least once every 3 months, the prevalence figure per 3 month period is equal to the prevalence per year. This isn't however plausible.

Case 3: Fracture of ankle

The incidence of a fracture of the ankle represents the total number of new cases of this disorder, that occurs in 3 months. Both the cases diagnosed by the general practitioner as well as the cases diagnosed in the hospital and about which the general practitioner was informed within the registration period are counted in the calculation. The incidence per year is obtained by multiplying with 4.

The prevalence of an ankle fracture represents the total number known cases of this disorder in a 3 month-period. The cases of an ankle fracture known at the beginning of the registration period are also included. For example, a patient with a healing fracture, who visits his doctor in the first week of the registration for retaping his ankle.

Only with the assumption that an ankle fracture on average has healed after 3 months, the prevalence per year is the fourfold of the prevalence per 3 months.

Case 4: Depression

The incidence of depression is the total number of new cases of depression within 3 months as far as these cases were diagnosed by the general practitioner himself or otherwise came to his knowledge. When a patient earlier in the past got through an episode of depression, than it is a matter of recidive and this will be included in the calculation of the incidence. The incidence per year can be obtained by multiplying with a factor 4.

With the prevalence of depression is meant the total number of the known cases of depression, as far as these cases came to the knowledge of the general practitioner. Though this regards the patients

- who at the beginning of the registration are known with a depression and visited in the following 3 months the general practice because of the depression or a disorder to which the depression as "underlying disease" has been considered relevant by the general practitioner and
- who are diagnosed as depression for the first time or as a recidive within the 3-months registration period.

Because it is unknown which part of the patients with a known depression will visit the general practice in 3 months, the prevalence figure of 3 months cannot be extrapolated to a figure on an annual base.

A.2 Clarification of the used abbreviations in the base tables

Abbreviation	Circumscription
total	
total abs./ /1000 pers.	weighted estimate of the absolute number of cases in the Netherlands weighted estimate of the number of cases per 1000 persons
sex	
male	weighted estimate of the number of cases per 1000 males
female	weighted estimate of the number of cases per 1000 females
age categories	
0- 4	weighted estimate of the number of cases per 1000 0- 4 years old
5-14	weighted estimate of the number of cases per 1000 5-14 years old
15-24	weighted estimate of the number of cases per 1000 15-24 years old
25-44	weighted estimate of the number of cases per 1000 25-44 years old
45-64	weighted estimate of the number of cases per 1000 45-64 years old
65-74	weighted estimate of the number of cases per 1000 65-74 years old
≥ 75	weighted estimate of the number of cases per 1000 ≥ 75 years old
sex/age categories	
0- 4 male	weighted estimate of the number of cases per 1000 0- 4 years old males
0- 4 female	weighted estimate of the number of cases per 1000 0- 4 years old females
5-14 male	weighted estimate of the number of cases per 1000 5-14 years old males
5-14 female	weighted estimate of the number of cases per 1000 5-14 years old females
15-24 male	weighted estimate of the number of cases per 1000 15-24 years old males
15-24 female	weighted estimate of the number of cases per 1000 15-24 years old females
25-44 male	weighted estimate of the number of cases per 1000 25-44 years old males
25-44 female	weighted estimate of the number of cases per 1000 25-44 years old females
45-64 male	weighted estimate of the number of cases per 1000 45-64 years old males
45-64 female	weighted estimate of the number of cases per 1000 45-64 years old females
65-74 male	weighted estimate of the number of cases per 1000 65-74 years old males
65-74 female	weighted estimate of the number of cases per 1000 65-74 years old females
≥ 75 male	weighted estimate of the number of cases per 1000 ≥ 75 years old males
≥ 75 female	weighted estimate of the number of cases per 1000 ≥ 75 years old females
degree of urbanization	
rural	weighted estimate of the number of cases per 1000 persons from rural areas
sub-urban	weighted estimate of the number of cases per 1000 persons from sub urban areas
urban	weighted estimate of the number of cases per 1000 persons from urban areas
big city	weighted estimate of the number of cases per 1000 persons from big cities
distance from practice-address to nearest hospital	
same place	weighted estimate of the number of cases per 1000 persons with practice address in the same town as the hospital
1-15 km	weighted estimate of the number of cases per 1000 persons with hospital out of town and less than 15 km distant from practice-address
≥ 15 km	weighted estimate of the number of cases per 1000 persons with hospital out of town and more than 15 km distant from practice-address
season	
spring	weighted estimate of the number of cases per 1000 persons in spring
summer	weighted estimate of the number of cases per 1000 persons in summer
autumn	weighted estimate of the number of cases per 1000 persons in autumn
winter	weighted estimate of the number of cases per 1000 persons in winter
insurance-mode	
public	weighted estimate of the number of cases per 1000 public insurants
private	weighted estimate of the number of cases per 1000 private insurants

general abbreviations

abd.	abdominal
abn.	abnormal, abnormality
abnorm.	abnormal
ac.	acute
aden.	adenoid
adv.	adverse
all.	allergy
anom.	anomalies
appear.	appearance
art.	arterial
aspec.	aspecific/nonspecific
attr.	attributed
bact.	bacterial
ben.	benign
bloodf.	bloodforming
cer.	cerebri, cerebral
cerv.	cervical
chron.	chronic
circ.	circulatory
complic.	complications, complicated, complicating
complt.	complaint(s)
cond.	conditions
cong.	congenital
COPD	chronic obstructive pulmonary diseases
defic.	deficiency
deliv.	delivery
dig., digest.	digestive
dis.	disease, disorder
disab.	disability
dist.	disturbance
dysf.	dysfunction
eff.	effect
end./endocr.	endocrine
excess.	excessive
excl.	exclusive
f.p./fam. plan.	family planning
fem.	female
fract.	fracture
gen./genit.	genital
gl.	gland, glandulae
hyperkin.	hyperkinetic
hypo/am.	hypo-/amenorrhea
hypochondr.	hypochondrical
impairm.	impairment
inf.	infection, infectious
inj.	injection
invol.	involuntary
involv.	involvement
irreg.	irregular
irrit.	irritable
isch.	ischemic
IUD	intra uterine device
lig.	ligament
mm	mamma
M/morb.	Morbus
mal.	malign, malignant
med.	medical
menstr.	menstruation
met./metab.	metabolic
mov.	movements
mult.	multiple
musc. skel.	musculo-skeletal
musc.	muscular, musculo-
NEC	not elsewhere classified
neighborh.	neighborhood
neopl.	neoplasm(s)
nerv.	nervous
neur.	neurological
non-obst.	non-obstetric
non-rheum.	non-rheumatic
NOS	not otherwise specified
nutr.	nutritional
OAC	oral anticonception
obst.	obstetric
obstr.	obstructive
org.	organs

periph.	peripheral
pernic.	pernicious
Pr/D/Pu/Fp	pregnancy, delivery, puerperium/family planning
pregn.	pregnancy
prim.	primary
probl.	problem(s)
puerp.	puerperium
pyog.	pyogenes
redund.	redundant
rel.	religion, religious
resp.	respiratory
sens.	sensation/sensibility
soc.	social
spontan.	spontaneous
streptoc.	streptococcus
subcut.	subcutaneous
supp.	suppurative
surg.	surgical
sympt.	symptom(s)
syndr.	syndrome
syst.	system
thyrotox	thyrotoxicosis
tr.	tract
traum.	traumatic
treatm.	treatment
tymp.	tympanic
ulc.	ulcerosa
undesc.	undescended
unkn	unknown
unspec.	unspecified
URTI	upper respiratory tract infection
urin.	urinary
vasc.	vascular
vert.	vertiginous
w/o	without
w/wo	with or without
w.	with
wks	weeks
♀	woman
♂	man

A.3 BASIC TABLE A: MORBIDITY IN GENERAL PRACTICE BY SEX AND AGE

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
001	pain: generalized/unspec.	i	3803	0.3	0.2	0.3	0.2	0.1	0.1	0.2	0.4	0.3	1.0
		p	52184	3.6	2.5	4.6	0.2	0.2	0.7	2.1	6.1	10.2	14.7
002	chills/fever	i	8988	0.6	0.6	0.6	5.0	0.8	0.1	0.3	0.3	0.3	0.5
		p	11737	0.8	0.8	0.8	5.8	0.9	0.1	0.4	0.3	0.5	1.6
003	general weak-/tiredness	i	34029	2.3	1.7	2.9	1.5	2.3	3.1	2.1	1.7	2.9	3.8
		p	77874	5.3	3.0	7.5	2.8	2.7	4.1	4.2	5.5	10.7	17.3
004	fainting (syncope)	i	12412	0.8	0.6	1.0	0.2	0.6	1.4	0.6	0.9	0.9	1.7
		p	15305	1.0	0.8	1.3	0.3	0.7	1.6	0.7	1.2	1.4	2.0
005	symptoms infant	i	2736	0.2	0.1	0.2	3.0
		p	3045	0.2	0.2	0.2	3.4
006	allergy NEC	i	41419	2.8	2.2	3.4	4.7	2.6	3.8	3.0	1.8	2.1	1.8
		p	89572	6.1	5.0	7.2	8.3	6.7	8.0	6.3	4.8	3.9	3.2
007	feeling/behaving old/senile	i	3092	0.2	0.2	0.3	.	.	.	0.0	0.0	0.5	3.0
		p	23055	1.6	1.0	2.1	.	.	.	0.0	0.3	2.9	24.5
008	fear of disease/death	i	7107	0.5	0.3	0.6	0.2	0.0	0.4	0.7	0.5	0.6	0.6
		p	18148	1.2	0.8	1.7	0.5	0.1	0.7	1.7	1.8	1.7	1.3
009	disability/impairment NOS	i	304	0.0	0.0	0.0	.	.	0.0	0.0	.	0.0	0.3
		p	2524	0.2	0.1	0.2	.	0.0	0.0	0.1	0.1	0.5	1.5
010	general sympt./complt. NEC	i	17038	1.2	1.0	1.3	0.8	0.6	1.0	1.4	1.3	1.1	1.4
		p	30725	2.1	1.7	2.5	1.2	0.8	1.5	2.2	2.5	2.5	5.3
011	tuberculosis, general, excl. lung	i	145	0.0	0.0	0.0	.	.	0.1
		p	1121	0.1	0.1	0.1	0.1	.	0.1	0.0	0.1	0.2	0.2
012	measles	i	2714	0.2	0.2	0.2	1.5	0.5	0.1	0.0	.	.	0.1
		p	2874	0.2	0.2	0.2	1.6	0.5	0.1	0.0	.	.	0.1
013	chickenpox	i	8952	0.6	0.6	0.7	5.2	1.5	0.3	0.2	0.0	.	.
		p	9705	0.7	0.6	0.7	5.5	1.6	0.4	0.2	0.0	.	0.0
014	malaria	i	294	0.0	0.0	0.0	.	.	.	0.0	0.0	.	.
		p	304	0.0	0.0	0.0	.	.	.	0.0	0.0	.	.
015	rubella	i	12796	0.9	0.9	0.8	6.4	2.8	0.2	0.3	0.0	.	.
		p	13697	0.9	0.9	0.9	6.6	3.1	0.2	0.3	0.1	.	.
016	infectious mononucleosis	i	6818	0.5	0.4	0.5	0.2	0.7	1.5	0.3	0.1	.	.
		p	10999	0.7	0.6	0.9	0.4	1.1	2.3	0.5	0.1	.	.
017	viral dis. with exanthems NEC	i	7209	0.5	0.5	0.5	5.5	0.9	0.1	0.1	.	.	.
		p	8530	0.6	0.6	0.6	6.3	1.1	0.2	0.1	0.0	.	.
018	viral dis. NOS/NEC	i	85998	5.9	5.4	6.3	25.9	7.5	4.8	4.4	3.6	3.2	3.3
		p	91154	6.2	5.7	6.7	26.8	7.9	5.1	4.6	4.0	3.4	3.9
019	infectious dis. NOS/NEC	i	2902	0.2	0.1	0.3	0.4	0.2	0.1	0.2	0.1	0.4	0.5
		p	5610	0.4	0.3	0.4	0.7	0.3	0.2	0.4	0.3	0.4	1.0
020	carcinomatosis (prim. site unkn)	i	1254	0.1	0.1	0.1	.	.	0.1	0.0	0.1	0.3	0.7
		p	4137	0.3	0.3	0.3	.	.	0.1	0.0	0.6	1.1	1.3
021	accident/injury NOS	i	11049	0.8	0.8	0.7	0.9	0.7	0.8	0.6	0.7	0.6	1.4
		p	17892	1.2	1.3	1.2	0.9	0.9	1.5	1.1	1.2	1.5	1.6
022	mult. trauma/internal injury	i	260	0.0	0.0	0.0	.	.	0.0	0.0	.	0.1	.
		p	429	0.0	0.0	0.0	.	.	0.1	0.0	.	0.1	0.0
023	late effects of trauma	i	2455	0.2	0.2	0.1	0.1	0.0	0.1	0.2	0.2	0.2	0.0
		p	7529	0.5	0.6	0.4	0.3	0.1	0.5	0.7	0.6	0.7	0.5
024	poisoning by/abuse of med. agents	i	1270	0.1	0.0	0.1	0.1	.	0.1	0.1	0.0	0.1	0.4
		p	3518	0.2	0.2	0.3	0.3	0.0	0.3	0.2	0.2	0.2	1.1
025	poisoning by benzodiazepines	i	146	0.0	0.0	0.0	.	.	.
		p	245	0.0	0.0	0.0	.	.	.	0.0	0.0	0.0	.

A GENERAL AND UNSPECIFIED

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
001	i	0.3	.	0.1	0.2	0.1	0.1	0.2	0.3	0.3	0.5	0.2	0.4	0.6	1.2
	p	0.4	.	0.3	0.2	0.6	0.7	1.4	2.9	4.7	7.4	7.5	12.4	10.9	16.8
002	i	4.6	5.4	0.8	0.9	0.1	0.1	0.2	0.4	0.3	0.2	0.6	0.1	0.5	0.5
	p	5.5	6.2	0.9	0.9	0.2	0.1	0.3	0.4	0.3	0.4	0.8	0.3	2.3	1.3
003	i	1.6	1.5	2.1	2.5	2.2	4.0	1.3	2.9	1.5	1.9	1.6	3.9	3.9	3.8
	p	2.3	3.2	2.6	2.8	2.8	5.4	2.5	6.0	3.0	7.9	4.6	15.3	8.6	21.9
004	i	0.2	0.2	0.4	0.8	1.0	1.7	0.3	0.9	0.9	1.0	0.7	1.1	1.9	1.5
	p	0.4	0.2	0.5	0.9	1.2	2.0	0.4	1.0	1.1	1.2	1.0	1.7	2.2	1.9
005	i	2.3	3.9
	p	2.8	3.9
006	i	5.6	3.8	2.5	2.8	2.6	5.0	2.0	4.0	1.2	2.4	1.3	2.7	2.5	1.5
	p	9.9	6.7	6.2	7.1	6.2	9.8	4.3	8.3	3.5	6.1	2.8	4.8	3.9	2.8
007	i	0.0	0.0	0.1	0.4	0.6	3.7	2.6
	p	0.1	0.1	0.5	2.7	3.0	22.1	25.8
008	i	0.2	0.2	0.1	0.0	0.2	0.5	0.5	1.0	0.3	0.6	0.6	0.6	0.2	0.8
	p	0.6	0.3	0.1	0.1	0.4	1.0	1.0	2.4	1.1	2.4	1.4	1.9	0.5	1.7
009	i	0.0	.	0.0	0.0	.	.	.	0.0	0.5	0.1
	p	.	.	.	0.0	0.0	0.1	0.2	0.0	0.0	0.1	0.6	0.3	0.9	1.8
010	i	0.7	0.8	0.7	0.6	0.7	1.3	1.1	1.7	1.3	1.4	1.3	0.9	1.5	1.3
	p	1.1	1.3	0.8	0.8	1.2	1.9	1.8	2.7	2.2	2.7	2.2	2.7	4.4	5.7
011	i	0.1	0.0
	p	0.3	.	.	.	0.1	0.0	0.0	0.1	0.1	0.2	0.1	0.2	0.1	0.2
012	i	1.4	1.6	0.1	0.9	0.2	0.1	0.0	0.2	.
	p	1.5	1.7	0.1	1.0	0.2	0.1	0.0	0.2	.
013	i	4.7	5.7	1.3	1.6	0.3	0.4	0.1	0.2	.	0.0
	p	5.1	5.9	1.5	1.7	0.4	0.4	0.2	0.2	.	0.0	.	.	.	0.0
014	i	0.0	0.0	0.1
	p	0.0	0.1	0.1
015	i	7.0	5.9	2.7	2.8	0.4	0.0	0.2	0.4	0.0	0.1
	p	7.0	6.2	2.7	3.5	0.4	0.0	0.2	0.4	0.0	0.1
016	i	0.3	0.2	0.7	0.7	1.3	1.8	0.3	0.3	0.0	0.1
	p	0.4	0.5	1.0	1.2	1.6	3.1	0.4	0.6	0.1	0.1
017	i	5.3	5.8	0.8	1.1	0.1	0.2	0.1	0.1
	p	6.0	6.5	1.0	1.2	0.2	0.2	0.1	0.1	0.1
018	i	27.3	24.5	6.6	8.3	3.4	6.2	3.8	5.1	3.2	4.0	3.1	3.2	2.4	3.8
	p	28.2	25.4	7.2	8.7	3.4	6.7	4.0	5.3	3.7	4.3	3.1	3.6	3.1	4.3
019	i	0.2	0.6	0.2	0.1	0.1	0.1	0.1	0.3	0.1	0.2	0.5	0.3	0.2	0.7
	p	0.6	0.8	0.2	0.3	0.3	0.2	0.3	0.5	0.1	0.4	0.5	0.4	1.4	0.8
020	i	0.1	0.0	0.0	0.1	0.0	0.6	0.1	0.6	0.8
	p	0.1	0.0	0.0	0.6	0.6	1.7	0.7	1.2	1.4
021	i	0.9	0.9	0.7	0.7	1.2	0.5	0.8	0.5	0.7	0.7	0.1	1.0	1.0	1.6
	p	0.9	0.9	1.0	0.9	2.1	0.9	1.1	1.1	1.1	1.3	0.9	2.0	1.2	1.9
022	i	0.1	0.0	0.0	0.1	.	.
	p	0.1	0.1	0.1	0.1	.	0.0
023	i	0.2	0.1	0.0	0.0	0.2	0.1	0.2	0.2	0.3	0.2	0.3	0.2	0.0	.
	p	0.2	0.4	0.1	0.1	0.5	0.5	0.8	0.5	0.7	0.5	0.9	0.6	0.7	0.4
024	i	0.1	0.1	.	.	0.0	0.2	0.0	0.1	0.0	0.0	.	0.2	.	0.7
	p	0.4	0.3	.	0.1	0.1	0.5	0.1	0.3	0.2	0.1	0.0	0.3	1.6	0.9
025	i	0.1
	p	0.0	0.1	.	.	0.1	.	.

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
026	poisoning by med. agents NOS	i	1508	0.1	0.1	0.1	0.2	.	0.1	0.0	0.1	0.2	0.7
		p	2800	0.2	0.2	0.2	0.3	.	0.2	0.1	0.2	0.4	1.0
027	adv. eff. med. agent proper dose: all.	i	9612	0.7	0.5	0.8	1.0	0.4	0.4	0.6	0.7	0.6	1.4
		p	13025	0.9	0.6	1.2	1.2	0.6	0.4	0.9	0.9	1.1	2.6
028	adv. eff. OAC	i	9412	0.6	.	1.3	.	0.1	2.1	0.8	0.0	.	.
		p	12706	0.9	.	1.7	.	0.1	2.8	1.1	0.1	.	.
029	adv. eff. med. agent NEC	i	18751	1.3	0.8	1.7	1.2	0.1	0.8	0.9	1.8	2.3	4.5
		p	34031	2.3	1.6	3.0	1.9	0.3	1.2	1.6	3.3	5.5	7.6
030	toxic eff. other substances	i	2663	0.2	0.2	0.2	0.3	0.1	0.1	0.2	0.1	0.3	0.0
		p	4715	0.3	0.3	0.4	0.6	0.2	0.2	0.3	0.3	0.3	0.3
031	complic. surg./med. treatm.: hematoma	i	484	0.0	0.0	0.0	0.1	0.0	0.0	0.1	0.0	0.0	.
		p	1195	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.0	0.3	0.2
032	complic. surg./med. treatm.: wounds	i	2272	0.2	0.2	0.1	0.0	0.0	0.2	0.1	0.3	0.2	.
		p	5585	0.4	0.4	0.4	0.0	0.1	0.3	0.3	0.7	0.5	0.4
033	complic. surg./med. treatm.: NEC	i	490	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.2
		p	35253	2.4	2.1	2.7	0.9	0.8	0.9	1.9	3.6	6.3	6.3
034	adv. eff. physical factors NEC	i	4560	0.3	0.2	0.4	0.3	0.6	0.3	0.4	0.2	0.0	0.2
		p	9790	0.7	0.4	0.9	0.9	0.8	0.7	0.7	0.6	0.3	0.4
035	adv. eff. prosthetic device	i	1480	0.1	0.0	0.2	.	0.0	.	0.0	0.2	0.2	0.6
		p	3022	0.2	0.1	0.3	.	0.0	0.1	0.1	0.4	0.7	0.8
036	mult. congenital syndromes	i	189	0.0	0.0	0.0	0.1	0.0
		p	3802	0.3	0.3	0.2	1.4	0.4	0.2	0.3	0.0	0.1	.
037	abnormal results investigation NEC	i	1045	0.1	0.0	0.1	0.1	0.0	0.0	0.0	0.2	0.1	.
		p	3062	0.2	0.1	0.3	0.1	0.0	0.1	0.1	0.4	0.6	0.3
038	toxoplasmosis	i	110	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	574	0.0	0.0	0.1	.	.	0.1	0.1	.	.	.
039	premat./immature liveborn infant	i	270	0.0	0.0	0.0	0.3
		p	1014	0.1	0.1	0.1	1.0
040	all perinatal morbidity	i	1126	0.1	0.1	0.1	1.2
		p	3232	0.2	0.2	0.2	3.5
041	perinatal mortality	i	85	0.0	0.0	.	0.1
		p	0
042	death (excl. perinatal)	i	12965	0.9	1.0	0.8	0.3	0.0	0.0	0.1	0.8	2.9	8.5
		p	0
043	no disease	i	158074	10.8	8.3	13.2	20.7	8.3	9.9	10.1	10.3	12.2	12.0
		p	270343	18.4	13.6	23.1	28.1	11.0	13.3	16.0	19.9	29.4	35.7
044	disease NEC	i	3610	0.2	0.2	0.3	0.1	0.2	0.2	0.3	0.3	0.3	0.4
		p	6639	0.5	0.3	0.6	0.2	0.2	0.3	0.4	0.8	0.6	0.9

A GENERAL AND UNSPECIFIED

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
026	i	0.3	0.1	.	.	0.1	0.0	0.0	0.0	0.2	0.0	.	0.4	0.5	0.8
	p	0.3	0.3	.	.	0.3	0.0	0.1	0.1	0.3	0.1	0.2	0.6	0.5	1.3
027	i	1.6	0.4	0.6	0.3	0.3	0.6	0.5	0.8	0.1	1.3	0.2	1.0	1.9	1.1
	p	1.9	0.5	0.6	0.6	0.3	0.6	0.6	1.1	0.2	1.6	0.6	1.5	2.2	2.8
028	i	.	.	.	0.2	.	4.2	.	1.6	.	0.1
	p	.	.	.	0.2	.	5.5	.	2.3	.	0.1
029	i	1.2	1.1	0.0	0.2	0.5	1.1	0.5	1.4	1.3	2.4	1.4	2.9	4.2	4.6
	p	1.7	2.0	0.3	0.2	0.6	1.8	1.0	2.3	2.3	4.2	4.3	6.4	7.0	8.0
030	i	0.3	0.3	0.2	0.1	0.1	0.2	0.2	0.3	0.1	0.2	0.4	0.1	0.1	.
	p	0.5	0.7	0.2	0.2	0.1	0.4	0.3	0.4	0.3	0.4	0.4	0.1	0.2	0.4
031	i	.	0.1	0.0	.	.	0.0	0.1	0.1	.	0.0	0.1	.	.	.
	p	.	0.1	0.0	0.1	.	0.0	0.1	0.1	0.0	0.1	0.5	0.1	0.3	0.2
032	i	0.1	.	0.0	0.0	0.2	0.2	0.2	0.1	0.3	0.3	0.3	0.1	.	.
	p	0.1	.	0.1	0.1	0.3	0.4	0.3	0.4	0.8	0.6	0.5	0.5	0.5	0.4
033	i	0.0	0.0	.	0.0	0.1	.	.	0.0	0.0	0.0	0.0	.	.	0.3
	p	1.0	0.8	0.5	1.0	1.0	0.9	1.4	2.3	3.2	4.0	7.2	5.6	6.5	6.1
034	i	0.2	0.3	0.1	1.0	0.0	0.5	0.3	0.4	0.2	0.3	0.0	0.0	0.3	0.2
	p	0.6	1.2	0.4	1.3	0.4	0.9	0.4	1.0	0.3	0.8	0.2	0.4	0.3	0.4
035	i	.	.	0.0	0.0	.	.	0.0	0.0	0.1	0.3	.	0.4	0.1	1.0
	p	.	.	0.0	0.0	0.1	0.1	0.0	0.2	0.3	0.4	0.8	0.5	0.1	1.2
036	i	0.0	0.3	0.1
	p	1.2	1.6	0.5	0.2	0.2	0.2	0.3	0.2	0.0	0.0	0.1	.	.	.
037	i	.	0.2	.	0.1	.	0.1	0.0	0.1	0.0	0.3	.	0.2	.	.
	p	.	0.3	.	0.1	0.2	0.1	0.0	0.2	0.3	0.5	0.2	1.0	0.3	0.3
038	i	0.0
	p	0.1	0.0	0.2
039	i	0.3	0.3
	p	1.2	0.9
040	i	1.4	1.1
	p	3.8	3.3
041	i	0.2
	p
042	i	0.5	0.0	0.0	.	0.0	0.0	0.1	0.2	0.8	0.8	5.0	1.2	10.6	7.4
	p
043	i	20.7	20.6	8.0	8.6	5.9	14.0	6.9	13.6	7.5	13.0	10.9	13.1	12.0	12.1
	p	27.5	28.8	11.0	11.1	7.8	19.0	10.8	21.3	14.7	25.0	24.3	33.3	27.4	40.1
044	i	.	0.2	0.1	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.6	0.3
	p	.	0.4	0.1	0.2	0.2	0.4	0.3	0.5	0.4	1.2	0.7	0.5	1.0	0.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
045	enlarged lymph gland(s)	i	14170	1.0	0.8	1.1	1.6	1.6	1.7	0.9	0.3	0.3	0.1
		p	17731	1.2	1.1	1.3	1.8	1.9	2.1	1.2	0.4	0.5	0.1
046	sympt. lymph glands(s) NEC	i	381	0.0	0.0	0.0	0.1	0.0	0.1	0.0	0.0	.	0.0
		p	728	0.0	0.0	0.1	0.2	0.0	0.1	0.0	0.0	0.0	0.0
047	fear of blood/lymph disease	i	3184	0.2	0.2	0.2	.	0.1	0.2	0.4	0.1	.	0.1
		p	4134	0.3	0.3	0.3	.	0.1	0.3	0.5	0.2	0.1	0.1
048	disability/impairment blood	i	0
		p	0
049	sympt./complt. blood NOS/NEC	i	73	0.0	.	0.0	0.0	0.1
		p	246	0.0	0.0	0.0	.	.	0.0	.	0.0	0.0	0.1
050	acute lymphadenitis/-angitis	i	8770	0.6	0.6	0.6	1.4	1.0	0.8	0.5	0.2	0.5	0.1
		p	9649	0.7	0.7	0.6	1.7	1.1	0.9	0.6	0.3	0.4	0.1
051	chron. aspec. lymphadenitis	i	1290	0.1	0.1	0.1	0.2	0.0	0.1	0.2	0.0	.	0.1
		p	1472	0.1	0.1	0.1	0.2	0.0	0.1	0.2	0.0	0.0	0.1
052	malignant lymphoma	i	277	0.0	0.0	0.0	.	.	0.0	0.0	0.0	0.0	0.0
		p	2161	0.1	0.1	0.2	.	.	0.1	0.2	0.2	0.3	0.1
053	leukemia	i	434	0.0	0.0	0.1	0.0	.	0.0	.	0.1	.	0.1
		p	3882	0.3	0.3	0.2	0.1	0.2	0.0	0.0	0.4	0.7	1.7
054	malignant neopl. blood NEC	i	757	0.1	0.0	0.1	.	.	0.0	0.0	0.1	0.5	0.0
		p	3804	0.3	0.2	0.3	.	.	0.0	0.1	0.4	1.3	1.0
055	benigne neopl. blood/bloodform. org.	i	216	0.0	0.0	0.0	0.0	.	0.0	0.0	0.0	.	.
		p	304	0.0	0.0	0.0	0.0	.	0.0	0.0	0.0	.	.
056	ruptured spleen	i	45	0.0	0.0	0.0	.	.	.
		p	45	0.0	0.0	0.0	.	.	.
057	injuries blood/bloodform. org. NEC	i	0
		p	22	0.0	.	0.0	0.0
058	hered. haemolytic anaemias	i	33	0.0	0.0	.	0.0
		p	583	0.0	0.0	0.1	0.0	0.0	0.1	0.1	.	.	.
059	cong. anom. blood/bloodf. org. NEC	i	64	0.0	.	0.0
		p	539	0.0	0.0	0.0	0.2	0.1	.	0.0	0.0	.	.
060	iron deficiency anemia	i	16120	1.1	0.5	1.7	0.8	1.4	1.0	1.4	0.5	0.8	1.7
		p	67408	4.6	1.7	7.4	3.6	2.6	2.7	5.4	3.3	4.6	16.3
061	pernic./other deficiency anemia	i	2303	0.2	0.1	0.2	0.0	.	0.1	0.1	0.1	0.3	1.0
		p	29629	2.0	1.2	2.8	0.1	0.0	0.3	0.7	2.6	6.3	14.5
062	anemia NOS/NEC	i	7485	0.5	0.2	0.8	0.3	0.4	0.4	0.6	0.5	0.3	1.2
		p	27862	1.9	0.8	2.9	0.7	1.5	1.4	1.7	2.0	3.0	5.3
063	hemophilia/coagulation disorders	i	1275	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.2	0.0	0.1
		p	5746	0.4	0.4	0.4	0.5	0.3	0.1	0.1	0.7	1.0	1.0
064	abnormal white cells	i	65	0.0	.	0.0	0.0	.	.
		p	638	0.0	0.0	0.1	0.0	0.0	.	0.1	0.1	0.1	.
065	unexplained abn. blood test	i	650	0.0	0.1	0.0	.	0.0	0.1	0.0	0.1	0.1	0.1
		p	2025	0.1	0.2	0.1	.	0.1	0.1	0.1	0.2	0.3	0.5
066	abn. blood test NOS	i	562	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
		p	3148	0.2	0.2	0.3	0.2	0.0	0.1	0.1	0.5	0.4	0.5
067	splenomegaly	i	164	0.0	.	0.0	0.2
		p	432	0.0	.	0.1	.	.	.	0.0	.	0.2	0.3
068	HIV-infection (AIDS/ARS)	i	39	0.0	0.0	.	.	.	0.0
		p	417	0.0	0.1	0.0	.	.	0.0	0.0	.	0.1	.
069	diseases blood/bloodf. org. NOS/NEC	i	745	0.1	0.0	0.1	.	.	0.1	0.0	0.1	.	0.0
		p	3564	0.2	0.2	0.2	0.2	0.1	0.1	0.2	0.4	0.4	0.8

B BLOOD, BLOODFORM. ORG., LYMPHATICS, SPLEEN

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
045	i	1.7	1.5	1.8	1.5	1.5	1.9	0.5	1.4	0.3	0.3	0.4	0.2	0.1	0.1
	p	2.0	1.5	2.1	1.8	1.9	2.2	0.6	1.8	0.3	0.4	0.9	0.2	0.1	0.1
046	i	0.2	.	.	0.0	0.1	0.0	0.0	0.0	.	0.0	.	.	0.0	.
	p	0.2	0.2	0.0	0.0	0.1	0.1	0.0	0.1	.	0.0	.	0.1	0.0	.
047	i	.	.	0.0	0.2	0.2	0.3	0.4	0.5	0.1	0.2	.	.	0.1	0.0
	p	.	.	0.0	0.2	0.2	0.3	0.6	0.5	0.2	0.3	0.2	.	0.1	0.0
048	i
	p
049	i	0.0	.	0.1
	p	0.1	0.0	0.1	0.0	.	0.1
050	i	1.3	1.4	1.1	0.8	0.8	0.9	0.5	0.4	0.1	0.4	0.6	0.4	0.3	.
	p	1.7	1.8	1.3	0.8	0.8	0.9	0.6	0.6	0.2	0.4	0.6	0.3	0.3	.
051	i	0.0	0.4	0.0	0.1	0.1	0.1	0.2	0.2	.	0.0	.	.	.	0.1
	p	0.0	0.4	0.0	0.1	0.1	0.2	0.2	0.2	.	0.0	0.0	.	.	0.1
052	i	0.0	.	.	0.1	0.0	.	0.0	.	.	0.0
	p	0.1	.	0.2	0.2	0.2	0.3	0.4	0.2	0.2	0.1
053	i	.	0.0	.	.	0.0	.	.	.	0.0	0.2	.	.	.	0.2
	p	0.0	0.2	0.2	0.2	0.0	.	0.1	0.0	0.5	0.3	0.9	0.5	2.6	1.2
054	i	0.1	0.0	.	0.0	0.1	0.4	0.5	.	0.0
	p	0.0	0.1	0.1	0.1	0.4	0.4	1.7	1.1	0.8	1.2
055	i	0.1	.	.	.	0.0	0.0	0.0	0.0	0.1
	p	0.1	.	.	.	0.0	0.0	0.0	0.0	0.1
056	i	0.0
	p	0.0
057	i
	p	0.0
058	i	0.0
	p	0.0	0.1	0.1	.	.	0.2	0.0	0.1
059	i
	p	0.3	.	0.0	0.1	.	.	0.0	0.1	.	0.0
060	i	0.6	0.9	1.7	1.1	0.0	2.0	0.2	2.7	0.2	0.8	0.4	1.2	2.0	1.6
	p	4.1	3.1	3.3	1.9	0.2	5.3	0.6	10.4	1.0	5.6	2.1	6.5	10.0	19.7
061	i	0.0	.	.	.	0.0	0.2	.	0.3	0.2	.	0.2	0.5	0.4	1.4
	p	0.1	0.1	0.0	0.0	0.0	0.6	0.3	1.2	1.9	3.3	5.3	7.0	10.7	16.6
062	i	0.3	0.3	0.4	0.5	0.1	0.7	0.1	1.0	0.2	0.8	0.1	0.5	0.6	1.5
	p	0.8	0.6	1.6	1.3	0.3	2.4	0.3	3.2	0.4	3.5	2.8	3.1	3.9	6.1
063	i	0.2	0.0	.	0.2	0.0	0.0	0.0	0.0	0.2	0.3	0.0	.	.	0.2
	p	0.7	0.4	0.4	0.2	0.2	0.1	0.1	0.1	0.6	0.7	1.4	0.7	1.1	0.9
064	i	0.0
	p	.	0.1	.	0.0	.	.	0.1	0.0	0.0	0.1	.	0.2	.	.
065	i	.	.	0.0	.	0.0	0.1	0.0	0.0	0.2	0.0	.	0.1	.	0.1
	p	.	.	0.1	0.0	0.1	0.2	0.1	0.0	0.3	0.1	0.5	0.1	0.2	0.6
066	i	.	0.2	0.0	.	.	0.1	0.0	0.0	0.1	.	.	0.1	0.1	.
	p	0.2	0.2	0.0	.	.	0.2	0.1	0.1	0.3	0.6	0.3	0.4	0.6	0.5
067	i	0.3
	p	0.0	.	.	.	0.3	.	.	0.5
068	i	0.0
	p	0.1	.	0.1	0.0	.	.	0.1	.	.	.
069	i	0.1	0.0	0.0	0.0	0.0	0.2	.	.	0.0	0.0
	p	0.3	0.0	0.1	0.1	0.1	0.1	0.2	0.1	0.4	0.4	0.3	0.5	1.0	0.6

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
070	generalized abd. pain/cramps	i	30073	2.0	1.6	2.5	3.2	3.8	2.3	1.6	1.5	1.6	1.3
		p	47636	3.2	2.2	4.3	3.9	4.5	2.6	2.7	3.0	4.0	4.7
071	stomach pain	i	11011	0.8	0.8	0.7	0.1	0.1	0.7	0.8	1.0	1.2	1.6
		p	63831	4.3	4.5	4.2	0.2	0.2	1.4	3.0	7.9	11.6	13.7
072	heartburn	i	5007	0.3	0.2	0.4	.	0.1	0.2	0.4	0.3	0.8	0.5
		p	20712	1.4	1.2	1.6	.	0.1	0.4	1.2	2.3	3.7	4.2
073	peri-anal itching	i	4487	0.3	0.3	0.3	0.1	0.0	0.3	0.3	0.6	0.4	0.3
		p	7638	0.5	0.6	0.4	0.2	0.1	0.3	0.5	1.0	0.7	0.7
074	localised abd. pain	i	8273	0.6	0.5	0.7	0.1	0.3	0.4	0.7	0.6	0.7	0.7
		p	12281	0.8	0.7	1.0	0.2	0.4	0.6	0.9	1.1	1.1	1.6
075	flatulence/belching	i	2680	0.2	0.1	0.3	.	0.1	0.2	0.2	0.3	0.1	0.2
		p	3923	0.3	0.2	0.4	.	0.1	0.3	0.2	0.4	0.4	0.8
076	nausea	i	3657	0.2	0.2	0.3	0.1	0.0	0.3	0.2	0.3	0.4	0.6
		p	9548	0.7	0.5	0.8	0.2	0.1	0.5	0.5	0.9	1.2	2.0
077	vomiting	i	3205	0.2	0.2	0.2	0.6	0.1	0.2	0.1	0.1	0.2	0.3
		p	6306	0.4	0.4	0.5	0.7	0.3	0.3	0.3	0.2	0.9	0.8
078	diarrhoea	i	17540	1.2	1.1	1.3	3.5	0.6	0.8	1.1	1.0	1.0	3.0
		p	29374	2.0	1.7	2.3	5.5	0.9	1.3	1.6	1.8	2.6	4.8
079	constipation	i	18074	1.2	0.8	1.7	2.8	0.6	0.8	0.8	1.0	1.9	4.8
		p	87685	6.0	3.8	8.1	5.0	1.2	1.3	2.4	5.7	16.0	42.6
080	jaundice	i	265	0.0	0.0	0.0	0.0	0.0	0.0	.	.	0.1	0.0
		p	367	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.0
081	blood vomiting	i	86	0.0	0.0	0.0	0.1
		p	189	0.0	0.0	0.0	.	.	.	0.0	.	.	0.2
082	melena	i	1378	0.1	0.1	0.1	0.0	.	0.0	0.1	0.1	0.4	0.3
		p	3310	0.2	0.2	0.2	0.0	.	0.0	0.1	0.4	0.9	0.8
083	rectal bleeding	i	1230	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.2	0.2
		p	2105	0.1	0.2	0.1	0.1	0.0	0.1	0.1	0.1	0.4	0.6
084	fecal incontinence	i	258	0.0	0.0	0.0	0.0	0.1	0.2
		p	1024	0.1	0.1	0.1	.	.	0.0	.	0.1	0.2	0.5
085	defecation anomalies NEC	i	2771	0.2	0.1	0.2	0.4	0.1	0.1	0.1	0.3	0.1	0.3
		p	3973	0.3	0.2	0.3	0.5	0.1	0.2	0.2	0.4	0.4	0.7
086	sympt./complt. of teeth, gums	i	4266	0.3	0.2	0.3	1.7	0.3	0.3	0.2	0.1	0.2	0.0
		p	10522	0.7	0.8	0.7	2.9	0.8	0.8	0.7	0.2	0.4	0.2
087	sympt./complt. mouth, tongue, lip	i	5724	0.4	0.3	0.5	1.4	0.4	0.2	0.3	0.3	0.4	0.2
		p	9855	0.7	0.5	0.8	1.7	0.8	0.6	0.5	0.6	0.7	0.6
088	swallowing problems	i	5195	0.4	0.2	0.5	0.4	0.0	0.3	0.4	0.6	0.4	0.2
		p	7143	0.5	0.3	0.7	0.5	0.1	0.3	0.5	0.9	0.6	0.3
089	fear of digestive diseases	i	1759	0.1	0.1	0.1	.	0.0	.	0.2	0.1	0.3	0.1
		p	3786	0.3	0.2	0.3	.	0.1	0.0	0.3	0.3	0.7	0.2
090	digestive disability/impairment	i	8	0.0	.	0.0	.	.	0.0
		p	8	0.0	.	0.0	.	.	0.0
091	sympt./complt. digestive NEC	i	3921	0.3	0.2	0.3	0.2	0.2	0.1	0.3	0.3	0.1	0.4
		p	6565	0.4	0.4	0.5	0.2	0.2	0.1	0.4	0.7	0.9	0.6
092	infectious diarrhoea/dysentery	i	5542	0.4	0.3	0.4	1.1	0.3	0.4	0.4	0.3	0.4	0.0
		p	7328	0.5	0.4	0.6	1.5	0.4	0.5	0.5	0.3	0.4	0.2
093	mumps	i	12609	0.9	0.9	0.9	5.0	3.2	0.3	0.2	0.0	0.0	0.2
		p	13001	0.9	0.9	0.9	5.1	3.2	0.3	0.3	0.0	0.0	0.2
094	infectious hepatitis	i	371	0.0	0.0	0.0	.	.	0.0	0.1	0.0	.	.
		p	1773	0.1	0.1	0.1	.	0.1	0.0	0.2	0.2	0.0	0.1

D DIGESTIVE

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
070	i	3.0	3.4	3.2	4.5	1.5	3.0	1.1	2.1	1.1	1.9	1.4	1.8	0.3	1.8
	p	4.1	3.6	3.4	5.6	1.5	3.7	1.6	4.0	2.2	3.7	2.4	5.2	2.4	5.8
071	i	0.0	0.1	0.0	0.1	0.5	0.8	1.0	0.6	1.2	0.8	1.1	1.2	2.1	1.4
	p	0.2	0.1	0.2	0.2	1.0	1.8	3.1	2.8	8.7	7.0	14.3	9.4	15.1	12.9
072	i	.	.	0.1	0.1	0.1	0.3	0.3	0.4	0.3	0.4	0.2	1.3	0.4	0.6
	p	.	.	0.1	0.1	0.1	0.6	0.9	1.4	2.8	1.7	2.8	4.5	3.3	4.7
073	i	0.1	.	.	0.1	0.3	0.3	0.3	0.2	0.7	0.5	0.1	0.6	0.1	0.4
	p	0.2	0.1	0.0	0.1	0.4	0.3	0.8	0.3	1.1	0.8	0.4	0.9	0.6	0.7
074	i	0.2	.	0.6	0.0	0.2	0.7	0.6	0.9	0.3	0.9	1.0	0.6	0.5	0.8
	p	0.4	.	0.6	0.1	0.4	0.9	0.7	1.1	0.8	1.4	1.4	1.0	1.4	1.7
075	i	.	.	.	0.2	0.1	0.4	0.1	0.2	0.1	0.4	0.2	0.0	0.1	0.2
	p	.	.	.	0.2	0.1	0.4	0.1	0.3	0.2	0.5	0.6	0.2	0.9	0.8
076	i	0.0	0.1	0.1	0.0	0.3	0.4	0.0	0.3	0.3	0.3	0.3	0.5	0.4	0.7
	p	0.2	0.2	0.1	0.1	0.3	0.6	0.3	0.7	0.9	0.9	1.1	1.4	0.6	2.7
077	i	0.8	0.5	0.1	0.2	0.3	0.0	0.1	0.1	0.1	0.1	0.3	0.1	.	0.4
	p	0.8	0.6	0.3	0.3	0.4	0.1	0.2	0.3	0.1	0.2	1.3	0.5	.	1.2
078	i	3.5	3.5	0.7	0.6	0.7	0.9	1.2	1.0	0.8	1.2	0.4	1.4	3.0	3.0
	p	5.8	5.3	1.1	0.8	1.1	1.5	1.6	1.7	1.5	2.2	1.1	3.8	4.1	5.1
079	i	2.9	2.8	0.5	0.7	0.4	1.2	0.3	1.3	0.4	1.6	1.7	2.0	4.4	5.1
	p	5.1	4.9	0.9	1.5	0.5	2.2	1.1	3.6	3.5	7.9	13.6	17.9	35.5	46.4
080	i	0.0	.	0.1	.	.	0.0	0.1	0.1	.	0.0
	p	0.0	0.0	0.1	.	.	0.0	0.0	.	0.1	.	0.1	0.1	.	0.0
081	i	0.2	0.0
	p	0.0	0.3	0.1
082	i	.	0.1	.	.	.	0.0	0.1	0.0	0.2	0.1	0.2	0.5	0.2	0.3
	p	.	0.1	.	.	.	0.0	0.2	0.0	0.4	0.3	0.7	1.1	0.9	0.8
083	i	.	0.2	.	0.0	0.1	.	0.1	0.0	0.1	0.1	0.2	0.1	.	0.3
	p	.	0.2	.	0.0	0.1	0.1	0.2	0.0	0.1	0.1	0.4	0.4	0.5	0.6
084	i	0.0	0.0	0.1	0.0	0.4	0.1
	p	0.1	.	.	0.2	0.0	0.5	0.0	0.5	0.5
085	i	0.3	0.5	0.1	0.1	0.0	0.2	0.1	0.2	0.2	0.3	0.1	0.1	0.3	0.3
	p	0.5	0.6	0.1	0.1	0.1	0.2	0.1	0.2	0.4	0.4	0.1	0.7	0.5	0.8
086	i	1.4	2.0	0.2	0.3	0.1	0.4	0.3	0.2	0.0	0.1	0.3	0.2	0.0	0.0
	p	2.8	3.0	1.1	0.6	0.8	0.8	0.7	0.7	0.2	0.1	0.3	0.5	0.1	0.2
087	i	1.2	1.7	0.4	0.4	0.1	0.3	0.2	0.5	0.2	0.4	0.7	0.2	0.3	0.2
	p	1.7	1.7	0.8	0.9	0.5	0.6	0.3	0.8	0.4	0.8	1.0	0.5	0.3	0.7
088	i	0.4	0.3	0.0	0.0	0.1	0.4	0.2	0.5	0.4	0.9	0.6	0.3	0.0	0.2
	p	0.6	0.3	0.0	0.1	0.1	0.5	0.4	0.7	0.4	1.3	0.6	0.5	0.1	0.4
089	i	.	.	.	0.1	.	.	0.2	0.2	0.2	0.1	0.6	0.1	.	0.1
	p	.	.	.	0.1	.	0.1	0.4	0.3	0.3	0.4	0.8	0.6	.	0.3
090	i	0.0
	p	0.0
091	i	0.4	0.0	0.2	0.3	0.0	0.3	0.2	0.5	0.3	0.4	.	0.2	0.9	0.2
	p	0.4	0.0	0.2	0.3	0.0	0.3	0.3	0.6	0.7	0.7	1.0	0.8	0.9	0.5
092	i	1.1	1.1	0.2	0.4	0.3	0.6	0.4	0.3	0.2	0.3	0.0	0.7	0.0	0.0
	p	1.3	1.8	0.3	0.5	0.3	0.7	0.5	0.5	0.2	0.4	0.0	0.7	0.1	0.2
093	i	5.2	4.8	3.2	3.2	0.1	0.4	0.2	0.3	0.1	0.0	.	0.0	.	0.4
	p	5.2	5.1	3.3	3.2	0.2	0.4	0.3	0.3	0.1	0.0	.	0.0	.	0.4
094	i	0.1	0.0	0.1	0.0
	p	.	.	.	0.2	.	0.1	0.2	0.2	0.2	0.3	.	0.0	0.2	.

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
095	parasites/worms	i	48290	3.3	2.9	3.7	6.7	12.2	2.3	2.4	0.7	0.4	0.1
		p	50949	3.5	3.0	3.9	6.8	12.9	2.4	2.6	0.7	0.5	0.1
096	other presumed infections	i	96720	6.6	6.6	6.6	21.8	6.1	6.7	5.3	5.0	4.0	7.6
		p	102233	7.0	6.9	7.0	22.8	6.5	6.9	5.5	5.5	4.7	7.7
097	malign. neopl. stomach	i	527	0.0	0.0	0.0	0.0	0.2	0.3
		p	3640	0.2	0.3	0.2	.	.	.	0.0	0.4	0.8	1.8
098	malign. neopl. colon, rectum	i	920	0.0	0.0	0.0	0.2	0.3	0.6
		p	10599	0.7	0.6	0.9	.	.	0.0	0.1	1.1	2.7	5.6
099	malign. neopl. colon, rectum	i	425	0.0	0.0	0.0	.	.	.	0.0	0.1	0.1	0.1
		p	4508	0.3	0.3	0.3	.	.	.	0.1	0.6	0.7	2.0
100	malign. neopl. pancreas	i	503	0.0	0.0	0.0	0.0	0.3	0.2
		p	1601	0.1	0.1	0.1	.	.	.	0.0	0.2	0.5	0.7
101	malign. neopl. dig. syst. NEC	i	1002	0.1	0.1	0.1	0.1	0.4	0.4
		p	6426	0.4	0.4	0.5	.	.	.	0.1	0.5	1.7	3.5
102	benign. neoplasms dig. syst. NEC	i	452	0.0	0.0	0.0	0.1	0.0	.	0.0	0.0	0.1	0.2
		p	1389	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.1	0.4	0.3
103	neoplasms dig. syst. NEC	i	769	0.1	0.1	0.0	.	.	.	0.0	0.0	0.1	0.4
		p	2113	0.1	0.1	0.2	.	.	.	0.1	0.2	0.3	1.1
104	foreign body through orifice	i	1207	0.1	0.1	0.1	0.8	0.1	0.0	0.1	0.0	0.0	.
		p	1634	0.1	0.1	0.1	1.1	0.1	0.0	0.1	0.0	0.0	.
105	injuries dig. syst. NEC	i	203	0.0	0.0	0.0	0.1	.	0.0	0.0	.	.	.
		p	263	0.0	0.0	0.0	0.1	.	0.0	0.0	0.0	0.0	.
106	cong. anom. digest. system	i	566	0.0	0.0	0.0	0.4	0.0	.	0.0	.	.	.
		p	2200	0.1	0.2	0.1	0.8	0.4	0.1	0.0	0.1	0.1	.
107	disease of teeth/gums	i	13360	0.9	0.8	1.0	0.9	0.6	0.9	1.1	0.9	0.8	0.7
		p	20833	1.4	1.3	1.5	1.8	1.4	1.5	1.7	1.2	0.9	0.9
108	disease of mouth/tongue/lips	i	32400	2.2	1.9	2.5	4.9	2.8	2.2	1.7	2.2	1.8	1.7
		p	43043	2.9	2.4	3.5	6.1	3.4	2.6	2.3	2.9	2.4	3.5
109	disease of esophagus	i	8660	0.6	0.6	0.6	.	0.0	0.4	0.5	1.1	1.2	0.9
		p	16790	1.1	1.1	1.2	0.2	0.1	0.6	0.7	2.2	2.5	3.4
110	duodenal ulcer	i	9653	0.7	0.9	0.5	0.0	0.0	0.4	0.9	1.3	0.7	0.1
		p	29761	2.0	2.7	1.4	0.0	0.0	0.5	2.2	4.2	3.6	2.3
111	gastric ulcers NEC	i	2958	0.2	0.3	0.2	.	0.1	0.1	0.2	0.4	0.1	0.8
		p	12102	0.8	0.9	0.7	.	0.1	0.1	0.5	1.8	1.3	3.6
112	peptic ulcers NEC	i	1484	0.1	0.1	0.1	.	.	0.0	0.1	0.3	0.1	0.4
		p	5793	0.4	0.5	0.3	.	0.0	0.1	0.2	1.0	0.8	0.8
113	disord. stomach function	i	46081	3.1	3.2	3.1	1.7	1.1	3.3	3.7	3.5	3.7	3.9
		p	74467	5.1	5.3	4.8	2.0	1.2	4.1	5.2	7.2	8.1	7.9
114	appendicitis	i	6175	0.4	0.4	0.4	0.1	0.9	0.7	0.4	0.1	0.3	0.0
		p	7328	0.5	0.5	0.5	0.1	1.1	0.8	0.4	0.2	0.4	0.1
115	inguinal hernia	i	8407	0.6	0.9	0.2	2.0	0.3	0.2	0.4	0.6	0.7	1.6
		p	11613	0.8	1.3	0.3	2.5	0.4	0.4	0.5	0.8	1.5	2.0
116	hiatus (diaphr.) hernia	i	10787	0.7	0.6	0.9	.	0.0	0.4	0.6	1.1	2.0	2.1
		p	29973	2.0	1.3	2.7	.	0.1	0.5	1.0	3.5	6.4	9.0
117	abdominal hernia NEC	i	2929	0.2	0.2	0.2	0.4	.	0.1	0.1	0.3	0.6	0.3
		p	4164	0.3	0.3	0.3	0.4	.	0.1	0.2	0.4	1.1	0.7
118	diverticular dis. intestines	i	5417	0.4	0.2	0.5	.	.	0.0	0.1	0.6	1.6	2.0
		p	12887	0.9	0.4	1.3	.	.	0.0	0.2	1.2	3.4	6.4
119	irritable bowel syndrome	i	58799	4.0	2.2	5.8	0.1	2.0	4.8	4.5	4.9	5.0	3.0
		p	90079	6.1	3.3	8.9	0.1	2.2	6.7	6.8	8.0	8.3	6.5

D DIGESTIVE

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
095	i	5.4	8.0	9.6	14.9	1.8	2.8	2.4	2.4	0.6	0.7	0.2	0.5	0.0	0.2
	p	5.5	8.2	10.1	15.8	1.9	2.9	2.7	2.6	0.6	0.7	0.2	0.6	0.0	0.2
096	i	22.6	20.9	6.2	6.1	7.2	6.2	5.3	5.2	5.1	4.9	2.0	5.5	6.3	8.3
	p	23.8	21.7	6.6	6.4	7.2	6.5	5.4	5.6	5.5	5.5	2.3	6.6	6.5	8.3
097	i	0.0	.	0.1	0.3	0.3	0.3
	p	0.0	0.0	0.5	0.4	1.0	0.6	3.3	1.1
098	i	0.1	0.3	0.2	0.4	1.0	0.3
	p	0.0	.	0.1	0.0	0.9	1.2	2.7	2.8	5.7	5.6
099	i	0.1	.	0.1	0.1	0.0	0.1	0.2	0.1
	p	0.2	.	0.4	0.8	0.7	0.8	2.3	1.9
100	i	0.0	0.4	0.2	.	0.3
	p	0.0	.	0.2	0.1	0.6	0.4	0.9	0.6
101	i	0.2	0.0	0.4	0.4	0.2	0.5
	p	0.1	0.1	0.6	0.4	2.3	1.3	3.4	3.6
102	i	0.2	.	.	0.0	.	.	0.0	0.0	.	0.0	0.1	0.0	0.2	0.1
	p	0.2	.	.	0.0	.	0.0	0.0	0.1	0.0	0.2	0.5	0.3	0.3	0.4
103	i	0.1	0.0	0.1	.	0.1	0.2	0.5	0.3
	p	0.1	0.0	0.3	0.1	0.2	0.4	0.6	1.4
104	i	0.7	0.9	0.1	.	0.0	0.0	0.1	0.0	0.0	0.0	.	0.0	.	.
	p	1.1	1.1	0.2	.	0.0	0.0	0.1	0.0	0.0	0.1	.	0.0	.	.
105	i	0.1	0.1	.	.	0.1	.	.	0.0
	p	0.1	0.1	.	.	0.1	.	0.0	0.0	0.0	.	.	0.0	.	.
106	i	0.5	0.3	0.1	.	.	.	0.0	0.0
	p	0.8	0.7	0.7	0.1	0.0	0.2	0.0	0.1	0.1	0.0	.	0.2	.	.
107	i	1.1	0.7	0.6	0.6	0.9	0.9	0.8	1.4	0.8	1.1	0.9	0.8	0.4	0.9
	p	1.8	1.7	1.5	1.3	1.7	1.3	1.3	2.1	1.1	1.4	0.9	0.8	0.4	1.1
108	i	4.1	5.8	2.8	2.7	1.8	2.6	1.5	1.8	1.7	2.7	1.6	1.9	0.7	2.3
	p	5.3	6.9	3.3	3.6	2.1	3.2	2.1	2.6	1.9	3.9	1.8	2.9	1.7	4.5
109	i	.	.	.	0.1	0.6	0.3	0.4	0.6	1.3	1.0	0.8	1.4	1.2	0.8
	p	0.3	0.1	0.0	0.1	0.7	0.4	0.7	0.7	2.3	2.0	2.0	3.0	4.3	2.9
110	i	0.0	.	.	0.0	0.4	0.4	1.2	0.5	1.7	0.9	0.8	0.7	0.4	.
	p	0.0	.	0.0	0.0	0.6	0.5	3.2	1.2	6.0	2.5	4.2	3.1	2.2	2.3
111	i	.	.	0.1	.	0.1	0.0	0.2	0.1	0.4	0.4	0.2	0.1	1.7	0.3
	p	.	.	0.1	.	0.1	0.0	0.6	0.4	2.1	1.5	1.7	1.0	4.8	3.0
112	i	0.0	0.1	0.1	0.3	0.2	0.0	0.1	0.3	0.4
	p	.	.	.	0.0	0.2	0.0	0.3	0.2	1.3	0.8	0.9	0.6	0.9	0.8
113	i	1.9	1.5	0.9	1.3	3.5	3.2	3.7	3.7	3.6	3.3	4.0	3.5	3.8	3.9
	p	2.0	1.9	1.2	1.3	4.4	3.8	5.5	5.0	8.3	6.1	9.6	6.9	5.6	9.1
114	i	.	0.3	1.2	0.7	0.7	0.6	0.4	0.5	0.1	0.1	0.4	0.3	.	0.0
	p	.	0.3	1.4	0.9	0.7	0.9	0.4	0.5	0.2	0.1	0.4	0.4	0.2	0.1
115	i	2.4	1.5	0.4	0.3	0.4	.	0.6	0.1	1.2	0.0	1.4	0.2	4.1	0.3
	p	3.1	1.9	0.5	0.3	0.7	0.0	0.8	0.1	1.6	0.0	2.9	0.4	5.0	0.4
116	i	0.0	.	.	0.1	0.2	0.6	0.6	0.6	0.8	1.4	2.5	1.7	1.2	2.5
	p	0.0	.	0.2	0.1	0.3	0.7	0.9	1.0	2.2	4.7	5.2	7.3	4.8	11.2
117	i	0.4	0.3	.	.	0.1	.	0.2	0.1	0.3	0.4	1.1	0.3	0.1	0.4
	p	0.4	0.4	.	.	0.1	0.0	0.2	0.1	0.4	0.4	1.7	0.6	0.1	1.0
118	i	0.0	0.1	0.0	0.1	0.6	0.6	0.5	2.5	1.5	2.3
	p	0.0	0.1	0.1	0.3	0.9	1.4	1.8	4.7	3.2	8.1
119	i	0.1	.	1.3	2.6	1.9	7.9	2.3	6.8	3.4	6.4	2.6	6.9	1.4	3.8
	p	0.2	0.1	1.4	3.1	2.7	10.9	3.4	10.4	5.3	10.7	5.0	10.9	3.8	7.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
120	chron. enteritis/colitis ulc.	i	3069	0.2	0.2	0.2	0.1	.	0.1	0.4	0.3	0.1	0.1
		p	14445	1.0	0.8	1.2	0.1	0.0	0.6	1.2	1.6	1.1	2.0
121	anal fissure/perianal abscess	i	11085	0.8	0.9	0.6	0.9	0.3	0.7	1.0	0.9	0.5	0.3
		p	15971	1.1	1.4	0.8	1.3	0.4	0.8	1.4	1.5	0.7	0.4
122	hepatomegaly	i	211	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	.
		p	503	0.0	0.0	0.0	.	.	.	0.0	0.1	0.1	0.1
123	cirrhosis/liver disease NEC	i	2708	0.2	0.3	0.1	0.1	0.1	0.2	0.2	0.2	0.3	0.4
		p	10172	0.7	0.8	0.6	0.1	0.1	0.4	0.5	1.2	1.7	1.3
124	diseases of gall-bladder/biliary duct	i	8326	0.6	0.3	0.8	0.0	0.0	0.1	0.4	1.1	1.6	1.6
		p	19698	1.3	0.7	1.9	0.0	0.0	0.1	0.8	2.5	4.3	5.0
125	diseases digestive system NEC	i	7293	0.5	0.4	0.6	0.4	0.1	0.1	0.5	0.6	1.1	1.8
		p	17297	1.2	0.9	1.5	0.5	0.4	0.2	0.8	1.8	3.0	4.0

D DIGESTIVE

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
120	i	0.1	.	.	.	0.0	0.1	0.4	0.3	0.1	0.6	0.2	0.1	0.1	0.2
	p	0.1	.	.	0.0	0.2	1.1	1.1	1.3	1.5	1.7	0.7	1.3	1.2	2.4
121	i	0.9	1.0	0.5	0.2	0.6	0.9	1.3	0.7	1.0	0.7	0.8	0.2	.	0.5
	p	1.2	1.4	0.6	0.3	0.7	1.0	2.0	0.9	1.9	1.0	1.2	0.3	0.1	0.5
122	i	0.0	.	.	0.1	0.1	0.0	.	.
	p	0.0	0.1	.	0.1	0.1	0.0	.	0.1
123	i	.	0.2	0.0	0.1	0.3	0.1	0.3	0.1	0.4	0.0	0.3	0.3	0.1	0.5
	p	.	0.2	0.1	0.1	0.7	0.1	0.6	0.4	1.8	0.6	1.4	2.0	0.6	1.8
124	i	.	0.0	.	0.1	.	0.1	0.3	0.6	0.5	1.7	1.4	1.9	1.7	1.5
	p	.	0.0	.	0.1	.	0.2	0.4	1.1	1.1	3.9	3.3	5.1	4.0	5.6
125	i	0.4	0.3	0.0	0.3	0.1	0.2	0.3	0.6	0.5	0.7	0.4	1.6	2.0	1.6
	p	0.7	0.3	0.0	0.7	0.2	0.3	0.7	1.0	1.5	2.2	2.4	3.5	3.6	4.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
126	complaints of vision	i	12422	0.8	0.7	0.9	0.3	0.9	0.6	0.8	1.0	1.2	1.4
		p	35586	2.4	2.0	2.8	1.6	2.9	1.3	1.6	3.0	4.4	5.8
127	fear of eye disease	i	355	0.0	0.0	0.0	.	0.0	0.1	0.0	0.0	0.0	.
		p	472	0.0	0.0	0.0	.	0.0	0.1	0.0	0.0	0.0	0.0
128	disability/impairment eye	i	19	0.0	.	0.0	.	.	0.0
		p	515	0.0	0.0	0.0	.	0.0	0.0	0.0	0.1	0.0	0.0
129	sympt./complt. eye NEC	i	15488	1.1	0.9	1.2	0.9	0.7	1.1	1.1	1.1	1.7	1.1
		p	34260	2.3	1.8	2.8	1.6	1.5	1.7	1.9	2.8	4.1	6.0
130	infectious conjunctivitis	i	26574	1.8	1.6	2.0	4.6	1.8	2.2	1.5	1.3	1.5	1.7
		p	28810	2.0	1.7	2.2	5.2	1.9	2.2	1.6	1.5	1.8	2.2
131	allergic conjunctivitis	i	79393	5.4	4.8	6.0	7.2	5.2	6.3	5.1	4.9	5.1	5.3
		p	88535	6.0	5.2	6.9	7.3	5.4	6.7	5.5	5.7	6.4	7.8
132	blepharitis/stye/chalazion	i	18930	1.3	1.3	1.2	1.2	1.0	1.6	1.3	1.4	0.6	1.2
		p	21661	1.5	1.5	1.4	1.2	1.1	1.8	1.5	1.6	1.1	1.4
133	infection of eye NEC	i	7033	0.5	0.4	0.6	0.5	0.3	0.6	0.5	0.6	0.4	0.4
		p	13761	0.9	0.8	1.1	1.0	0.5	0.8	0.8	0.9	1.8	2.2
134	corneal ulcer	i	664	0.0	0.1	0.0	.	.	.	0.1	0.1	.	.
		p	814	0.1	0.1	0.0	.	.	0.0	0.1	0.1	0.0	.
135	neoplasms of eye/adnexa	i	424	0.0	0.0	0.0	.	0.0	0.0	0.0	0.1	0.1	.
		p	1392	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.1	0.5	0.0
136	contusion/abrasion/black eye	i	9552	0.7	0.8	0.5	0.4	0.5	0.4	0.9	0.6	0.3	0.7
		p	10880	0.7	0.9	0.6	0.4	0.6	0.5	1.1	0.7	0.5	0.9
137	foreign body in eye	i	20885	1.4	2.6	0.3	0.5	0.5	1.9	2.4	0.9	0.4	0.3
		p	23468	1.6	2.9	0.4	0.5	0.5	2.2	2.8	1.0	0.4	0.6
138	injuries eye NEC	i	2902	0.2	0.3	0.1	0.3	0.3	0.1	0.1	0.3	0.3	0.1
		p	4174	0.3	0.4	0.2	0.3	0.3	0.3	0.2	0.3	0.4	0.2
139	blocked lacrimal duct of infants	i	543	0.0	0.0	0.0	0.4
		p	701	0.0	0.1	0.0	0.5
140	cong. anom. of eye NEC	i	352	0.0	0.0	0.0	0.1	0.0
		p	751	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.0	.	.
141	refractive errors	i	14155	1.0	0.8	1.1	0.3	1.4	1.2	0.7	1.3	0.9	0.4
		p	64538	4.4	3.6	5.1	1.1	5.3	3.9	3.3	6.2	6.1	4.8
142	cataract	i	2640	0.2	0.1	0.2	.	.	0.0	0.0	0.1	0.8	1.7
		p	17889	1.2	0.7	1.7	.	0.0	0.0	0.1	0.7	5.1	12.8
143	glaucoma	i	2740	0.2	0.1	0.2	.	.	0.0	0.1	0.3	0.8	0.9
		p	19709	1.3	1.0	1.7	.	0.1	0.1	0.2	1.5	5.5	10.7
144	blindness, all types	i	124	0.0	.	0.0	.	.	0.0	.	.	0.0	0.1
		p	3393	0.2	0.2	0.3	.	.	0.1	0.1	0.1	0.7	2.4
145	strabismus	i	2182	0.1	0.1	0.2	1.5	0.2	0.1	0.0	0.0	.	.
		p	6301	0.4	0.4	0.4	3.3	1.3	0.1	0.1	0.1	0.0	.
146	retinopathy	i	2359	0.2	0.1	0.2	.	.	0.0	0.1	0.2	0.7	0.8
		p	6948	0.5	0.4	0.6	0.0	.	0.1	0.2	0.5	2.6	2.4
147	diseases of eye NEC	i	9930	0.7	0.6	0.7	0.4	0.2	0.4	0.7	0.9	1.5	0.9
		p	18321	1.2	1.1	1.4	0.7	0.5	0.7	0.9	1.6	3.0	3.9

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
126	i	0.2	0.4	0.9	0.9	0.5	0.8	0.7	1.0	1.1	0.9	0.8	1.4	1.1	1.5
	p	1.8	1.4	2.8	2.9	1.1	1.4	1.4	1.9	2.9	3.2	3.1	5.3	3.1	7.2
127	i	.	.	.	0.1	0.0	0.1	0.0	0.0	0.0	.	.	0.0	.	.
	p	.	.	.	0.1	0.0	0.1	0.0	0.0	0.0	.	.	0.0	0.0	.
128	i	0.0
	p	.	.	.	0.1	.	0.0	0.0	0.0	0.0	0.1	.	0.1	0.0	0.0
129	i	1.0	0.8	0.5	0.9	1.3	0.9	0.8	1.4	0.8	1.3	1.1	2.1	1.3	0.9
	p	1.7	1.4	1.1	1.9	1.7	1.7	1.5	2.3	2.3	3.2	2.7	5.2	3.8	7.1
130	i	4.9	4.2	1.4	2.1	1.5	2.9	1.2	1.9	1.5	1.1	1.5	1.6	1.4	1.8
	p	5.0	5.3	1.6	2.1	1.5	2.9	1.3	1.9	1.8	1.2	1.9	1.7	1.4	2.6
131	i	7.3	7.0	4.4	6.1	4.9	7.7	4.2	6.0	5.2	4.7	4.4	5.7	4.4	5.8
	p	7.5	7.2	4.3	6.5	5.0	8.5	4.4	6.7	6.1	5.3	5.5	7.1	6.2	8.7
132	i	0.9	1.4	0.9	1.1	1.6	1.5	1.1	1.6	2.0	0.9	1.0	0.3	1.4	1.1
	p	0.9	1.5	1.1	1.2	1.8	1.8	1.3	1.8	2.1	1.0	1.4	0.8	1.4	1.3
133	i	0.3	0.7	0.2	0.4	0.4	0.7	0.4	0.5	0.5	0.6	0.2	0.6	0.1	0.7
	p	0.6	1.5	0.5	0.5	0.7	1.0	0.6	0.9	0.9	0.9	2.4	1.3	1.3	2.7
134	i	0.1	0.1	0.1
	p	0.0	0.1	0.1	0.1	0.0	.	0.0	.	.
135	i	.	.	0.0	0.0	0.0	0.0	0.0	0.0	0.1	0.1	.	0.1	.	.
	p	0.1	.	0.0	0.1	0.1	0.0	0.0	0.1	0.1	0.1	0.3	0.6	0.1	.
136	i	0.2	0.6	0.9	0.2	0.6	0.2	1.1	0.8	0.6	0.6	0.2	0.5	0.7	0.8
	p	0.2	0.6	1.0	0.2	0.7	0.2	1.2	0.9	0.8	0.6	0.2	0.7	0.9	0.9
137	i	0.6	0.4	0.7	0.3	3.4	0.4	4.4	0.4	1.8	0.1	0.5	0.3	0.6	0.1
	p	0.6	0.4	0.7	0.3	3.8	0.5	4.9	0.5	1.8	0.2	0.5	0.3	1.2	0.2
138	i	0.3	0.3	0.4	0.2	0.2	0.1	0.2	0.1	0.5	0.1	0.4	0.2	.	0.1
	p	0.3	0.4	0.4	0.2	0.4	0.1	0.3	0.2	0.5	0.1	0.4	0.4	.	0.3
139	i	0.3	0.4
	p	0.4	0.7
140	i	0.1	0.0	0.0	0.1
	p	0.2	0.1	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
141	i	0.2	0.5	0.9	1.8	1.3	1.1	0.5	0.9	0.9	1.7	0.6	1.2	0.8	0.2
	p	0.6	1.5	4.7	5.9	3.6	4.1	2.5	4.1	5.0	7.5	5.1	6.9	4.5	4.9
142	i	0.0	0.0	0.0	0.1	0.2	1.1	0.6	0.6	2.3
	p	.	.	.	0.1	.	0.0	0.1	0.0	0.7	0.8	4.5	5.6	7.6	15.6
143	i	0.0	.	0.0	0.1	0.3	0.3	0.6	0.9	0.9	1.0
	p	.	.	0.1	0.1	0.0	0.2	0.2	0.2	1.3	1.7	4.9	5.9	10.2	10.9
144	i	0.0	0.0	.	0.2
	p	0.1	0.0	0.1	0.0	0.1	0.1	0.3	1.0	1.4	3.0
145	i	1.5	1.6	0.1	0.3	0.1	0.1	0.0	0.0	0.0	0.0
	p	3.3	3.3	1.2	1.3	0.1	0.2	0.1	0.2	0.1	0.0	.	0.0	.	.
146	i	0.1	0.1	0.1	0.2	0.2	0.7	0.8	0.4	0.9
	p	.	0.1	.	.	0.1	0.1	0.2	0.1	0.5	0.5	2.2	2.9	2.4	2.4
147	i	0.7	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	0.9	1.7	1.4	1.2	0.8
	p	0.8	0.6	0.5	0.5	0.7	0.7	0.6	1.2	1.6	1.5	3.1	2.9	3.2	4.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
148	earache	i	3738	0.3	0.2	0.3	1.1	0.4	0.2	0.2	0.2	0.3	0.1
		p	8693	0.6	0.5	0.7	2.4	0.9	0.3	0.4	0.4	0.7	0.3
149	hearing complt.	i	2441	0.2	0.2	0.2	0.4	0.1	0.1	0.2	0.2	0.1	0.1
		p	6419	0.4	0.4	0.4	1.0	0.6	0.1	0.4	0.4	0.2	1.0
150	ringing/buzzing/tinnitus	i	2797	0.2	0.2	0.2	.	.	0.0	0.2	0.3	0.9	0.3
		p	4244	0.3	0.3	0.2	.	.	0.0	0.2	0.5	1.0	0.6
151	blood/discharge from ear	i	1763	0.1	0.2	0.1	0.8	0.3	0.0	0.0	0.1	0.1	.
		p	3614	0.2	0.3	0.2	1.1	0.7	0.0	0.1	0.2	0.1	0.1
152	fear of ear disease	i	26	0.0	0.0	.	.	.	0.0	.	.	.	0.0
		p	277	0.0	0.0	0.0	0.2	0.0	.	0.0	.	.	0.0
153	disability/impairment of ear	i	26	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	33	0.0	.	0.0	.	.	.	0.0	.	.	.
154	symp./complt. of ear NEC	i	4018	0.3	0.2	0.3	0.6	0.5	0.3	0.3	0.1	0.1	0.2
		p	6537	0.4	0.5	0.4	0.8	0.7	0.3	0.4	0.4	0.5	0.5
155	otitis externa	i	50202	3.4	3.7	3.2	2.8	3.0	2.2	4.3	4.0	3.4	1.8
		p	67925	4.6	4.9	4.4	3.8	3.7	3.1	5.4	5.6	5.1	3.6
156	acute otitis media/myringitis	i	97708	6.7	6.6	6.7	51.2	16.8	2.8	1.7	1.4	1.1	0.7
		p	102157	7.0	6.8	7.1	52.2	17.6	2.9	1.9	1.5	1.3	0.9
157	serous otitis media/glue ear	i	21105	1.4	1.5	1.4	8.4	3.4	0.9	0.7	0.4	0.5	0.3
		p	29633	2.0	2.0	2.0	10.6	6.2	1.0	0.8	0.5	0.5	0.5
158	eustachian tube dysfunction	i	32787	2.2	1.9	2.6	3.4	3.3	2.1	2.3	1.7	1.4	1.1
		p	40382	2.8	2.3	3.2	4.5	4.1	2.5	2.9	2.1	1.7	1.6
159	infections of ear NEC	i	4944	0.3	0.3	0.4	1.0	0.5	0.3	0.3	0.3	0.2	0.2
		p	9367	0.6	0.6	0.7	1.6	1.3	0.5	0.5	0.5	0.3	0.2
160	neoplasm of ear	i	256	0.0	0.0	0.0	.	0.0	0.0	.	.	.	0.1
		p	423	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	0.1	0.1
161	foreign body in ear	i	1806	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.1	0.1	0.3
		p	2083	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.3
162	perforation tymp. membrane	i	2256	0.2	0.2	0.1	0.3	0.3	0.2	0.2	0.0	0.1	0.0
		p	4702	0.3	0.3	0.3	0.5	0.5	0.4	0.3	0.1	0.3	0.1
163	superficial injury of ear	i	470	0.0	0.0	0.0	0.0	0.1	0.0	0.0	.	.	0.2
		p	679	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.0	.	0.2
164	injuries of ear NEC	i	890	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.0	.
		p	1061	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.1
165	cong. anom. of ear	i	416	0.0	0.0	0.0	0.1	0.1	0.0	.	0.0	.	.
		p	638	0.0	0.1	0.0	0.2	0.1	0.0	0.0	0.0	.	.
166	ear wax (excessive)	i	121818	8.3	8.4	8.2	3.3	2.6	5.1	9.0	10.8	13.9	16.8
		p	135848	9.3	9.2	9.3	3.6	2.8	5.3	9.7	12.4	15.9	20.4
167	vertiginous syndromes	i	8671	0.6	0.5	0.7	.	.	0.1	0.4	1.2	2.0	1.5
		p	16608	1.1	0.8	1.4	.	.	0.1	0.6	2.0	3.8	4.9
168	otosclerosis	i	701	0.0	0.0	0.1	.	.	0.0	0.0	0.0	0.3	0.1
		p	1337	0.1	0.1	0.1	.	.	0.1	0.1	0.1	0.3	0.3
169	presbycusis	i	5346	0.4	0.4	0.4	0.3	1.7	3.4
		p	8050	0.5	0.5	0.6	0.5	2.6	5.1
170	acoustic trauma	i	1188	0.1	0.1	0.0	.	.	0.0	0.1	0.2	0.0	.
		p	1310	0.1	0.2	0.0	.	.	0.0	0.1	0.2	0.0	.
171	deafness NEC	i	4539	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.4	0.2	0.0
		p	10525	0.7	0.7	0.7	1.1	0.7	0.5	0.6	0.7	0.8	1.8
172	disease of ear/mastoid NEC	i	4084	0.3	0.3	0.3	0.2	0.3	0.2	0.3	0.4	0.4	0.2
		p	6563	0.4	0.5	0.4	0.4	0.4	0.3	0.4	0.6	0.6	0.5

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
148	i	1.2	0.9	0.2	0.6	0.1	0.2	0.0	0.3	0.1	0.3	0.3	0.3	0.1	0.1
	p	2.4	2.4	0.5	1.3	0.3	0.3	0.4	0.5	0.4	0.5	0.7	0.7	0.1	0.5
149	i	0.3	0.6	0.1	0.2	0.0	0.1	0.2	0.1	0.2	0.2	0.0	0.1	0.4	0.0
	p	1.3	0.8	0.5	0.8	0.1	0.1	0.4	0.3	0.5	0.4	0.1	0.3	0.8	1.1
150	i	0.0	0.2	0.1	0.3	0.3	1.1	0.7	0.5	0.2
	p	0.0	0.0	0.3	0.1	0.6	0.5	1.1	1.0	1.1	0.3
151	i	0.9	0.7	0.4	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	.	.	.
	p	1.3	0.8	1.0	0.4	0.1	0.0	0.1	0.1	0.3	0.1	0.1	0.2	0.2	0.1
152	i	0.0	0.0	.
	p	0.2	0.1	0.0	.	.	.	0.0	0.0	0.0	.
153	i	0.0
	p	0.0
154	i	0.4	0.8	0.5	0.6	0.1	0.4	0.3	0.2	0.1	0.2	0.1	0.0	0.5	0.1
	p	0.5	1.0	0.6	0.8	0.2	0.5	0.5	0.3	0.4	0.3	0.5	0.4	1.1	0.1
155	i	3.4	2.3	2.8	3.2	2.4	2.0	4.5	4.0	4.2	3.8	4.6	2.4	2.2	1.6
	p	4.5	3.1	3.7	3.7	3.4	2.7	5.2	5.6	5.6	5.6	6.7	3.9	5.6	2.5
156	i	51.8	50.5	15.5	18.0	2.4	3.2	1.5	2.0	1.5	1.3	1.3	1.0	0.6	0.8
	p	52.3	52.2	16.1	19.2	2.6	3.3	1.6	2.2	1.5	1.6	1.6	1.1	0.9	0.9
157	i	9.5	7.3	3.3	3.6	0.7	1.1	0.6	0.8	0.4	0.3	0.8	0.3	0.7	0.2
	p	12.8	8.3	5.2	7.2	0.8	1.2	0.7	0.9	0.5	0.5	0.8	0.3	0.9	0.3
158	i	2.3	4.4	3.2	3.4	1.3	3.0	2.0	2.7	1.7	1.8	1.2	1.6	1.1	1.2
	p	3.4	5.6	4.1	4.1	1.5	3.5	2.3	3.4	1.9	2.4	1.4	1.9	1.3	1.7
159	i	0.8	1.3	0.3	0.7	0.2	0.3	0.4	0.2	0.3	0.3	0.1	0.3	0.3	0.1
	p	1.5	1.8	1.2	1.5	0.5	0.5	0.5	0.5	0.4	0.5	0.1	0.5	0.3	0.1
160	i	.	.	.	0.1	0.1	0.2	.
	p	.	.	.	0.1	0.1	.	.	0.0	0.0	.	0.2	.	0.2	.
161	i	0.1	0.1	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.3	.	0.6	0.1
	p	0.1	0.1	0.2	0.3	0.1	0.1	0.1	0.1	0.2	0.1	0.3	.	0.6	0.1
162	i	0.1	0.4	0.4	0.1	0.1	0.2	0.3	0.1	.	0.0	0.2	0.1	.	0.0
	p	0.2	0.8	0.5	0.4	0.3	0.5	0.4	0.3	0.0	0.2	0.2	0.3	0.4	0.0
163	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.3
	p	0.0	0.0	0.1	0.1	0.0	0.1	0.0	0.0	.	0.1	.	.	.	0.3
164	i	.	0.0	0.1	.	0.0	0.1	0.1	0.1	0.1	0.0	0.0	.	.	.
	p	.	0.0	0.1	.	0.0	0.1	0.1	0.1	0.2	0.0	0.0	.	.	0.1
165	i	0.3	.	0.1	0.1	.	0.0	.	.	0.0
	p	0.3	.	0.1	0.1	.	0.0	0.0	.	0.1
166	i	3.2	3.4	2.0	3.2	4.6	5.6	9.3	8.8	12.2	9.4	15.6	12.5	17.6	16.4
	p	3.4	3.8	1.9	3.6	4.7	5.8	9.8	9.5	13.5	11.4	17.2	14.9	22.8	19.2
167	i	0.1	0.1	0.3	0.5	1.0	1.4	1.7	2.1	1.5	1.5
	p	0.1	0.1	0.5	0.7	1.7	2.3	3.1	4.3	3.2	5.8
168	i	0.1	0.1	0.0	0.0	0.0	0.0	0.4	0.2	0.1
	p	0.0	0.1	0.1	0.1	0.0	0.1	0.1	0.4	0.2	0.3
169	i	0.4	0.2	1.7	1.7	4.3	2.9
	p	0.6	0.3	2.4	2.8	6.8	4.2
170	i	0.1	.	0.1	0.1	0.4	0.0	0.1	.	.	.
	p	0.1	.	0.1	0.1	0.5	0.0	0.1	.	.	.
171	i	0.2	0.7	0.2	0.4	0.5	0.2	0.4	0.2	0.4	0.5	0.4	.	.	0.0
	p	0.8	1.4	0.5	0.8	0.7	0.3	0.7	0.4	0.5	0.9	1.3	0.5	1.7	1.9
172	i	0.3	0.2	0.3	0.3	0.2	0.1	0.3	0.2	0.3	0.5	0.5	0.3	0.3	0.2
	p	0.4	0.4	0.4	0.4	0.3	0.2	0.5	0.3	0.6	0.7	0.5	0.7	0.6	0.4

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals abs.	/1000 pers.	sex		age						
					male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
173	complaints of heart	i	7180	0.5	0.3	0.6	.	0.1	0.3	0.4	0.9	1.1	0.6
		p	17606	1.2	0.9	1.5	0.0	0.1	0.3	0.9	2.7	2.7	2.0
174	edema of extremities	i	12452	0.8	0.4	1.3	0.2	0.0	0.1	0.6	1.1	2.1	4.6
		p	68683	4.7	2.0	7.3	0.3	0.1	0.2	1.6	6.3	13.5	35.3
175	fear of dis. circ. system	i	7513	0.5	0.5	0.5	.	.	0.1	0.7	1.0	0.6	0.7
		p	13212	0.9	0.8	1.0	.	.	0.1	1.1	1.6	1.7	1.6
176	disability/impairment circ. syst. NEC	i	0
		p	130	0.0	0.0	0.0	0.1	0.1
177	sympt./complt. circ. syst. NEC	i	2328	0.2	0.2	0.2	0.0	.	0.1	0.1	0.3	0.2	0.7
		p	5136	0.4	0.3	0.4	0.0	0.1	0.1	0.2	0.5	1.1	1.5
178	infections dis. circ. syst.	i	205	0.0	0.0	0.0	.	.	0.0	0.0	0.0	0.0	.
		p	1161	0.1	0.1	0.1	.	.	0.1	0.0	0.2	0.1	0.2
179	neoplasm circ. syst.	i	374	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	0.1	.
		p	756	0.1	0.0	0.1	.	0.0	0.1	0.1	0.0	0.1	.
180	injuries heart/circ. syst.	i	27	0.0	0.0	0.0	.	0.0	0.0
		p	310	0.0	0.0	0.0	.	0.0	.	.	0.0	0.2	0.0
181	cong. anom. heart/circ. syst.	i	249	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	.
		p	4083	0.3	0.2	0.4	0.7	0.4	0.4	0.1	0.1	0.5	0.2
182	rheumatic fever/heart dis.	i	292	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	.	.
		p	1331	0.1	0.1	0.1	.	0.1	0.2	0.1	0.1	.	0.2
183	acute myocardial infarction	i	6581	0.5	0.5	0.4	.	.	.	0.1	0.8	2.0	2.6
		p	0
184	angina pectoris	i	18736	1.3	1.3	1.3	.	.	0.0	0.5	2.2	4.6	6.4
		p	116735	8.0	7.8	8.1	.	.	0.1	0.9	12.5	35.3	49.6
185	healed myocardial infarction	i	0
		p	70603	4.8	6.9	2.8	.	.	.	0.7	9.5	21.1	21.6
186	chron. ischemic heart dis. NEC	i	2403	0.2	0.1	0.2	.	.	.	0.0	0.4	0.3	0.9
		p	73762	5.0	6.1	4.0	.	.	.	0.3	8.6	23.3	28.2
187	heart failure	i	12593	0.9	0.7	1.0	0.0	.	0.0	0.1	0.3	2.8	10.4
		p	93835	6.4	4.9	7.8	0.0	0.1	0.1	0.3	3.0	21.6	78.4
188	atrial fibrillation/flutter	i	2977	0.2	0.1	0.3	.	.	.	0.1	0.2	1.1	1.3
		p	34835	2.4	2.1	2.7	.	.	.	0.2	1.7	11.0	22.0
189	paroxysmal tachycardia	i	3551	0.2	0.2	0.3	.	0.1	0.2	0.3	0.4	0.4	0.1
		p	9678	0.7	0.4	0.9	.	0.1	0.2	0.5	1.3	1.9	1.3
190	ectopic beats, all types	i	2569	0.2	0.2	0.2	0.0	.	0.0	0.2	0.3	0.2	0.6
		p	6017	0.4	0.3	0.5	0.0	0.0	0.0	0.2	0.9	0.9	1.7
191	heart murmur NOS	i	965	0.1	0.1	0.1	0.6	0.1	0.1	0.0	.	0.1	.
		p	2657	0.2	0.2	0.1	1.2	0.4	0.1	0.1	0.0	0.1	.
192	pulmonary heart dis.	i	108	0.0	0.0	0.1	0.1
		p	1747	0.1	0.1	0.1	.	.	.	0.0	0.2	0.5	0.9
193	heart valve dis. NOS, non-rheum.	i	496	0.0	0.0	0.0	.	0.0	0.0	0.0	0.1	0.1	0.1
		p	10557	0.7	0.6	0.8	.	0.1	0.2	0.3	1.1	2.5	3.2
194	heart disease NEC	i	6194	0.4	0.3	0.5	.	0.1	0.0	0.2	0.6	1.0	2.8
		p	40793	2.8	2.5	3.1	0.1	0.1	0.2	0.9	3.7	10.1	18.4
195	elevated blood pressure w.o. hypert.	i	2833	0.2	0.2	0.2	.	0.0	0.0	0.2	0.4	0.4	0.1
		p	20697	1.4	1.0	1.8	.	0.0	0.3	1.0	2.8	4.7	3.1
196	uncomplicated hypertension	i	45815	3.1	2.5	3.8	.	0.0	0.3	1.7	6.1	10.8	9.8
		p	658779	44.9	32.9	56.5	.	0.3	2.4	16.0	96.5	166.3	151.4
197	hypertension with invol.targ. organs	i	5972	0.4	0.4	0.4	.	.	.	0.0	0.5	2.0	2.9
		p	103972	7.1	6.3	7.8	.	.	0.1	0.6	8.3	33.3	53.8

K CIRCULATORY

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
173	i	.	.	.	0.3	0.2	0.4	0.3	0.5	0.6	1.2	1.0	1.1	0.3	0.7
	p	.	0.1	.	0.3	0.2	0.5	0.9	0.9	2.0	3.4	2.7	2.7	1.3	2.4
174	i	0.4	.	0.0	0.1	0.1	0.1	0.2	1.1	0.5	1.7	1.5	2.6	2.2	5.9
	p	0.4	0.1	0.1	0.1	0.2	0.2	0.4	3.0	2.2	10.4	7.9	17.9	23.0	41.8
175	i	0.1	0.2	0.9	0.5	0.9	1.0	0.6	0.7	0.8	0.7
	p	0.1	0.2	1.3	0.9	1.2	2.0	1.2	2.0	1.5	1.7
176	i
	p	0.1	.	.	0.1
177	i	0.0	0.0	.	.	0.0	0.1	0.2	0.0	0.2	0.3	0.3	0.1	0.3	0.9
	p	0.0	0.0	.	0.1	0.1	0.2	0.3	0.2	0.5	0.4	1.0	1.2	1.4	1.5
178	i	0.0	0.0	.	.	0.1	.	0.1	.	.
	p	0.1	0.1	0.1	0.0	0.1	0.3	.	0.1	.	0.2
179	i	.	.	0.1	.	0.0	.	.	0.1	.	0.0	0.0	0.1	.	.
	p	.	.	0.1	0.0	0.0	0.1	.	0.1	.	0.1	0.0	0.2	.	.
180	i	.	.	0.0	0.0
	p	.	.	0.0	0.1	0.0	0.4	.	0.0	0.0
181	i	0.0	0.0	0.0	.	.	0.0	.	0.1	.	0.0	.	0.0	.	.
	p	0.9	0.6	0.3	0.5	0.3	0.4	0.0	0.3	0.1	0.2	0.2	0.8	0.3	0.1
182	i	.	.	.	0.0	.	0.0	0.0	.	0.1
	p	.	.	0.2	0.0	0.3	0.1	0.1	0.0	0.1	0.0	.	.	0.0	0.3
183	i	0.1	0.0	1.2	0.3	2.2	1.7	2.9	2.4
	p
184	i	0.0	0.7	0.2	2.6	1.9	4.5	4.7	6.1	6.6
	p	0.0	0.1	1.2	0.5	15.3	9.7	39.9	31.7	48.3	50.3
185	i
	p	1.3	0.1	15.5	3.6	34.5	10.7	31.4	16.5
186	i	0.0	.	0.4	0.5	0.5	0.2	0.3	1.2
	p	0.4	0.2	13.0	4.3	33.3	15.6	32.7	25.8
187	i	.	0.0	.	.	0.0	0.0	0.1	0.1	0.3	0.3	3.6	2.3	9.9	10.7
	p	.	0.0	0.1	0.1	0.1	0.1	0.3	0.3	3.0	3.1	25.6	18.5	69.7	82.9
188	i	0.1	0.1	0.1	0.2	0.6	1.5	1.1	1.3
	p	0.3	0.1	1.9	1.6	11.1	10.9	23.2	21.4
189	i	.	.	.	0.1	0.1	0.2	0.2	0.3	0.3	0.6	0.5	0.2	0.2	0.1
	p	.	.	0.0	0.1	0.1	0.3	0.3	0.6	0.8	1.8	1.6	2.2	1.1	1.5
190	i	.	0.0	.	.	0.0	0.0	0.2	0.1	0.3	0.4	0.2	0.3	0.8	0.5
	p	.	0.0	.	0.0	0.0	0.0	0.2	0.2	0.6	1.2	0.8	1.0	2.0	1.5
191	i	0.8	0.4	0.1	0.1	0.1	0.0	0.0	0.1	.	.
	p	1.9	0.5	0.5	0.3	0.1	0.1	0.1	0.1	.	0.1	.	0.2	.	.
192	i	0.1	.	0.1	.
	p	0.0	0.2	0.1	0.1	0.7	0.3	1.1	0.8
193	i	.	.	0.0	.	.	0.0	.	0.0	0.1	0.1	.	0.3	0.1	0.1
	p	.	.	0.1	.	0.2	0.3	0.3	0.4	1.0	1.1	2.5	2.6	2.3	3.7
194	i	.	.	0.1	0.1	0.1	.	0.1	0.3	0.5	0.7	0.9	1.2	3.3	2.5
	p	.	0.1	0.1	0.2	0.3	0.1	1.1	0.7	3.4	4.0	9.9	10.2	19.3	17.9
195	i	.	.	0.0	.	0.1	0.0	0.3	0.2	0.3	0.5	0.4	0.4	0.1	0.1
	p	.	.	0.0	.	0.2	0.4	1.0	0.9	2.0	3.5	2.9	6.1	1.8	3.7
196	i	.	.	0.0	0.0	0.3	0.2	2.0	1.4	5.1	7.1	7.2	13.5	6.9	11.4
	p	.	.	0.2	0.4	2.1	2.7	16.9	15.0	76.9	115.6	120.1	201.9	100.5	178.5
197	i	0.0	0.0	0.6	0.4	2.2	1.9	2.9	2.9
	p	0.0	0.1	0.6	0.6	9.5	7.0	37.3	30.1	49.0	56.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
198	(postural) hypotension	i	10216	0.7	0.5	0.9	.	.	0.9	0.6	0.6	1.2	2.3
		p	16173	1.1	0.8	1.4	.	.	1.2	0.9	1.0	2.2	3.6
199	transient cerebral ischemia	i	8798	0.6	0.6	0.6	.	.	0.0	0.0	0.5	2.6	5.8
		p	24136	1.6	1.6	1.7	.	.	0.0	0.1	1.5	8.1	14.0
200	cerebrovascular diseases	i	9886	0.7	0.5	0.9	0.0	.	0.0	0.1	0.8	2.5	5.5
		p	46526	3.2	3.0	3.3	0.1	0.0	0.0	0.3	3.3	12.6	28.4
201	atherosclerosis, excl. heart/brain	i	1609	0.1	0.1	0.1	.	.	.	0.0	0.1	0.9	0.5
		p	13390	0.9	1.0	0.8	.	.	0.0	0.1	1.0	5.0	6.2
202	arterial vascular diseases NEC	i	8746	0.6	0.7	0.5	.	.	0.2	0.2	1.2	2.0	2.6
		p	45719	3.1	3.6	2.6	.	.	0.3	0.6	4.2	14.0	19.8
203	pulmonary embolism	i	1186	0.1	0.0	0.1	.	.	0.0	0.0	0.1	0.2	0.4
		p	5000	0.3	0.2	0.5	.	.	0.0	0.2	0.3	1.5	2.2
204	(thrombo) phlebitis	i	6454	0.4	0.3	0.6	.	.	0.1	0.2	0.9	1.7	1.1
		p	10556	0.7	0.4	1.0	.	.	0.1	0.3	1.4	2.8	2.4
205	venous thrombosis	i	2795	0.2	0.2	0.2	.	.	0.0	0.1	0.3	0.5	0.9
		p	15854	1.1	1.0	1.2	.	.	0.1	0.4	1.6	4.7	5.0
206	varicous veins of legs	i	18747	1.3	0.5	2.0	.	0.1	0.6	1.1	2.3	2.3	3.7
		p	52605	3.6	1.5	5.6	.	0.1	0.8	2.5	5.7	10.2	14.7
207	various ulcer of skin	i	4381	0.3	0.2	0.4	.	.	.	0.0	0.3	1.1	2.6
		p	16849	1.1	0.6	1.7	.	.	0.0	0.2	1.0	3.8	11.6
208	hemorrhoids	i	24219	1.7	1.6	1.7	.	0.1	0.9	2.4	2.6	1.6	1.5
		p	54279	3.7	3.5	3.9	.	0.2	1.4	4.6	6.2	5.2	6.4
209	dis. circulatory system NEC	i	7883	0.5	0.4	0.7	.	0.1	0.3	0.4	0.7	1.7	1.1
		p	26438	1.8	1.7	1.9	0.1	0.3	0.5	0.9	2.4	6.7	8.0

K CIRCULATORY

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
198	i	0.4	1.3	0.3	0.9	0.4	0.8	1.3	1.2	3.7	1.6
	p	0.5	1.9	0.4	1.5	0.8	1.3	2.4	2.1	5.8	2.4
199	i	0.0	.	0.0	0.0	0.7	0.4	3.0	2.3	6.9	5.2
	p	0.0	0.1	0.1	0.1	2.2	0.7	9.0	7.5	15.6	13.1
200	i	0.1	.	.	.	0.0	0.1	0.0	0.1	0.9	0.8	2.4	2.6	3.4	6.6
	p	0.3	.	.	0.0	0.0	0.1	0.3	0.3	4.4	2.2	15.3	10.5	28.8	28.1
201	i	0.0	0.0	0.1	1.4	0.5	0.2	0.6
	p	0.0	.	0.1	0.1	1.2	0.8	6.9	3.5	8.2	5.2
202	i	0.1	0.3	0.1	0.2	1.4	0.9	3.0	1.2	4.2	1.7
	p	0.1	0.5	0.5	0.7	5.1	3.2	21.3	8.3	28.8	15.0
203	i	0.1	.	0.1	0.0	0.2	0.1	0.3	0.3	0.5
	p	0.1	0.1	0.3	0.2	0.3	1.6	1.4	0.9	2.9
204	i	0.1	0.0	0.1	0.2	0.8	1.0	0.8	2.5	0.2	1.7
	p	0.2	0.0	0.2	0.4	1.2	1.6	0.8	4.3	0.6	3.4
205	i	0.1	0.0	0.1	0.1	0.4	0.3	0.5	0.6	1.3	0.7
	p	0.1	0.2	0.4	0.4	1.7	1.5	4.5	4.9	5.5	4.8
206	i	.	.	0.0	0.1	0.2	1.0	0.3	2.0	1.4	3.1	0.7	3.5	1.2	5.0
	p	.	.	0.0	0.1	0.3	1.3	0.8	4.3	3.2	8.1	3.2	15.5	8.7	17.9
207	i	0.0	0.0	0.3	0.4	1.0	1.3	1.9	3.0
	p	0.0	0.1	0.1	0.2	0.6	1.5	1.9	5.2	8.0	13.5
208	i	.	.	0.2	.	0.9	1.0	2.4	2.4	2.3	2.8	1.9	1.4	1.6	1.4
	p	.	.	0.2	0.1	1.1	1.7	4.6	4.7	6.1	6.2	4.3	6.0	7.3	5.9
209	i	.	.	0.1	0.2	0.2	0.5	0.3	0.6	0.6	0.9	1.3	2.0	1.1	1.1
	p	0.0	0.1	0.3	0.2	0.5	0.5	0.7	1.1	2.5	2.4	7.2	6.4	9.6	7.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex	age							
			abs.	/1000 pers.		male	female	0-4	5-14	15-24	25-44	45-64	65-74
210	neck symp./complt., excl. headache	i	10843	0.7	0.7	0.8	0.1	0.2	0.7	0.8	1.1	1.2	0.6
		p	18693	1.3	1.1	1.5	0.2	0.2	0.9	1.4	2.1	1.7	1.6
211	back symp./complt.	i	19699	1.3	1.3	1.4	.	0.3	1.4	1.8	1.6	1.3	1.9
		p	54072	3.7	3.3	4.1	0.2	0.4	2.3	4.0	6.1	5.2	6.6
212	low back symp./complt. w/o radiation	i	5148	0.4	0.3	0.4	.	0.0	0.4	0.5	0.6	0.0	0.1
		p	8500	0.6	0.4	0.7	.	0.0	0.4	0.8	1.0	0.2	0.3
213	chest symp./complt.	i	12733	0.9	0.9	0.8	0.1	0.2	0.6	1.1	1.4	0.7	0.8
		p	16807	1.1	1.2	1.1	0.1	0.3	0.8	1.4	1.8	1.3	1.5
214	flank sympt.complt.	i	997	0.1	0.1	0.1	.	0.0	0.1	0.1	0.0	0.2	.
		p	1824	0.1	0.1	0.1	0.1	0.0	0.1	0.2	0.0	0.2	0.1
215	axilla sympt./complt.	i	322	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	.
		p	338	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	.
216	jaw sympt./complt.	i	4459	0.3	0.3	0.3	0.1	0.2	0.4	0.3	0.3	0.5	0.4
		p	6908	0.5	0.4	0.5	0.2	0.4	0.5	0.5	0.4	0.7	0.5
217	shoulder sympt./complt.	i	4633	0.3	0.3	0.3	.	0.1	0.2	0.5	0.4	0.3	0.3
		p	10110	0.7	0.7	0.7	.	0.1	0.3	0.8	1.1	1.0	1.5
218	arm sympt./complt.	i	7364	0.5	0.5	0.5	0.1	0.1	0.3	0.7	0.9	0.2	0.5
		p	13556	0.9	0.8	1.1	0.1	0.1	0.4	1.1	1.7	0.9	1.4
219	elbow sympt./complt.	i	1325	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	.	0.0
		p	2307	0.2	0.2	0.1	0.1	0.2	0.2	0.2	0.2	0.0	0.1
220	wrist sympt./complt.	i	1250	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.1	.
		p	2189	0.1	0.1	0.2	0.0	0.2	0.1	0.1	0.2	0.2	0.0
221	hand/finger sympt./complt.	i	6963	0.5	0.6	0.4	0.3	0.3	0.5	0.6	0.5	0.4	0.3
		p	10792	0.7	0.8	0.7	0.4	0.5	0.7	0.8	0.8	0.9	0.7
222	hip sympt./complt.	i	2826	0.2	0.2	0.2	0.1	0.0	0.1	0.1	0.4	0.3	0.3
		p	8670	0.6	0.4	0.8	0.2	0.1	0.1	0.3	0.9	2.3	2.1
223	leg/thigh sympt./complt.	i	8104	0.6	0.4	0.7	0.5	0.2	0.3	0.4	0.7	0.9	1.7
		p	19986	1.4	1.0	1.7	0.9	0.3	0.6	0.7	2.0	3.5	5.6
224	knee sympt./complt.	i	11036	0.8	0.9	0.6	0.1	0.5	1.3	0.8	0.8	0.4	0.5
		p	19602	1.3	1.4	1.3	0.1	0.6	1.7	1.3	1.5	1.6	2.5
225	ankle sympt./complt.	i	2592	0.2	0.2	0.2	0.1	0.3	0.3	0.2	0.1	0.2	0.0
		p	4278	0.3	0.3	0.3	0.1	0.3	0.4	0.3	0.2	0.3	0.7
226	foot/toe sympt./complt.	i	10187	0.7	0.6	0.8	0.1	0.7	1.0	0.6	0.8	0.5	1.1
		p	16338	1.1	1.1	1.2	0.6	0.8	1.3	0.9	1.3	1.2	2.0
227	muscle pain/fibrositis	i	212037	14.4	13.3	15.5	0.7	4.0	13.5	18.6	18.8	16.3	14.3
		p	281779	19.2	16.8	21.5	1.1	4.3	15.5	22.5	26.6	27.2	29.2
228	(mult.) muscle sympt./complt. NEC	i	29939	2.0	1.9	2.2	0.0	1.0	1.6	2.7	2.9	1.9	1.1
		p	43573	3.0	2.7	3.3	0.4	1.2	2.3	3.5	4.8	3.1	2.5
229	(mult.) joints sympt./complt. NEC	i	9403	0.6	0.5	0.7	0.2	0.2	0.4	0.8	1.0	0.6	0.8
		p	22839	1.6	1.0	2.1	0.2	0.3	0.5	1.1	2.7	3.7	4.9
230	limbs sympt./complt. NOS	i	550	0.0	0.0	0.0	.	0.1	.	0.0	0.0	0.1	.
		p	700	0.0	0.1	0.0	.	0.1	.	0.0	0.1	0.1	.
231	fear of dis. musculoskeletal syst.	i	1338	0.1	0.1	0.1	.	0.1	0.1	0.1	0.1	0.1	.
		p	2267	0.2	0.1	0.2	.	0.2	0.1	0.2	0.2	0.1	0.2
232	disability/impairment musc. skel. syst.	i	13635	0.9	0.8	1.1	1.3	1.1	1.5	0.8	0.5	0.8	0.6
		p	29607	2.0	1.8	2.2	1.8	2.5	2.5	1.3	1.4	3.1	5.2
233	sympt./complt. musc. skel. syst.	i	1779	0.1	0.1	0.2	0.0	0.1	0.1	0.1	0.2	0.1	0.0
		p	4016	0.3	0.2	0.3	0.0	0.2	0.3	0.2	0.5	0.6	0.2
234	infections musc. skel. syst.	i	2304	0.2	0.2	0.1	0.3	0.2	0.1	0.1	0.2	0.2	0.2
		p	3877	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.4	0.4	0.2

L MUSCULOSKELETAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
210	i	0.0	0.2	0.1	0.3	0.6	0.8	0.8	0.9	0.9	1.2	1.3	1.1	0.5	0.6
	p	0.1	0.3	0.1	0.3	0.7	1.0	1.2	1.7	1.8	2.5	1.8	1.6	1.7	1.5
211	i	.	.	0.3	0.3	1.5	1.2	1.6	1.9	1.6	1.6	1.0	1.5	1.3	2.2
	p	.	0.4	0.3	0.5	2.6	2.0	3.9	4.1	5.4	6.9	3.9	6.2	4.0	8.0
212	i	.	.	0.0	0.0	0.4	0.4	0.2	0.8	0.5	0.7	0.1	0.0	.	0.2
	p	.	.	0.0	0.0	0.4	0.4	0.5	1.2	0.9	1.2	0.1	0.3	.	0.4
213	i	0.1	.	0.1	0.3	0.6	0.7	1.2	1.0	1.5	1.4	0.9	0.5	0.9	0.7
	p	0.2	.	0.3	0.3	0.9	0.7	1.4	1.3	1.8	1.8	1.6	1.0	1.2	1.7
214	i	.	.	.	0.0	0.2	0.1	0.1	0.1	0.0	0.0	.	0.3	.	.
	p	0.2	.	.	0.0	0.2	0.1	0.1	0.3	0.0	0.0	.	0.3	0.3	0.1
215	i	0.0	0.1	.	0.0	.	0.2	.	.
	p	0.0	0.1	.	0.0	.	0.2	.	.
216	i	0.2	.	0.4	.	0.3	0.5	0.2	0.3	0.2	0.3	0.5	0.5	0.2	0.6
	p	0.2	0.1	0.5	0.2	0.4	0.7	0.4	0.7	0.3	0.5	0.8	0.6	0.3	0.6
217	i	.	.	0.1	0.0	0.3	0.2	0.5	0.4	0.4	0.4	0.3	0.3	0.2	0.4
	p	.	.	0.1	0.0	0.4	0.3	0.8	0.7	1.1	1.2	1.1	0.9	1.1	1.7
218	i	0.2	0.1	.	0.2	0.1	0.4	0.7	0.6	0.8	1.0	0.2	0.2	0.4	0.5
	p	0.2	0.1	0.0	0.2	0.2	0.6	1.2	1.0	1.4	2.1	0.5	1.1	0.5	1.9
219	i	0.1	.	0.1	0.2	0.2	0.1	0.1	0.0	0.2	0.1	.	.	0.1	.
	p	0.1	.	0.1	0.3	0.3	0.1	0.3	0.1	0.3	0.1	0.0	.	0.2	.
220	i	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.2	.	.	.
	p	0.0	0.1	0.1	0.3	0.2	0.1	0.1	0.1	0.2	0.3	0.2	0.1	0.1	.
221	i	0.7	.	0.3	0.4	0.6	0.4	0.6	0.5	0.7	0.3	0.4	0.4	0.5	0.1
	p	0.7	0.2	0.5	0.5	0.8	0.5	0.9	0.8	1.0	0.7	0.9	1.0	0.6	0.8
222	i	0.2	0.0	0.0	0.0	0.1	0.1	0.1	0.2	0.3	0.4	0.2	0.5	0.1	0.4
	p	0.2	0.2	0.0	0.2	0.1	0.1	0.2	0.3	0.6	1.3	1.8	2.6	1.0	2.6
223	i	0.3	0.7	0.3	0.2	0.3	0.4	0.3	0.6	0.6	0.9	0.7	1.0	1.4	1.9
	p	0.6	1.2	0.4	0.2	0.6	0.5	0.6	0.8	1.4	2.7	1.6	4.9	6.0	5.3
224	i	0.2	0.1	0.5	0.6	1.7	0.9	1.1	0.6	0.8	0.7	0.1	0.6	0.2	0.6
	p	0.2	0.1	0.6	0.7	2.1	1.3	1.7	0.8	1.4	1.7	0.2	2.6	1.4	3.1
225	i	0.3	.	0.3	0.4	0.4	0.1	0.2	0.1	.	0.2	.	0.3	.	0.1
	p	0.3	.	0.3	0.4	0.5	0.3	0.4	0.1	0.1	0.3	0.3	0.3	.	1.0
226	i	0.1	0.1	0.6	0.7	1.0	1.1	0.6	0.5	0.7	0.8	0.3	0.7	0.2	1.6
	p	0.7	0.5	0.7	0.8	1.3	1.3	1.1	0.8	1.4	1.2	0.4	1.9	0.8	2.6
227	i	1.2	0.2	3.4	4.5	12.5	14.6	17.3	20.0	17.0	20.5	16.0	16.6	13.1	15.0
	p	1.4	0.8	3.9	4.7	14.0	17.0	20.8	24.3	22.9	30.3	22.8	30.6	23.1	32.5
228	i	0.1	.	0.8	1.2	1.6	1.6	2.6	2.8	2.6	3.2	1.2	2.4	0.1	1.6
	p	0.6	0.3	0.9	1.4	2.2	2.3	3.3	3.7	4.2	5.3	2.5	3.5	1.4	3.1
229	i	.	0.3	0.3	0.1	0.5	0.2	0.6	1.0	0.9	1.1	0.6	0.6	0.4	1.0
	p	.	0.3	0.3	0.3	0.6	0.4	0.9	1.4	1.8	3.6	1.5	5.4	2.9	6.0
230	i	.	.	.	0.2	.	.	0.1	.	0.1	.	0.2	.	.	.
	p	.	.	.	0.2	.	.	0.1	.	0.2	0.0	0.2	.	.	.
231	i	.	.	0.1	0.1	0.1	0.1	0.0	0.2	0.1	0.0	0.1	0.0	.	.
	p	.	.	0.1	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.1	0.0	.	0.3
232	i	1.9	0.8	1.2	1.1	0.9	2.1	0.7	0.9	0.4	0.7	0.3	1.2	0.7	0.5
	p	2.5	1.1	2.8	2.1	1.6	3.3	1.3	1.2	1.2	1.6	3.7	2.7	3.6	6.0
233	i	.	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.1	0.3	.	0.2	.	0.1
	p	.	0.1	0.3	0.1	0.3	0.2	0.1	0.3	0.3	0.6	0.4	0.7	0.5	0.1
234	i	0.5	.	0.3	0.1	0.1	0.1	0.2	0.1	0.0	0.3	0.3	0.2	0.1	0.2
	p	0.6	0.0	0.3	0.2	0.2	0.2	0.3	0.2	0.2	0.6	0.3	0.5	0.2	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
235	neoplasms musc. skel. syst.	i	1489	0.1	0.0	0.2	.	0.0	0.1	0.1	0.1	0.1	0.1
		p	4280	0.3	0.2	0.4	.	0.1	0.3	0.3	0.3	0.4	0.6
236	fracture: radius/ulna	i	5638	0.4	0.3	0.5	0.5	0.9	0.4	0.1	0.4	0.6	0.3
		p	8225	0.6	0.4	0.7	0.6	1.2	0.4	0.2	0.6	0.9	1.0
237	fracture: tibia/fibula	i	3265	0.2	0.2	0.2	0.2	0.3	0.5	0.1	0.1	0.1	0.1
		p	6160	0.4	0.4	0.4	0.3	0.5	0.8	0.4	0.3	0.3	0.2
238	fracture: hand/foot bones	i	12193	0.8	0.9	0.8	0.4	1.5	0.9	0.7	0.7	1.0	0.5
		p	14694	1.0	1.1	0.9	0.6	1.7	1.1	0.9	0.8	1.3	0.7
239	fracture: collum femoris	i	1883	0.1	0.1	0.2	.	.	0.1	0.0	0.0	0.2	1.9
		p	3701	0.3	0.2	0.3	.	.	0.1	0.0	0.2	0.7	2.9
240	fracture: femur NEC	i	1178	0.1	0.0	0.1	0.1	0.1	0.1	0.0	0.0	0.2	0.3
		p	1864	0.1	0.1	0.2	0.1	0.1	0.1	0.0	0.1	0.3	0.8
241	fracture: skull	i	257	0.0	0.0	0.0	.	0.0	.	0.0	0.0	0.0	0.0
		p	384	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	0.0	0.0
242	fracture: facial bones	i	1292	0.1	0.1	0.0	0.1	0.2	0.2	0.0	0.0	.	0.1
		p	1910	0.1	0.2	0.1	0.1	0.2	0.2	0.1	0.0	.	0.1
243	fracture: spine	i	1013	0.1	0.1	0.1	.	.	0.1	0.0	0.1	.	0.5
		p	2992	0.2	0.2	0.2	.	0.1	0.2	0.1	0.2	0.7	0.9
244	fracture: rib	i	2155	0.1	0.1	0.2	.	0.0	0.0	0.2	0.2	0.4	0.4
		p	2607	0.2	0.2	0.2	.	0.0	0.0	0.2	0.3	0.5	0.6
245	fracture: clavícula	i	1824	0.1	0.1	0.1	0.6	0.2	0.2	0.0	0.1	0.1	0.0
		p	2495	0.2	0.2	0.2	0.6	0.3	0.3	0.1	0.1	0.2	0.1
246	fracture: humerus	i	1019	0.1	0.0	0.1	.	0.2	0.0	0.0	0.1	0.3	0.1
		p	2144	0.1	0.1	0.2	0.1	0.2	0.0	0.0	0.2	0.5	0.4
247	fracture: pelvis	i	382	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	0.2
		p	834	0.1	0.1	0.1	.	.	.	0.0	0.1	0.2	0.2
248	fracture: NEC	i	8557	0.6	0.5	0.6	0.4	0.8	0.7	0.5	0.4	0.7	0.5
		p	11652	0.8	0.8	0.8	0.4	0.9	1.1	0.7	0.6	0.9	0.9
249	sprains/strains ankle(s)	i	46920	3.2	3.6	2.8	1.2	3.5	6.0	3.2	2.2	2.0	0.9
		p	56987	3.9	4.3	3.5	1.4	4.0	7.1	4.1	2.7	2.4	1.5
250	sprains/strains knee(s)	i	33464	2.3	2.7	1.9	0.4	2.5	4.0	2.5	1.6	0.7	1.6
		p	39117	2.7	3.1	2.3	0.4	2.7	4.7	2.8	2.2	1.0	1.9
251	sprains/strains shoulder/arm	i	20990	1.4	1.4	1.4	0.7	1.9	2.4	1.2	1.3	0.8	0.9
		p	24577	1.7	1.7	1.7	0.7	1.9	2.5	1.4	1.8	1.1	1.5
252	sprains/strains wrist/hand	i	40817	2.8	3.1	2.5	1.5	5.3	4.3	2.4	2.0	1.0	1.0
		p	45748	3.1	3.5	2.7	1.9	5.7	4.8	2.7	2.3	1.0	1.2
253	sprains/strains foot/toe	i	32805	2.2	2.5	1.9	1.4	3.4	3.2	2.1	1.8	1.3	0.9
		p	35642	2.4	2.7	2.2	1.4	3.5	3.5	2.3	2.1	1.4	1.0
254	sprains/strains spine, excl. cerv. spine	i	10979	0.7	0.7	0.8	.	0.6	1.1	0.9	0.5	0.5	1.1
		p	13241	0.9	0.9	0.9	.	0.6	1.2	1.2	0.7	0.5	1.6
255	sprains/strains cerv. spine	i	5256	0.4	0.3	0.4	0.1	0.2	0.6	0.5	0.2	0.1	0.0
		p	6926	0.5	0.5	0.5	0.1	0.3	0.8	0.7	0.3	0.1	0.1
256	whiplash calf	i	5628	0.4	0.5	0.3	.	0.0	0.1	0.7	0.6	0.2	0.1
		p	6344	0.4	0.5	0.3	.	0.0	0.2	0.7	0.7	0.2	0.1
257	sprains/strains musc. skel. syst. NEC	i	53142	3.6	4.1	3.2	0.9	3.0	4.1	3.7	3.7	3.8	6.1
		p	61422	4.2	4.6	3.8	1.1	3.1	4.5	4.3	4.3	4.7	7.7
258	(sub)luxation, dislocation	i	7246	0.5	0.5	0.5	0.9	0.3	0.8	0.5	0.3	0.5	0.3
		p	9772	0.7	0.7	0.7	1.1	0.4	0.9	0.6	0.5	0.7	0.6
259	cong. luxation of hip	i	1428	0.1	0.1	0.1	1.1	.	0.0	0.0	0.1	0.0	0.1
		p	3438	0.2	0.1	0.4	2.5	0.0	0.0	0.1	0.1	0.1	0.6

L MUSCULOSKELETAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
235	i	.	.	.	0.1	0.0	0.2	0.0	0.3	0.1	0.1	.	0.2	.	0.2
	p	.	.	0.0	0.1	0.3	0.3	0.2	0.5	0.2	0.5	0.4	0.5	0.4	0.6
236	i	0.3	0.7	0.9	0.9	0.4	0.4	0.2	0.0	0.2	0.6	0.3	0.9	.	0.5
	p	0.3	0.8	1.1	1.2	0.4	0.4	0.3	0.1	0.3	0.9	0.3	1.4	0.0	1.4
237	i	0.3	0.1	0.2	0.4	0.7	0.4	0.1	0.2	0.1	0.2	0.0	0.1	.	0.2
	p	0.3	0.3	0.4	0.6	1.1	0.5	0.4	0.4	0.2	0.4	0.2	0.4	.	0.4
238	i	0.3	0.4	1.6	1.4	1.2	0.6	0.9	0.5	0.6	0.7	0.8	1.1	0.0	0.8
	p	0.6	0.6	1.9	1.5	1.5	0.7	1.1	0.7	0.7	0.9	0.8	1.6	0.0	1.1
239	i	0.1	0.0	0.0	.	0.0	0.0	0.1	0.3	1.5	2.1
	p	0.1	0.0	0.0	.	0.1	0.2	0.5	0.9	1.9	3.4
240	i	0.3	.	0.1	0.1	0.0	0.1	0.0	0.0	.	0.1	0.2	0.3	0.1	0.5
	p	0.3	.	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.2	0.2	0.4	0.2	1.1
241	i	.	.	.	0.0	.	.	0.0	0.0	.	0.0	.	0.0	0.1	.
	p	.	.	0.0	0.0	0.0	.	0.0	0.0	.	0.1	0.0	0.0	0.1	0.0
242	i	.	0.1	0.2	0.2	0.4	.	0.1	0.0	0.0	0.0	.	.	.	0.1
	p	.	0.1	0.3	0.2	0.5	0.0	0.3	0.0	0.0	0.0	.	.	.	0.1
243	i	0.2	.	0.1	0.0	0.0	0.1	.	.	0.3	0.6
	p	.	.	0.1	.	0.2	0.2	0.1	0.1	0.1	0.2	0.9	0.5	0.8	1.0
244	i	.	.	0.0	.	0.0	.	0.2	0.1	0.3	0.2	0.0	0.7	.	0.6
	p	.	.	0.0	.	0.0	0.0	0.2	0.1	0.4	0.2	0.0	0.8	.	0.9
245	i	0.7	0.5	0.1	0.2	0.2	0.2	0.1	.	0.1	0.0	0.0	0.2	.	0.1
	p	0.7	0.5	0.2	0.3	0.3	0.3	0.1	0.1	0.1	0.0	0.1	0.2	0.1	0.1
246	i	.	.	0.1	0.3	.	0.0	0.0	0.0	0.0	0.1	0.4	0.3	.	0.2
	p	0.2	0.0	0.2	0.3	0.0	0.0	0.0	0.0	0.0	0.4	0.4	0.5	0.1	0.6
247	i	0.0	0.0	.	0.0	0.1	.	0.0	0.3
	p	0.1	0.0	0.1	0.0	0.1	0.3	0.0	0.3
248	i	0.2	0.6	0.6	1.1	0.8	0.6	0.5	0.5	0.5	0.3	0.1	1.2	0.2	0.6
	p	0.2	0.6	0.6	1.2	1.3	0.8	0.8	0.6	0.7	0.5	0.4	1.3	0.3	1.3
249	i	1.0	1.3	3.3	3.8	7.0	4.9	4.2	2.2	1.9	2.6	1.5	2.3	0.1	1.3
	p	1.4	1.4	3.6	4.3	8.2	5.9	5.3	2.9	2.3	3.1	1.5	3.1	0.4	2.1
250	i	0.3	0.5	2.6	2.4	5.0	3.0	3.2	1.7	1.6	1.7	0.5	0.9	1.0	1.8
	p	0.3	0.5	2.7	2.6	5.8	3.5	3.6	2.0	2.0	2.4	0.6	1.3	1.3	2.3
251	i	0.4	0.9	1.7	2.1	2.3	2.5	1.2	1.2	1.5	1.1	0.6	1.0	1.2	0.7
	p	0.5	0.9	1.7	2.1	2.4	2.6	1.5	1.4	2.0	1.5	1.0	1.2	1.7	1.4
252	i	1.9	1.1	5.1	5.5	5.1	3.5	2.9	1.9	1.9	2.0	1.2	0.8	0.3	1.4
	p	2.4	1.3	5.7	5.7	5.8	3.9	3.2	2.1	2.2	2.4	1.2	0.9	0.4	1.7
253	i	1.3	1.5	3.3	3.6	4.2	2.2	2.5	1.7	2.0	1.6	0.9	1.6	0.4	1.2
	p	1.3	1.5	3.4	3.7	4.3	2.6	2.7	1.9	2.2	2.0	0.9	1.7	0.4	1.2
254	i	.	.	0.4	0.7	0.8	1.3	1.1	0.7	0.4	0.6	0.5	0.6	0.8	1.2
	p	.	.	0.4	0.8	0.9	1.5	1.5	0.8	0.6	0.7	0.5	0.6	0.8	2.1
255	i	.	0.2	0.0	0.4	0.7	0.6	0.5	0.5	0.2	0.2	0.1	0.1	0.1	.
	p	.	0.2	0.1	0.4	0.8	0.8	0.6	0.8	0.4	0.3	0.1	0.1	0.1	0.2
256	i	.	.	0.0	.	0.3	.	0.7	0.6	0.7	0.5	0.3	0.1	0.2	.
	p	.	.	0.0	.	0.4	0.1	0.8	0.7	0.9	0.6	0.3	0.1	0.2	.
257	i	1.0	0.7	3.4	2.6	5.2	2.9	5.0	2.3	3.1	4.2	3.6	4.0	4.0	7.2
	p	1.0	1.2	3.4	2.7	5.7	3.2	5.8	2.8	3.7	4.8	4.3	5.0	4.8	9.2
258	i	0.9	1.0	0.2	0.5	0.9	0.7	0.6	0.4	0.1	0.4	0.5	0.6	0.4	0.2
	p	1.0	1.2	0.2	0.7	0.9	0.9	0.8	0.5	0.5	0.6	0.8	0.6	0.6	0.6
259	i	0.7	1.5	.	.	0.0	0.0	0.0	0.0	.	0.1	.	0.0	.	0.2
	p	1.3	3.8	.	0.0	0.0	0.0	0.0	0.1	.	0.2	.	0.2	0.4	0.7

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
260	cong. anom. musc. skel. syst. NEC	i	2303	0.2	0.2	0.1	0.7	0.2	0.2	0.1	0.1	.	.
		p	5787	0.4	0.5	0.3	1.8	0.5	0.3	0.4	0.2	0.1	0.0
261	syndromes cervical spine	i	19294	1.3	1.0	1.6	0.2	0.4	0.5	1.7	2.2	1.6	1.4
		p	30193	2.1	1.5	2.6	0.2	0.5	0.6	2.4	3.8	3.2	2.2
262	osteoarthritis of cervical spine	i	11617	0.8	0.5	1.0	.	.	0.0	0.2	1.4	3.6	3.6
		p	23420	1.6	1.1	2.0	.	.	0.0	0.4	3.0	6.3	7.6
263	osteoarthritis of spine NEC	i	8711	0.6	0.4	0.8	.	.	0.0	0.2	1.3	2.0	2.2
		p	22207	1.5	1.0	2.0	.	.	0.1	0.5	3.2	4.4	7.1
264	acquired deformities of spine	i	10004	0.7	0.6	0.7	0.1	1.1	1.2	0.8	0.2	0.2	0.4
		p	22683	1.5	1.4	1.7	0.1	2.3	2.2	1.8	0.8	0.8	1.8
265	back pain with radiating sympt.	i	6753	0.5	0.5	0.4	.	.	0.2	0.8	0.7	0.3	0.4
		p	12434	0.8	0.9	0.8	.	.	0.4	1.4	1.2	0.9	0.6
266	lumbago	i	96167	6.6	7.7	5.4	.	0.3	5.0	9.5	9.7	6.2	4.6
		p	129036	8.8	10.0	7.6	.	0.4	6.0	12.4	14.1	8.6	6.6
267	sciatica	i	15490	1.1	1.0	1.1	.	0.0	0.2	1.1	2.3	1.9	1.3
		p	24969	1.7	1.4	2.0	.	0.0	0.3	1.6	3.7	3.6	2.2
268	slipped lumbar disc	i	18703	1.3	1.5	1.1	.	0.0	0.6	1.9	2.4	0.9	0.5
		p	45564	3.1	3.6	2.7	.	0.0	1.1	4.1	6.1	3.6	2.7
269	ganglion joint/tendon	i	9883	0.7	0.5	0.8	0.1	0.3	1.0	0.9	0.6	0.2	0.3
		p	11897	0.8	0.7	1.0	0.2	0.4	1.1	1.1	0.8	0.4	0.4
270	rheumatoid arthritis	i	7975	0.5	0.3	0.8	0.0	0.1	0.1	0.3	1.2	1.2	1.8
		p	44673	3.0	1.6	4.4	0.1	0.2	0.3	1.3	5.1	11.0	13.7
271	spondylitis (ankylosing)	i	1430	0.1	0.1	0.1	.	0.1	0.1	0.1	0.1	0.1	.
		p	5350	0.4	0.4	0.3	.	0.1	0.1	0.5	0.5	1.1	0.2
272	osteoarthritis hip	i	9987	0.7	0.5	0.9	.	.	0.0	0.1	1.2	2.9	3.9
		p	27153	1.9	1.0	2.7	.	.	0.0	0.2	2.6	7.6	13.6
273	osteoarthritis knee	i	16099	1.1	0.6	1.6	.	.	0.0	0.3	1.9	4.4	5.4
		p	36919	2.5	1.2	3.8	.	.	0.0	0.6	3.2	9.1	19.0
274	osteoarthritis NEC	i	15605	1.1	0.6	1.5	.	.	0.0	0.4	1.8	2.9	6.8
		p	47020	3.2	1.5	4.9	.	.	0.2	0.6	4.9	11.0	22.7
275	arthritis in allied conditions NEC	i	15818	1.1	0.9	1.3	0.2	0.5	0.9	1.2	1.1	2.0	1.9
		p	31789	2.2	1.5	2.8	0.3	0.6	1.1	1.9	2.6	4.6	8.0
276	shoulder syndrome	i	43679	3.0	2.8	3.1	.	0.2	1.2	2.9	7.0	3.6	3.2
		p	63275	4.3	4.0	4.6	0.0	0.2	1.6	3.7	10.6	5.9	5.6
277	dis. bursa, excl. shoulder	i	20376	1.4	1.8	1.0	.	0.3	1.0	1.6	2.1	1.8	2.2
		p	24914	1.7	2.1	1.3	.	0.4	1.2	1.9	2.7	2.3	2.7
278	epicondylitis lateralis	i	25355	1.7	1.9	1.5	0.0	0.0	0.4	2.6	3.7	1.3	0.2
		p	34434	2.3	2.6	2.1	0.0	0.0	0.5	3.3	5.4	1.8	0.3
279	tendinitis/synovitis	i	63577	4.3	4.5	4.2	0.0	2.0	4.1	5.8	5.4	3.9	3.2
		p	78363	5.3	5.3	5.4	0.0	2.2	5.1	7.1	6.9	5.0	3.7
280	osteocondrosis	i	3843	0.3	0.3	0.2	0.0	1.0	0.4	0.2	0.0	0.1	.
		p	7283	0.5	0.6	0.4	0.0	1.5	0.8	0.4	0.1	0.1	.
281	osteoporosis	i	2834	0.2	0.1	0.3	.	.	.	0.0	0.2	0.9	1.5
		p	21165	1.4	0.4	2.4	.	.	.	0.2	2.1	5.5	10.2
282	ac. traum. meniscus/lig. knee	i	17885	1.2	1.5	0.9	.	0.2	1.9	1.6	1.4	1.0	0.2
		p	20196	1.4	1.7	1.0	.	0.2	2.2	1.8	1.7	1.1	0.2
283	retropatellar chondropathy	i	14111	1.0	0.9	1.0	.	1.0	2.2	1.1	0.5	0.2	0.1
		p	19080	1.3	1.2	1.4	.	1.4	3.0	1.4	0.7	0.2	0.2
284	chron. internal knee derangement	i	4148	0.3	0.3	0.2	.	0.2	0.3	0.4	0.4	0.1	0.2
		p	6344	0.4	0.5	0.4	.	0.2	0.4	0.5	0.5	0.7	0.5

L MUSCULOSKELETAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
260	i	0.7	0.7	0.3	0.1	0.1	0.2	0.2	0.1	0.1	0.1
	p	2.2	1.5	0.6	0.4	0.4	0.3	0.4	0.3	0.2	0.2	.	0.2	0.1	0.0
261	i	0.1	0.4	0.3	0.5	0.4	0.6	1.1	2.3	1.8	2.6	1.5	1.8	1.7	1.2
	p	0.1	0.4	0.4	0.7	0.4	0.8	1.5	3.4	2.9	4.6	2.9	3.4	2.0	2.3
262	i	0.0	.	0.2	0.2	1.1	1.7	2.6	4.3	2.5	4.1
	p	0.0	.	0.4	0.5	2.8	3.3	4.4	7.8	4.8	9.0
263	i	0.0	0.1	0.2	0.3	1.2	1.3	0.5	3.1	1.8	2.4
	p	0.1	0.1	0.6	0.5	3.0	3.4	1.7	6.4	3.1	9.2
264	i	.	0.2	0.7	1.5	1.5	0.8	0.8	0.8	0.1	0.3	0.1	0.3	0.1	0.6
	p	.	0.3	1.5	3.0	2.3	2.0	1.8	1.8	0.6	1.0	0.5	1.1	0.7	2.4
265	i	0.1	0.3	1.1	0.5	0.6	0.7	0.3	0.3	1.1	0.0
	p	0.3	0.5	1.8	1.0	1.0	1.5	1.0	0.8	1.4	0.2
266	i	.	.	0.4	0.3	5.7	4.3	11.6	7.4	11.2	8.3	6.8	5.8	4.7	4.5
	p	.	.	0.5	0.3	7.0	5.0	14.6	10.0	15.5	12.8	8.5	8.6	6.0	6.9
267	i	.	.	0.1	0.0	0.1	0.4	1.2	0.9	1.7	3.0	2.1	1.7	2.3	0.7
	p	.	.	0.1	0.0	0.1	0.5	1.6	1.7	2.9	4.6	3.3	3.9	2.8	1.9
268	i	.	.	0.0	.	0.7	0.5	2.3	1.5	2.8	2.0	0.5	1.2	0.3	0.7
	p	.	.	0.1	0.0	1.3	0.8	5.1	3.1	7.2	5.0	3.0	4.1	1.1	3.6
269	i	0.0	0.2	0.1	0.6	0.6	1.4	1.0	0.9	0.5	0.8	0.2	0.2	0.1	0.4
	p	0.0	0.5	0.1	0.7	0.7	1.5	1.1	1.1	0.6	1.0	0.5	0.4	0.1	0.6
270	i	0.0	.	0.0	0.1	.	0.1	0.2	0.4	0.7	1.8	0.4	1.8	0.7	2.3
	p	0.1	0.1	0.3	0.2	0.0	0.6	0.8	1.8	3.3	6.9	7.2	13.9	5.8	18.0
271	i	.	.	0.2	0.0	0.1	.	0.1	0.1	0.1	0.1	0.2	0.0	.	.
	p	.	.	0.2	0.0	0.1	0.1	0.5	0.5	0.6	0.4	2.0	0.4	0.0	0.2
272	i	0.0	0.0	0.1	0.1	0.8	1.6	1.9	3.7	4.2	3.7
	p	0.0	0.0	0.2	0.2	1.8	3.3	4.3	10.2	8.7	16.2
273	i	0.0	0.1	0.4	0.3	1.2	2.6	2.5	6.0	1.4	7.5
	p	0.0	0.1	0.8	0.5	2.2	4.2	4.8	12.4	6.1	25.9
274	i	0.1	.	0.4	0.4	1.2	2.3	1.6	3.9	2.9	8.8
	p	0.2	0.2	0.6	0.6	3.1	6.6	4.5	16.0	9.3	29.9
275	i	0.3	0.1	0.6	0.3	0.7	1.2	1.4	0.9	0.7	1.6	0.9	2.9	1.0	2.4
	p	0.4	0.2	0.7	0.5	0.9	1.4	1.9	2.0	1.7	3.4	2.6	6.1	2.7	10.7
276	i	.	.	0.1	0.2	0.9	1.4	2.7	3.1	7.1	6.9	3.4	3.7	2.3	3.7
	p	0.0	.	0.1	0.4	1.4	1.7	3.5	4.0	10.3	10.8	5.6	6.2	3.5	6.6
277	i	.	.	0.4	0.3	1.4	0.7	2.3	0.9	2.3	1.8	2.2	1.4	3.6	1.5
	p	.	.	0.5	0.3	1.6	0.8	2.5	1.2	3.0	2.4	2.6	2.0	3.7	2.2
278	i	0.0	.	0.0	0.0	0.2	0.5	2.8	2.3	4.0	3.4	1.9	0.8	.	0.2
	p	0.0	.	0.0	0.0	0.3	0.7	3.9	2.6	5.6	5.2	2.8	1.0	.	0.4
279	i	0.0	0.1	1.5	2.4	3.8	4.5	6.6	5.0	5.3	5.6	3.9	3.9	2.7	3.5
	p	0.0	0.1	1.6	2.8	4.6	5.6	7.8	6.5	6.4	7.3	4.4	5.4	3.0	4.1
280	i	0.1	.	1.1	0.8	0.6	0.3	0.2	0.1	0.0	0.0	.	0.1	.	.
	p	0.1	.	1.5	1.5	1.1	0.6	0.4	0.4	0.2	0.0	.	0.1	.	.
281	i	0.0	0.0	0.1	0.3	0.3	1.3	0.8	1.9
	p	0.2	0.3	0.7	3.5	1.2	8.8	3.3	13.9
282	i	.	.	0.1	0.3	2.4	1.3	2.1	1.0	1.6	1.3	1.0	1.1	.	0.3
	p	.	.	0.1	0.3	2.9	1.4	2.3	1.2	1.9	1.4	1.0	1.2	0.1	0.3
283	i	.	.	0.9	1.1	2.1	2.4	1.0	1.1	0.4	0.6	0.1	0.2	0.1	0.1
	p	.	.	1.1	1.7	2.9	3.1	1.3	1.5	0.7	0.8	0.1	0.2	0.3	0.1
284	i	.	.	0.2	0.2	0.3	0.2	0.5	0.3	0.4	0.3	0.0	0.2	.	0.3
	p	.	.	0.2	0.3	0.5	0.3	0.7	0.4	0.5	0.4	0.8	0.7	.	0.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
285	acquired deformities of limbs	i	25365	1.7	1.4	2.1	1.9	2.5	1.3	1.4	1.9	1.9	2.3
		p	37875	2.6	2.1	3.1	2.7	3.3	2.1	2.0	2.7	3.2	4.3
286	dis. musc. skel. system NEC	i	20549	1.4	1.2	1.6	0.2	1.1	1.5	1.7	1.7	1.0	0.8
		p	32407	2.2	2.0	2.4	0.4	1.4	1.9	2.4	2.9	2.3	2.8

L MUSCULOSKELETAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
285	i	2.2	1.6	2.4	2.5	1.2	1.5	1.1	1.7	1.3	2.5	0.9	2.7	1.6	2.8
	p	2.9	2.5	3.1	3.6	1.8	2.3	1.8	2.3	2.0	3.4	1.6	4.4	1.9	5.6
286	i	.	0.4	0.9	1.3	1.2	1.7	1.6	1.8	1.3	2.2	1.0	1.0	0.7	0.8
	p	0.4	0.4	1.0	1.8	1.7	2.2	2.4	2.5	2.6	3.3	2.3	2.4	2.4	3.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals abs.	/1000 pers.	sex		age						
					male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
287	headache/facial pain	i	23008	1.6	1.2	1.9	0.2	1.2	2.0	1.8	1.7	1.3	1.3
		p	77413	5.3	3.5	7.0	0.2	1.8	3.7	5.5	8.3	8.2	7.8
288	sensibility disturbance	i	5279	0.4	0.4	0.3	.	0.1	0.3	0.3	0.6	0.6	0.6
		p	11221	0.8	0.7	0.8	0.0	0.1	0.3	0.5	1.2	2.0	2.6
289	restless legs syndrome	i	5413	0.4	0.3	0.5	.	.	0.0	0.2	0.9	1.1	0.7
		p	14671	1.0	0.6	1.4	.	.	0.0	0.3	2.1	3.4	4.3
290	convulsions	i	1859	0.1	0.2	0.1	1.8	0.1
		p	2798	0.2	0.2	0.2	2.7	0.1
291	symptoms sense/smell/taste	i	275	0.0	0.0	0.0	.	.	.	0.0	0.1	0.0	.
		p	283	0.0	0.0	0.0	.	.	.	0.0	0.1	0.0	.
292	vertigo	i	23140	1.6	1.1	2.1	0.0	0.3	1.0	1.2	2.5	3.3	4.7
		p	61455	4.2	2.4	5.9	0.1	0.5	1.2	1.9	5.8	10.9	26.2
293	paralysis	i	1317	0.1	0.1	0.0	.	.	0.0	0.1	0.2	0.3	0.0
		p	7086	0.5	0.6	0.4	0.3	0.1	0.1	0.3	0.8	1.2	1.5
294	sympt./complt. speech	i	2947	0.2	0.2	0.2	0.4	0.6	0.2	0.1	0.0	0.1	0.3
		p	5811	0.4	0.4	0.3	0.7	1.2	0.3	0.2	0.1	0.3	0.5
295	fear of neurological disease	i	616	0.0	0.0	0.1	.	0.1	.	0.1	0.0	0.0	.
		p	1446	0.1	0.1	0.1	.	0.1	0.0	0.2	0.0	0.2	0.1
296	disability/impairment neurol. syst.	i	30	0.0	.	0.0	0.0	.	.
		p	274	0.0	0.0	0.0	.	0.0	.	0.0	0.0	0.0	0.2
297	sympt./complt. neurol. syst. NEC	i	903	0.1	0.1	0.1	0.0	.	.	0.0	0.2	0.3	.
		p	2612	0.2	0.2	0.2	0.0	0.0	0.0	0.1	0.3	0.8	0.3
298	poliomyelitis	i	0
		p	1094	0.1	0.0	0.1	.	.	.	0.1	0.1	0.3	0.3
299	meningitis/encefalitis	i	542	0.0	0.1	0.0	0.4	0.0	0.0	0.0	.	.	.
		p	1369	0.1	0.1	0.1	0.5	0.1	0.0	0.0	0.1	0.0	0.2
300	tetanus	i	0
		p	13	0.0	0.0	.	.	0.0
301	infections neur. syst. NEC	i	84	0.0	.	0.0	.	.	0.0	0.0	.	.	.
		p	297	0.0	0.0	0.0	.	.	0.0	0.0	0.0	.	0.1
302	malignant neoplasm neur. syst.	i	471	0.0	0.1	0.0	0.1	.	.	0.0	0.0	0.3	.
		p	2492	0.2	0.2	0.1	0.1	.	0.1	0.1	0.3	0.4	0.6
303	benigne neopl. neur. syst.	i	555	0.0	0.0	0.0	.	.	0.0	0.0	0.1	0.2	.
		p	1595	0.1	0.1	0.1	0.1	.	0.0	0.0	0.2	0.2	0.4
304	unspecified neopl. neur. syst.	i	611	0.0	0.0	0.0	.	0.0	0.1	0.0	0.0	0.1	0.1
		p	1641	0.1	0.1	0.1	.	0.0	0.1	0.1	0.2	0.1	0.1
305	concussion: commotio cer.	i	8056	0.5	0.6	0.5	1.2	1.1	0.7	0.5	0.1	0.2	0.4
		p	9982	0.7	0.7	0.7	1.3	1.3	0.8	0.6	0.2	0.5	0.7
306	concussion: contusio cer.	i	1386	0.1	0.1	0.1	0.2	0.1	0.2	0.0	0.0	0.0	0.1
		p	2175	0.1	0.2	0.1	0.2	0.2	0.3	0.1	0.1	0.2	0.1
307	head injury NOS/NEC, w/o skull fract.	i	1501	0.1	0.1	0.1	0.5	0.3	0.0	0.1	0.0	.	0.1
		p	1746	0.1	0.1	0.2	0.5	0.3	0.0	0.1	0.0	.	0.1
308	injuries neur. syst. NEC	i	774	0.1	0.0	0.1	.	0.0	0.0	0.0	0.1	0.2	0.1
		p	1323	0.1	0.1	0.1	.	0.0	0.0	0.1	0.2	0.2	0.2
309	cong. anom. neur. syst.	i	296	0.0	0.0	0.0	0.2	.	0.0	0.0	0.0	.	.
		p	3542	0.2	0.2	0.2	0.7	0.4	0.4	0.1	0.1	0.1	0.1
310	multiple sclerosis	i	844	0.1	0.1	0.1	.	.	.	0.1	0.1	0.0	.
		p	6856	0.5	0.4	0.5	.	.	0.0	0.6	0.9	1.0	0.4
311	Parkinsonism	i	1728	0.1	0.2	0.1	.	.	.	0.0	0.1	0.7	0.9
		p	19870	1.4	1.1	1.6	.	.	.	0.2	1.0	5.1	13.3

N NEUROLOGICAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
287	i	0.2	0.2	1.0	1.4	1.2	2.8	1.4	2.1	1.4	1.9	1.5	1.2	0.7	1.6
	p	0.2	0.2	1.4	2.3	2.5	5.0	3.7	7.4	6.4	10.1	4.6	11.0	3.3	10.1
288	i	.	.	0.1	0.1	0.3	0.2	0.4	0.2	0.7	0.6	0.1	1.0	0.8	0.5
	p	0.0	0.0	0.1	0.1	0.4	0.3	0.6	0.4	1.1	1.3	1.0	2.7	3.3	2.2
289	i	0.1	0.1	0.3	0.7	1.2	0.9	1.1	0.3	0.8
	p	0.0	0.1	0.2	0.4	1.2	2.9	2.7	3.9	2.8	5.2
290	i	2.2	1.4	0.1	0.1
	p	3.1	2.2	0.1	0.1
291	i	0.0	0.0	0.1	0.0	0.1	0.0	.	.
	p	0.0	0.0	0.1	0.0	0.1	0.0	.	.
292	i	0.0	0.0	0.1	0.5	0.7	1.3	0.8	1.6	1.7	3.3	1.8	4.4	5.4	4.3
	p	0.0	0.1	0.1	0.9	0.8	1.7	1.3	2.4	3.4	8.1	5.2	15.3	21.9	28.6
293	i	0.0	0.0	0.1	0.0	0.3	0.1	0.5	0.2	0.1	0.0
	p	0.4	0.2	0.1	0.1	0.2	0.1	0.3	0.2	0.9	0.8	1.9	0.7	2.7	0.9
294	i	0.2	0.6	0.9	0.3	0.1	0.3	0.1	0.1	.	0.1	0.2	0.0	0.4	0.3
	p	0.6	0.8	1.7	0.7	0.3	0.4	0.2	0.2	0.0	0.2	0.3	0.3	0.9	0.3
295	i	.	.	.	0.1	.	.	0.1	0.1	0.0	0.0	.	0.0	.	.
	p	.	.	.	0.1	0.0	0.0	0.1	0.2	0.0	0.1	.	0.3	0.3	.
296	i	0.0
	p	.	.	.	0.0	.	.	0.0	.	0.0	0.0	.	0.1	0.2	0.1
297	i	.	0.0	0.0	0.0	0.1	0.3	0.7	.	.	.
	p	0.0	0.1	0.1	.	0.0	0.0	0.1	0.1	0.4	0.3	0.9	0.7	0.3	0.4
298	i
	p	0.1	0.0	0.1	0.1	.	0.6	0.1	0.5
299	i	0.6	0.1	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.9	0.1	0.1	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.1	.	0.5	.
300	i
	p	0.0
301	i	0.0	.	0.0
	p	0.0	0.0	0.1	0.0	0.2
302	i	.	0.2	0.0	.	0.1	.	0.6	.	.	.
	p	0.0	0.2	.	.	0.1	.	0.1	0.1	0.4	0.3	0.8	0.1	1.3	0.2
303	i	0.0	.	0.0	0.0	.	0.1	0.4	.	.	.
	p	0.2	.	.	.	0.0	.	0.1	0.0	0.0	0.4	0.5	0.0	0.5	0.4
304	i	.	.	0.0	.	0.1	0.0	.	0.1	0.1	0.0	0.0	0.2	0.2	.
	p	.	.	0.0	.	0.1	0.0	0.1	0.2	0.3	0.1	0.0	0.2	0.2	.
305	i	1.7	0.6	1.1	1.2	0.9	0.6	0.5	0.5	0.0	0.2	0.2	0.1	0.5	0.3
	p	1.9	0.6	1.2	1.4	0.9	0.8	0.6	0.6	0.2	0.2	0.2	0.7	0.5	0.7
306	i	0.1	0.3	0.2	0.0	0.2	0.3	0.1	0.0	0.0	0.1	0.1	.	.	0.1
	p	0.1	0.3	0.2	0.2	0.2	0.3	0.2	0.1	0.0	0.1	0.4	.	.	0.1
307	i	0.4	0.6	0.1	0.4	0.0	0.0	0.1	0.1	.	0.1	.	.	.	0.2
	p	0.4	0.6	0.1	0.4	0.0	0.0	0.1	0.2	.	0.1	.	.	.	0.2
308	i	.	.	0.0	.	0.1	.	0.0	0.1	0.1	0.1	0.1	0.3	.	0.2
	p	.	.	0.1	.	0.1	0.0	0.1	0.1	0.1	0.2	0.1	0.3	0.3	0.2
309	i	.	0.3	.	.	0.1	.	.	0.0	.	0.0
	p	0.5	0.9	0.3	0.5	0.3	0.4	0.2	0.1	0.2	0.0	.	0.2	0.2	0.1
310	i	0.1	0.1	0.0	0.1	0.1	.	.	.
	p	0.1	0.5	0.7	0.8	1.0	0.9	1.0	.	0.6
311	i	0.0	.	0.1	0.0	1.4	0.2	1.1	0.9
	p	0.1	0.3	1.2	0.9	6.2	4.3	11.5	14.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
312	epilepsy, all types	i	4222	0.3	0.3	0.3	0.2	0.2	0.4	0.3	0.2	0.5	0.2
		p	50891	3.5	3.8	3.2	1.3	1.7	2.8	4.1	3.9	5.4	4.2
313	migraine	i	18002	1.2	0.6	1.9	.	0.5	1.4	1.8	1.6	0.5	0.4
		p	62359	4.2	1.9	6.5	.	0.9	2.8	5.3	7.8	4.0	2.7
314	facial paralysis, Bell's palsy	i	1383	0.1	0.1	0.1	.	0.0	0.1	0.1	0.2	0.0	0.4
		p	2080	0.1	0.1	0.1	.	0.0	0.2	0.1	0.3	0.1	0.5
315	trigeminal neuralgia	i	1582	0.1	0.1	0.2	.	.	0.0	0.1	0.2	0.3	0.1
		p	4033	0.3	0.1	0.4	.	.	0.0	0.2	0.5	0.7	1.1
316	carpal tunnel syndrome	i	4050	0.3	0.1	0.4	.	.	0.1	0.4	0.5	0.5	0.5
		p	6737	0.5	0.2	0.8	.	.	0.1	0.5	0.8	1.0	0.9
317	cluster headache	i	502	0.0	0.0	0.0	.	.	0.1	0.1	0.0	.	.
		p	965	0.1	0.1	0.1	.	.	0.1	0.1	0.1	0.1	.
318	dis. of periph. nerv. syst. NEC	i	32173	2.2	2.1	2.3	0.2	0.1	1.2	2.9	3.1	3.1	3.7
		p	56890	3.9	3.7	4.0	0.4	0.3	1.7	4.2	6.5	6.6	7.7
319	dis. of central nerv. syst. NEC	i	3808	0.3	0.3	0.2	0.1	0.0	0.1	0.2	0.2	1.0	0.9
		p	14310	1.0	1.0	1.0	0.5	0.3	0.4	0.7	1.3	2.6	3.0

N NEUROLOGICAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
312	i	0.0	0.4	0.3	0.1	0.3	0.5	0.3	0.3	0.2	0.1	0.9	0.2	0.3	0.2
	p	1.2	1.5	1.9	1.6	2.4	3.2	4.5	3.6	4.2	3.6	7.7	3.7	5.2	3.7
313	i	.	.	0.4	0.5	0.5	2.3	0.9	2.7	0.6	2.5	0.2	0.7	.	0.7
	p	.	.	0.9	1.0	1.4	4.3	2.4	8.2	2.8	12.6	1.5	5.8	1.8	3.2
314	i	.	.	0.0	.	0.1	.	0.1	0.1	0.2	0.2	0.1	.	0.1	0.6
	p	.	.	0.0	.	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.0	0.1	0.7
315	i	0.0	0.1	0.1	0.1	0.4	0.4	0.3	0.0	0.2
	p	0.1	0.1	0.3	0.3	0.7	0.4	1.0	0.2	1.5
316	i	0.0	0.1	0.2	0.5	0.1	0.8	0.0	0.8	0.3	0.6
	p	0.1	0.2	0.2	0.8	0.2	1.4	0.3	1.5	0.5	1.1
317	i	0.1	0.1	0.0	0.0	0.0
	p	0.1	0.1	0.0	0.0	0.2	0.2	.	.	.
318	i	0.4	.	0.1	0.1	0.9	1.5	3.0	2.9	2.9	3.3	3.1	3.1	3.0	4.0
	p	0.6	0.1	0.4	0.2	1.4	2.1	4.2	4.2	6.5	6.5	6.3	6.8	7.7	7.7
319	i	0.1	0.2	0.1	0.0	0.1	0.1	0.2	0.2	0.3	0.2	1.5	0.7	1.3	0.7
	p	0.4	0.5	0.1	0.5	0.5	0.3	0.7	0.7	1.4	1.3	2.7	2.5	4.2	2.4

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals abs.	/1000 pers.	sex		age						
					male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
320	feeling anxious/nervous/tense	i	57024	3.9	2.4	5.3	0.5	1.3	2.8	4.4	5.4	5.5	6.4
		p	307605	21.0	13.3	28.4	1.0	1.7	6.1	17.0	39.1	49.3	54.1
321	acute stress (transient) reaction	i	26541	1.8	1.3	2.3	0.0	0.3	1.7	2.6	2.3	1.4	1.5
		p	66992	4.6	3.2	5.9	0.0	0.3	2.7	5.7	7.1	6.7	6.0
322	feeling depressed	i	5612	0.4	0.3	0.5	.	.	0.2	0.4	0.6	0.7	0.7
		p	24390	1.7	1.2	2.1	.	.	0.6	1.5	3.0	3.6	4.0
323	feeling/behaving irritable	i	2938	0.2	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.5	0.8
		p	14879	1.0	0.8	1.2	0.3	0.2	0.3	0.5	1.3	2.8	5.5
324	suicide attempt	i	1663	0.1	0.1	0.1	.	0.1	0.0	0.1	0.2	0.2	0.2
		p	1889	0.1	0.2	0.1	.	0.1	0.0	0.2	0.2	0.2	0.2
325	disturbances of sleep	i	15100	1.0	0.8	1.3	1.2	0.1	0.5	0.8	1.4	2.6	2.7
		p	289739	19.7	11.4	27.8	2.6	0.5	1.5	6.8	32.2	71.7	104.5
326	sexual problems	i	4368	0.3	0.3	0.3	.	.	0.4	0.4	0.5	0.3	.
		p	7192	0.5	0.5	0.5	.	0.1	0.5	0.6	0.8	0.4	0.3
327	hypochondr. dis./probl. with being ill	i	2505	0.2	0.2	0.1	.	0.0	0.3	0.2	0.2	0.2	0.0
		p	12408	0.8	0.9	0.8	.	0.1	0.7	1.1	1.1	0.9	1.1
328	tension headache	i	31359	2.1	1.3	2.9	.	0.9	3.1	2.8	2.2	1.4	1.2
		p	47028	3.2	2.1	4.3	.	1.2	4.1	4.3	3.8	2.2	1.7
329	hyperventilation	i	30074	2.0	1.6	2.5	0.0	0.5	2.6	3.2	2.0	1.2	1.2
		p	46680	3.2	2.2	4.1	0.0	0.7	3.5	4.9	3.4	2.6	1.8
330	enuresis	i	3829	0.3	0.3	0.2	0.6	1.5	0.1	0.0	.	.	0.0
		p	13655	0.9	1.0	0.9	1.5	5.2	0.6	0.2	0.0	0.1	0.2
331	anorexia nervosa/boulimia	i	1837	0.1	0.0	0.2	.	0.1	0.4	0.1	0.0	.	.
		p	4199	0.3	0.0	0.5	.	0.1	0.9	0.3	0.1	0.0	.
332	eating/feeding probl. NEC	i	678	0.0	0.0	0.1	0.0	.	0.1	0.0	.	0.1	0.1
		p	1717	0.1	0.1	0.2	0.1	.	0.2	0.1	0.1	0.1	0.3
333	chronic alcohol abuse	i	5468	0.4	0.6	0.2	.	.	0.2	0.6	0.6	0.3	0.2
		p	26142	1.8	2.6	0.9	.	.	0.4	2.4	3.7	2.0	1.1
334	acute alcohol abuse	i	759	0.1	0.1	0.0	.	.	0.0	0.0	0.2	.	.
		p	1376	0.1	0.1	0.0	.	.	0.1	0.1	0.2	0.0	.
335	tobacco abuse	i	7211	0.5	0.5	0.4	.	0.0	0.3	0.9	0.6	0.2	0.0
		p	31086	2.1	2.2	2.0	.	0.1	1.4	3.1	3.6	1.4	1.0
336	drug abuse	i	676	0.0	0.1	0.0	.	.	0.2	0.0	.	.	.
		p	5311	0.4	0.5	0.2	.	.	0.6	0.7	0.2	0.1	.
337	medicinal abuse	i	484	0.0	0.0	0.1	.	.	0.0	0.0	0.0	0.1	0.0
		p	6393	0.4	0.3	0.6	.	.	0.1	0.5	0.8	0.8	1.1
338	abuse substances NEC	i	20	0.0	.	0.0	0.0
		p	539	0.0	0.0	0.1	0.0	.	0.0	0.0	0.0	0.0	0.1
339	dist. memory/concentration	i	1648	0.1	0.1	0.1	.	0.0	0.0	0.1	0.0	0.1	0.9
		p	7227	0.5	0.4	0.6	.	0.0	0.1	0.2	0.3	1.2	4.9
340	disturbance behaviour child	i	2424	0.2	0.2	0.1	1.3	0.6	0.0
		p	3770	0.3	0.3	0.2	1.8	0.9	0.1
341	sympt./complt. behaviour adolescent	i	1157	0.1	0.1	0.1	.	0.1	0.4
		p	2812	0.2	0.1	0.2	.	0.3	0.7
342	sympt./complt. adult	i	2161	0.1	0.1	0.2	.	.	0.2	0.1	0.2	0.2	0.4
		p	9038	0.6	0.6	0.6	.	.	0.6	0.6	0.7	1.0	1.7
343	phase of live problem adult	i	752	0.1	0.0	0.1	.	.	0.1	0.0	0.1	0.1	0.3
		p	4008	0.3	0.2	0.4	.	.	0.2	0.2	0.3	0.4	1.5
344	dist. learning/mental development	i	582	0.0	0.1	0.0	0.3	0.1	0.0
		p	3470	0.2	0.2	0.2	1.1	0.6	0.2	0.1	0.2	0.1	0.1

P PSYCHOLOGICAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
320	i	0.3	0.6	1.2	1.4	1.3	4.3	3.2	5.6	3.3	7.5	2.8	7.6	3.2	8.1
	p	0.6	1.3	1.5	2.0	4.1	8.2	11.1	23.2	25.8	52.2	34.2	61.0	34.9	64.3
321	i	0.0	.	0.1	0.5	0.9	2.5	2.1	3.2	1.9	2.7	0.8	1.9	0.8	1.8
	p	0.0	.	0.2	0.5	1.7	3.9	4.5	7.0	5.1	9.2	3.3	9.4	2.9	7.7
322	i	0.0	0.3	0.4	0.4	0.3	0.9	0.8	0.6	0.7	0.7
	p	0.5	0.7	1.1	1.9	2.2	3.8	2.4	4.6	2.5	4.8
323	i	.	0.3	0.2	0.0	0.1	0.3	0.1	0.1	0.1	0.3	0.7	0.3	1.1	0.7
	p	0.4	0.3	0.3	0.0	0.3	0.3	0.7	0.4	0.9	1.7	3.0	2.6	3.1	6.7
324	i	.	.	0.1	.	0.0	0.0	0.2	0.1	0.2	0.1	0.1	0.2	0.5	.
	p	.	.	0.1	.	0.0	0.0	0.2	0.1	0.2	0.2	0.1	0.2	0.5	0.0
325	i	0.9	1.4	0.3	0.0	0.2	0.7	0.8	0.8	1.1	1.6	1.2	3.7	1.4	3.4
	p	2.3	2.9	0.7	0.3	0.6	2.3	5.2	8.5	19.9	44.2	44.1	92.9	68.6	123.5
326	i	0.1	0.7	0.3	0.5	0.7	0.2	0.6	.	.	.
	p	.	.	.	0.1	0.1	0.9	0.5	0.8	1.2	0.3	0.8	.	0.5	0.2
327	i	.	.	.	0.0	0.5	0.2	0.1	0.2	0.3	0.1	.	0.3	0.1	.
	p	.	.	0.1	0.1	0.8	0.5	1.0	1.2	1.5	0.8	0.7	1.0	0.8	1.3
328	i	.	.	0.5	1.4	1.6	4.5	1.7	4.0	1.7	2.7	0.3	2.3	1.4	1.0
	p	.	.	0.7	1.8	2.1	6.1	2.7	5.9	2.9	4.6	0.8	3.2	1.6	1.7
329	i	0.0	0.0	0.4	0.5	1.5	3.7	2.7	3.6	1.4	2.5	0.5	1.8	0.9	1.3
	p	0.0	0.0	0.4	0.9	2.0	5.0	3.8	6.0	2.3	4.5	1.2	3.6	1.3	2.1
330	i	0.7	0.6	1.9	1.1	0.1	0.1	0.0	0.0	0.0
	p	1.6	1.5	5.9	4.5	0.5	0.7	0.1	0.3	0.0	0.0	0.0	0.1	.	0.4
331	i	.	.	.	0.2	.	0.8	0.1	0.2	.	0.1
	p	.	.	.	0.2	0.0	1.7	0.1	0.5	.	0.3	.	0.0	.	.
332	i	0.1	.	.	.	0.1	0.2	.	0.1	.	.	0.2	0.0	0.1	0.1
	p	0.1	0.2	.	.	0.1	0.3	.	0.3	0.1	0.1	0.2	0.0	0.1	0.4
333	i	0.3	0.2	0.9	0.3	0.8	0.4	0.8	.	0.3	0.2
	p	0.5	0.4	3.4	1.4	5.9	1.5	3.0	1.2	2.1	0.6
334	i	0.1	.	0.1	0.0	0.2	0.1
	p	0.1	.	0.2	0.1	0.2	0.1	0.1	.	.	.
335	i	.	.	.	0.1	0.3	0.3	0.9	0.9	0.8	0.5	0.4	.	0.1	.
	p	.	.	0.0	0.2	1.4	1.3	2.8	3.5	4.3	3.0	1.6	1.3	2.4	0.3
336	i	0.2	0.1	0.0	0.0
	p	0.5	0.6	1.1	0.3	0.2	0.2	0.2	.	.	.
337	i	0.0	.	0.1	0.1	0.0	.	0.1	.	0.0
	p	0.2	0.4	0.6	0.5	1.1	0.8	0.8	0.1	1.7
338	i	.	0.0
	p	.	0.0	.	.	0.0	0.1	0.0	0.1	.	0.1	0.0	0.1	.	0.2
339	i	.	.	0.1	.	0.1	.	0.1	0.1	0.0	0.0	0.1	0.2	1.1	0.9
	p	.	.	0.1	0.0	0.2	.	0.3	0.2	0.1	0.4	0.9	1.5	4.3	5.2
340	i	1.4	1.1	1.0	0.2	.	0.1
	p	1.9	1.6	1.6	0.2	0.1	0.1
341	i	.	.	0.0	0.1	0.4	0.4
	p	.	.	0.1	0.5	0.5	0.8
342	i	0.1	0.3	0.1	0.1	0.3	0.2	.	0.3	0.5	0.3
	p	0.4	0.8	0.7	0.5	0.8	0.7	1.5	0.7	1.6	1.8
343	i	0.0	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.0	0.4
	p	0.1	0.2	0.1	0.4	0.3	0.4	0.4	0.4	1.0	1.7
344	i	0.2	0.4	0.2	.	.	0.0
	p	0.9	1.3	0.9	0.2	0.2	0.2	0.0	0.2	0.1	0.2	0.2	.	.	0.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
345	fear of mental disorder	i	273	0.0	0.0	0.0	.	.	.	0.0	0.0	.	.
		p	475	0.0	0.0	0.1	.	.	0.0	0.0	0.0	.	0.1
346	psychic disability/impairment	i	37	0.0	.	0.0	.	.	0.0
		p	517	0.0	0.0	0.0	.	.	0.0	0.0	0.0	0.2	.
347	psychic sympt./complt. NEC	i	7170	0.5	0.2	0.7	0.3	0.3	0.5	0.5	0.6	0.6	0.4
		p	24320	1.7	0.8	2.4	0.3	0.4	1.3	1.7	2.0	3.8	2.9
348	dementia (pre)senile	i	2793	0.2	0.2	0.2	.	.	.	0.0	0.0	0.8	2.2
		p	15776	1.1	0.7	1.5	.	.	.	0.1	0.1	3.0	15.5
349	organic psychosis	i	700	0.0	0.1	0.0	.	.	.	0.1	0.1	0.2	0.0
		p	3022	0.2	0.2	0.2	.	0.1	0.0	0.2	0.4	0.3	0.6
350	schizophrenia, all types	i	1487	0.1	0.1	0.1	.	.	0.1	0.1	0.1	0.2	0.4
		p	8650	0.6	0.5	0.6	.	.	0.2	0.7	0.9	1.3	1.3
351	affective psychosis	i	3147	0.2	0.2	0.3	.	.	0.1	0.2	0.4	0.3	0.5
		p	13570	0.9	0.7	1.1	.	0.1	0.1	0.9	2.0	1.8	1.7
352	unspecified psychosis NEC	i	2009	0.1	0.1	0.2	.	.	0.2	0.2	0.1	0.3	0.1
		p	16662	1.1	0.9	1.3	.	0.1	0.4	1.2	1.6	2.6	2.9
353	anxiety disorder/anxiety state	i	3609	0.2	0.2	0.3	0.0	0.1	0.0	0.3	0.3	0.4	0.7
		p	10877	0.7	0.5	1.0	0.1	0.1	0.2	0.7	1.3	1.8	1.8
354	hysterical/hypochondriacal dis.	i	4310	0.3	0.2	0.4	.	0.0	0.2	0.3	0.5	0.4	0.6
		p	21040	1.4	1.0	1.9	.	0.1	0.6	1.3	2.3	3.6	3.6
355	depressive disorder	i	33150	2.3	1.3	3.2	.	0.1	1.2	2.7	3.6	4.2	3.0
		p	121941	8.3	4.6	11.9	.	0.2	2.4	8.2	14.1	18.7	20.7
356	mental retardation	i	342	0.0	0.0	0.0	0.1	0.0	.	0.0	0.0	.	.
		p	10934	0.7	0.7	0.8	1.0	0.7	0.8	0.7	0.9	0.4	0.5
357	neurasthenia, surmenage	i	51155	3.5	3.0	3.9	.	0.3	3.9	5.7	3.9	1.6	1.7
		p	90831	6.2	5.5	6.9	.	0.4	5.3	9.6	8.4	4.0	3.5
358	personality disorder	i	739	0.1	0.1	0.0	.	.	0.2	0.1	0.0	0.1	.
		p	3385	0.2	0.2	0.3	.	.	0.2	0.3	0.3	0.2	0.2
359	phobia/neurotic disorder NEC	i	6214	0.4	0.3	0.5	0.2	0.1	0.4	0.6	0.5	0.2	0.5
		p	27812	1.9	1.3	2.5	0.2	0.2	1.1	2.9	2.5	2.2	1.7
360	mental/psychological dis. NEC	i	7248	0.5	0.5	0.5	0.1	0.3	0.5	0.5	0.8	0.3	0.2
		p	29762	2.0	1.6	2.5	0.4	0.4	1.2	2.3	3.3	2.9	2.5

P PSYCHOLOGICAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
345	i	0.0	0.1	.	0.1
	p	0.1	0.0	0.1	0.0	0.1	.	.	.	0.2
346	i	0.0
	p	0.1	0.0	0.0	0.1	0.0	0.2	0.2	.	.
347	i	0.1	0.4	0.1	0.6	0.1	0.8	0.4	0.7	0.3	0.8	0.1	1.0	0.1	0.6
	p	0.1	0.4	0.1	0.8	0.7	1.9	1.2	2.1	1.0	2.9	0.7	6.2	1.0	4.0
348	i	0.0	0.0	0.0	.	1.0	0.6	2.3	2.2
	p	0.0	0.1	0.2	0.1	2.9	3.1	12.0	17.4
349	i	0.1	0.1	0.1	0.0	0.3	0.1	0.1	.
	p	.	.	0.1	.	0.0	0.0	0.2	0.3	0.6	0.2	0.4	0.2	0.5	0.6
350	i	0.1	0.0	0.1	0.0	0.1	0.1	.	0.4	0.3	0.4
	p	0.2	0.2	0.7	0.6	0.9	1.0	0.8	1.7	1.2	1.3
351	i	0.2	.	0.1	0.3	0.5	0.4	0.1	0.5	0.3	0.7
	p	.	.	.	0.1	0.2	.	0.5	1.3	2.1	2.0	1.0	2.4	1.1	2.0
352	i	0.1	0.2	0.1	0.3	0.1	0.1	0.0	0.5	0.1	0.1
	p	.	.	.	0.1	0.4	0.4	1.1	1.2	1.6	1.6	1.3	3.6	1.4	3.7
353	i	.	0.0	0.0	0.1	0.0	0.1	0.1	0.5	0.3	0.3	0.6	0.2	0.1	1.0
	p	0.1	0.0	0.1	0.2	0.1	0.3	0.3	1.0	0.9	1.6	1.5	1.9	0.8	2.4
354	i	.	.	0.0	0.0	0.1	0.2	0.2	0.5	0.4	0.6	0.5	0.3	0.4	0.7
	p	.	.	0.1	0.1	0.5	0.7	1.0	1.7	1.5	3.1	2.8	4.2	2.2	4.3
355	i	.	.	0.1	0.2	0.4	2.0	1.6	3.8	2.2	4.9	2.6	5.4	1.6	3.8
	p	.	.	0.3	0.2	1.1	3.7	4.9	11.6	8.5	19.6	10.0	25.4	11.5	25.6
356	i	0.2	.	0.0	0.0	.	.	0.0	0.0	0.0	0.0
	p	1.0	1.0	0.8	0.5	0.8	0.9	0.7	0.7	0.6	1.2	0.6	0.3	1.0	0.2
357	i	.	.	0.4	0.1	2.5	5.3	5.3	6.0	3.5	4.3	0.8	2.3	1.3	2.0
	p	.	.	0.5	0.3	3.6	7.1	8.8	10.5	8.3	8.6	2.3	5.3	2.3	4.2
358	i	0.2	0.1	0.1	0.0	0.0	0.0	.	0.1	.	.
	p	0.3	0.1	0.3	0.4	0.2	0.5	0.0	0.4	.	0.4
359	i	0.1	0.2	0.1	0.2	0.4	0.4	0.5	0.7	0.5	0.6	.	0.4	0.1	0.6
	p	0.2	0.2	0.1	0.4	0.7	1.5	1.8	4.0	2.0	3.0	1.3	2.9	0.6	2.3
360	i	0.0	0.2	0.2	0.4	0.5	0.6	0.6	0.5	0.9	0.6	0.5	0.1	.	0.3
	p	0.5	0.2	0.4	0.5	0.7	1.7	1.7	3.0	2.9	3.7	2.7	3.1	1.1	3.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
361	pain attr. to respiratory system	i	318	0.0	0.0	0.0	.	0.0	.	0.0	0.0	.	0.2
		p	473	0.0	0.0	0.1	.	0.0	.	0.0	0.0	0.1	0.2
362	shortness of breath, dyspnoe	i	4288	0.3	0.3	0.3	0.1	0.3	0.2	0.1	0.3	1.0	0.9
		p	13798	0.9	1.0	0.9	0.5	0.5	0.5	0.5	1.0	3.2	3.4
363	wheezing	i	169	0.0	.	0.0	0.1	0.0	.	.	0.0	.	0.0
		p	667	0.0	0.1	0.0	0.2	0.0	0.0	0.1	0.0	.	0.1
364	breathing problems NEC	i	1507	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.1	0.1
		p	2256	0.2	0.2	0.1	0.1	0.2	0.1	0.2	0.1	0.3	0.2
365	cough	i	133305	9.1	7.3	10.9	20.3	9.1	6.5	6.7	9.8	12.1	10.6
		p	137700	9.4	7.6	11.1	20.6	9.2	6.6	6.9	10.4	12.9	11.6
366	epistaxis/nose bleed	i	7029	0.5	0.6	0.4	0.4	1.2	0.6	0.1	0.5	0.4	0.7
		p	8614	0.6	0.7	0.5	0.6	1.3	0.7	0.2	0.5	0.8	1.2
367	sneezing/nasal congestion	i	4379	0.3	0.3	0.3	0.5	0.3	0.4	0.3	0.3	0.3	0.1
		p	10744	0.7	0.7	0.8	1.6	0.6	0.5	0.8	0.7	0.6	0.6
368	sympt./complt. nose NEC	i	3843	0.3	0.2	0.3	0.4	0.3	0.4	0.2	0.2	0.1	0.0
		p	6051	0.4	0.4	0.4	0.4	0.4	0.6	0.4	0.6	0.1	0.1
369	sympt./complt. sinus (incl. pain)	i	1889	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.0	.
		p	2564	0.2	0.1	0.2	0.1	0.1	0.3	0.3	0.1	0.0	.
370	sympt./complt. throat	i	19498	1.3	1.1	1.6	0.7	1.2	1.7	1.4	1.3	1.1	0.8
		p	38324	2.6	2.2	3.0	1.1	1.9	3.1	2.8	3.0	2.4	2.0
371	sympt./complt. tonsils	i	834	0.1	0.0	0.1	0.3	0.2	0.0	0.0	0.0	.	.
		p	1185	0.1	0.1	0.1	0.5	0.2	0.0	0.0	0.1	.	.
372	sympt./complt. voice/vocal cords	i	4354	0.3	0.1	0.5	0.1	0.3	0.5	0.2	0.4	0.3	0.2
		p	6250	0.4	0.2	0.6	0.2	0.4	0.6	0.3	0.6	0.6	0.3
373	haemoptysis	i	772	0.1	0.1	0.0	.	.	0.0	0.1	0.1	0.1	0.2
		p	1052	0.1	0.1	0.1	.	.	0.0	0.1	0.1	0.1	0.2
374	abnormal sputum/phlegm NEC	i	2906	0.2	0.2	0.2	0.6	0.2	0.1	0.2	0.2	0.3	0.4
		p	9188	0.6	0.5	0.7	1.1	0.5	0.2	0.3	0.5	1.8	2.9
375	fear of respiratory disease	i	1919	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.2	0.1	0.1
		p	3676	0.3	0.3	0.2	0.3	0.1	0.1	0.3	0.3	0.2	0.1
376	disability/impairment resp. syst.	i	133	0.0	0.0	0.0	.	0.1	0.0
		p	208	0.0	0.0	0.0	.	0.1	0.0	0.0	0.0	.	.
377	sympt./complt. resp. syst. NEC	i	3827	0.3	0.3	0.2	0.8	0.3	0.3	0.2	0.2	0.3	0.4
		p	7855	0.5	0.6	0.4	1.1	0.5	0.6	0.3	0.4	1.1	0.9
378	tuberculosis resp. (excl. A70)	i	336	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	.	.
		p	2571	0.2	0.2	0.2	0.1	0.1	0.0	0.1	0.4	0.3	0.6
379	whooping cough	i	4454	0.3	0.2	0.4	1.8	0.8	0.2	0.1	0.1	0.2	0.0
		p	6848	0.5	0.4	0.5	2.9	1.4	0.2	0.1	0.1	0.2	0.0
380	scarlet fever	i	3667	0.2	0.3	0.2	1.9	1.0	0.0	0.0	.	.	.
		p	4235	0.3	0.3	0.3	2.3	1.1	0.0	0.0	.	.	.
381	streptoc. inf. throat NEC	i	857	0.1	0.0	0.1	0.0	0.1	0.1	0.1	0.1	0.0	.
		p	1054	0.1	0.0	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0
382	boil/abscess nose	i	768	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.1	.
		p	850	0.1	0.0	0.1	0.1	0.1	0.0	0.0	0.1	0.1	0.1
383	URI (head cold)	i	451996	30.8	29.2	32.3	105.6	33.4	31.0	26.4	21.3	20.3	15.0
		p	491827	33.5	31.3	35.6	109.9	35.4	32.9	28.4	24.7	25.5	18.6
384	sinusitis acute/chronic	i	127580	8.7	7.5	9.9	4.6	5.4	9.2	13.5	7.6	3.4	2.4
		p	157505	10.7	9.3	12.1	5.4	6.8	10.9	16.5	9.8	4.8	3.4
385	acute tonsillitis/adenoiditis	i	87855	6.0	5.8	6.2	20.1	10.8	7.1	4.8	2.7	1.0	1.1
		p	90927	6.2	6.0	6.4	20.3	11.1	7.3	5.0	2.8	1.4	1.2

R RESPIRATORY

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
361	i	.	.	.	0.0	.	.	.	0.1	0.0	0.2
	p	.	.	.	0.0	.	.	0.0	0.1	0.0	.	.	0.1	.	0.4
362	i	0.1	0.0	0.2	0.4	0.2	0.1	0.1	0.1	0.4	0.2	1.0	1.1	2.0	0.4
	p	0.4	0.5	0.4	0.7	0.5	0.4	0.4	0.6	1.3	0.8	3.0	3.3	6.1	2.0
363	i	.	0.2	.	0.0	0.0	.	.	.	0.0
	p	0.1	0.3	.	0.0	.	0.1	0.1	.	.	0.0	.	.	0.2	0.0
364	i	0.1	0.1	0.2	0.1	0.1	0.1	0.2	0.1	0.0	0.0	.	0.2	0.3	0.1
	p	0.1	0.1	0.2	0.2	0.1	0.1	0.2	0.1	0.1	0.0	0.3	0.3	0.3	0.1
365	i	18.0	22.7	7.6	10.6	5.5	7.5	4.5	8.9	7.2	12.5	12.6	11.6	11.3	9.8
	p	18.4	23.0	7.7	10.9	5.5	7.8	4.9	9.0	7.7	13.2	13.5	12.6	12.8	10.9
366	i	0.4	0.5	1.3	1.1	1.0	0.1	0.2	0.1	0.6	0.4	0.2	0.6	0.8	0.6
	p	0.7	0.5	1.4	1.2	1.2	0.2	0.2	0.1	0.6	0.4	0.4	1.1	1.3	1.1
367	i	0.7	0.3	0.3	0.3	0.2	0.6	0.3	0.2	0.2	0.4	0.4	0.3	.	0.1
	p	1.2	2.0	0.6	0.6	0.4	0.6	0.8	0.8	0.6	0.7	0.4	0.8	0.5	0.6
368	i	0.3	0.6	0.1	0.6	0.5	0.3	0.2	0.2	0.3	0.2	0.1	0.1	0.0	0.0
	p	0.3	0.6	0.1	0.6	0.7	0.5	0.3	0.4	0.6	0.5	0.2	0.1	0.1	0.0
369	i	.	0.2	0.1	0.0	0.1	0.2	0.2	0.3	0.0	0.1	.	0.0	.	.
	p	.	0.2	0.1	0.1	0.2	0.3	0.2	0.3	0.0	0.2	.	0.0	.	.
370	i	0.3	1.1	0.8	1.6	1.2	2.3	1.3	1.5	1.2	1.4	0.3	1.7	0.6	0.9
	p	0.6	1.6	1.2	2.6	2.7	3.6	2.6	3.1	2.6	3.4	1.6	3.0	1.3	2.4
371	i	0.1	0.6	0.2	0.1	.	0.1	0.0	0.0	0.0	0.1
	p	0.3	0.7	0.2	0.3	0.0	0.1	0.0	0.0	0.0	0.1
372	i	0.1	0.2	0.2	0.5	0.1	0.9	0.1	0.3	0.2	0.5	0.3	0.2	.	0.3
	p	0.1	0.2	0.4	0.5	0.1	1.1	0.1	0.4	0.3	0.8	0.8	0.5	0.0	0.4
373	i	0.1	.	0.1	0.0	0.1	0.1	0.2	.	0.2	0.1
	p	0.1	.	0.1	0.1	0.2	0.1	0.2	.	0.2	0.2
374	i	0.8	0.5	0.2	0.2	0.0	0.1	0.2	0.1	0.1	0.2	0.4	0.2	0.5	0.4
	p	1.0	1.2	0.3	0.8	0.1	0.2	0.2	0.3	0.4	0.5	2.1	1.6	3.1	2.7
375	i	0.1	.	.	0.1	0.1	0.2	0.1	0.1	0.3	0.2	0.1	0.0	.	0.2
	p	0.6	0.0	0.1	0.1	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.1	.	0.2
376	i	.	.	0.1	0.0	0.0
	p	.	.	0.1	0.0	0.0	.	.	0.0	.	0.0
377	i	1.1	0.5	0.4	0.2	0.2	0.3	0.1	0.2	0.2	0.2	0.5	0.1	0.5	0.3
	p	1.5	0.8	0.6	0.4	0.7	0.5	0.3	0.4	0.6	0.3	2.0	0.3	0.8	0.9
378	i	.	0.1	0.0	.	.	0.0	0.0	.	0.0	0.0
	p	.	0.2	0.0	0.1	.	0.0	0.1	0.0	0.5	0.3	0.1	0.5	0.3	0.7
379	i	1.1	2.5	0.9	0.8	0.0	0.3	0.1	0.1	0.1	0.1	0.2	0.1	.	0.1
	p	1.7	4.1	1.7	1.1	0.0	0.5	0.1	0.1	0.1	0.1	0.2	0.2	.	0.1
380	i	2.0	1.8	1.1	0.9	0.0	0.0	.	0.0
	p	2.5	2.1	1.2	1.0	0.0	0.0	.	0.0
381	i	0.0	.	0.0	0.2	.	0.2	0.0	0.1	0.1	0.0	.	0.0	.	.
	p	0.2	.	0.0	0.2	.	0.2	0.0	0.1	0.1	0.0	.	0.0	.	0.0
382	i	.	0.1	.	0.1	0.0	.	0.0	0.1	0.1	0.1	0.2	.	.	.
	p	.	0.1	.	0.1	0.0	.	0.0	0.1	0.1	0.1	0.2	.	.	0.1
383	i	105.4	105.7	31.1	35.8	26.5	35.7	24.0	28.8	20.7	21.9	19.5	21.0	16.2	14.4
	p	109.6	110.1	33.0	37.9	28.0	38.0	25.5	31.5	23.3	26.0	22.9	27.6	19.4	18.2
384	i	4.0	5.2	5.0	5.7	6.8	11.6	11.4	15.6	6.7	8.6	2.6	4.1	2.9	2.2
	p	4.8	6.0	6.9	6.6	8.2	13.6	13.9	19.2	8.7	10.8	4.0	5.5	2.9	3.7
385	i	20.6	19.6	10.4	11.3	5.7	8.5	4.7	4.8	2.5	2.9	0.6	1.4	1.5	1.0
	p	20.8	19.8	10.6	11.6	5.9	8.7	4.9	5.1	2.6	3.1	1.0	1.6	1.6	1.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
386	acute laryngitis/tracheitis/croup	i	51964	3.5	3.3	3.8	9.2	3.3	3.2	3.4	3.2	2.6	2.2
		p	54709	3.7	3.5	4.0	9.9	3.4	3.2	3.6	3.5	2.8	2.3
387	acute bronchi(ol)itis	i	141965	9.7	9.8	9.6	30.9	10.1	6.4	6.9	9.5	10.2	11.1
		p	191330	13.0	13.3	12.8	36.6	13.2	8.1	9.0	13.9	16.5	18.0
388	influenza w/o pneumonia	i	90889	6.2	6.0	6.3	7.7	4.0	5.2	7.0	7.2	6.0	4.9
		p	92355	6.3	6.1	6.5	7.7	4.1	5.2	7.1	7.4	6.1	5.1
389	pneumonia	i	30974	2.1	2.3	1.9	6.0	2.4	0.9	1.3	1.8	3.8	5.0
		p	32449	2.2	2.4	2.0	6.0	2.4	0.9	1.4	1.8	3.9	5.5
390	pleurisy all (excl. R70)	i	781	0.1	0.1	0.0	.	.	0.1	0.1	0.1	0.1	.
		p	1585	0.1	0.1	0.1	.	.	0.1	0.1	0.1	0.4	0.1
391	infection resp. syst. NEC	i	10809	0.7	0.8	0.7	2.2	0.8	0.5	0.6	0.6	0.8	1.4
		p	18663	1.3	1.4	1.2	3.0	1.1	0.7	1.0	1.4	1.7	1.9
392	mal. neopl. bronchus/lung	i	1530	0.1	0.2	0.0	0.2	0.4	0.5
		p	14823	1.0	1.7	0.3	.	.	.	0.2	1.6	5.3	4.8
393	mal. neopl. resp. syst. NEC	i	554	0.0	0.0	0.0	.	.	0.0	0.0	0.1	0.1	0.1
		p	3772	0.3	0.4	0.2	.	.	0.0	0.0	0.5	0.9	1.1
394	ben. neopl. resp. syst.	i	379	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.1	0.1	.
		p	900	0.1	0.0	0.1	0.1	0.0	0.1	0.0	0.1	0.2	.
395	unspecified neopl. resp. syst.	i	468	0.0	0.1	0.0	.	.	.	0.0	0.0	0.0	0.2
		p	2055	0.1	0.1	0.2	.	0.0	0.0	0.1	0.2	0.5	0.5
396	foreign body resp. syst.	i	808	0.1	0.1	0.0	0.5	0.1	0.0	0.0	0.0	.	0.1
		p	911	0.1	0.1	0.0	0.6	0.1	0.0	0.0	0.0	.	0.1
397	injuries resp. syst. NEC	i	648	0.0	0.0	0.1	0.0	0.1	0.0	0.1	0.0	.	.
		p	698	0.0	0.0	0.1	0.0	0.1	0.1	0.1	0.0	.	.
398	cong. anom. resp. syst.	i	115	0.0	0.0	.	0.0	.	0.0	0.0	.	0.0	.
		p	445	0.0	0.0	0.0	0.1	.	0.0	0.0	0.1	0.0	.
399	hypertrophy/chron. inf. tonsills/aden.	i	9393	0.6	0.5	0.7	5.8	1.0	0.4	0.2	0.0	0.1	.
		p	14342	1.0	0.9	1.1	8.2	1.9	0.7	0.3	0.0	0.1	0.0
400	chronic bronchitis	i	14787	1.0	1.2	0.8	1.6	0.6	0.5	0.5	1.3	2.0	3.4
		p	153518	10.5	12.8	8.2	8.8	7.1	4.9	4.7	13.8	28.2	35.6
401	pleural effusion NOS	i	1026	0.1	0.1	0.1	.	0.1	0.0	0.0	0.1	0.4	.
		p	1529	0.1	0.1	0.1	.	0.1	0.0	0.0	0.2	0.5	0.1
402	emfysema	i	3008	0.2	0.3	0.1	.	.	0.1	0.0	0.4	0.8	1.1
		p	46250	3.2	4.9	1.5	.	0.1	0.1	0.4	3.6	16.6	20.1
403	asthma	i	26701	1.8	1.9	1.7	3.6	3.1	1.6	1.3	1.3	1.9	2.1
		p	149170	10.2	10.6	9.7	14.6	13.2	7.2	6.6	9.9	18.5	18.2
404	allergic rhinitis/hay fever	i	31500	2.1	1.9	2.4	0.8	1.9	3.5	2.9	1.1	1.1	0.8
		p	96600	6.6	6.5	6.6	1.9	7.1	10.1	8.2	4.4	3.0	2.8
405	diseases resp. syst. NEC	i	8464	0.6	0.7	0.5	0.6	0.3	0.6	0.7	0.7	0.5	0.5
		p	21915	1.5	1.9	1.1	0.9	0.6	1.3	1.4	1.9	3.1	1.9

R RESPIRATORY

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
386	i	9.3	9.1	3.5	3.1	2.9	3.5	3.0	3.8	2.8	3.6	2.5	2.6	1.3	2.7
	p	10.2	9.6	3.5	3.2	2.8	3.5	3.1	4.1	3.1	3.9	2.8	2.8	1.5	2.7
387	i	32.7	29.1	10.5	9.7	6.4	6.5	6.4	7.5	9.9	9.2	10.4	10.0	13.5	9.9
	p	37.6	35.6	13.8	12.6	8.1	8.0	8.5	9.5	14.8	13.0	16.7	16.4	24.1	14.8
388	i	8.8	6.4	3.4	4.7	5.0	5.3	7.4	6.6	6.6	7.9	3.8	7.8	3.9	5.4
	p	8.8	6.5	3.5	4.7	5.0	5.3	7.5	6.6	6.8	8.0	3.8	7.9	3.9	5.8
389	i	6.4	5.3	2.7	2.0	1.0	0.8	1.3	1.2	1.9	1.7	4.6	2.9	7.1	2.9
	p	6.5	5.5	2.7	2.1	1.0	0.8	1.5	1.3	1.9	1.8	4.8	3.1	8.0	4.1
390	i	0.1	0.0	0.1	0.0	0.1	0.0	0.2	0.1	.	.
	p	0.2	0.0	0.1	0.1	0.1	0.1	0.2	0.6	.	0.2
391	i	2.4	2.0	0.7	0.9	0.6	0.3	0.6	0.5	0.5	0.6	0.9	0.6	2.3	0.9
	p	3.1	2.9	1.0	1.3	0.9	0.5	1.2	0.9	1.3	1.4	2.1	1.4	3.0	1.3
392	i	0.5	.	0.6	0.0	0.9	0.1
	p	0.1	0.2	2.8	0.4	11.1	0.9	10.7	1.7
393	i	0.1	.	0.0	0.1	0.0	0.3	.	.	0.1
	p	0.1	0.0	0.1	0.8	0.3	1.7	0.3	2.4	0.5
394	i	.	0.2	0.0	.	0.0	.	0.0	.	0.0	0.1	0.1	0.0	.	.
	p	.	0.2	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.2	.	.
395	i	0.0	0.0	0.1	.	0.1	.	0.6	.
	p	.	.	0.0	.	0.1	.	0.0	0.2	0.3	0.1	0.2	0.7	0.6	0.5
396	i	0.7	0.4	0.2	0.0	0.0	.	.	0.0	.	0.0	.	.	0.0	0.1
	p	0.7	0.5	0.2	0.0	0.0	.	.	0.0	0.0	0.0	.	.	0.0	0.1
397	i	0.0	.	0.0	0.2	0.0	0.1	0.0	0.1	.	0.0
	p	0.0	.	0.0	0.2	0.0	0.1	0.0	0.1	.	0.0
398	i	0.0	.	.	.	0.0	.	0.0	.	.	.	0.1	.	.	.
	p	0.0	0.1	.	.	0.0	0.0	0.0	.	.	0.1	0.1	.	.	.
399	i	5.3	6.4	0.7	1.3	0.4	0.4	0.1	0.3	0.0	0.0	.	0.2	.	.
	p	8.8	7.6	1.6	2.3	0.4	0.9	0.2	0.5	0.0	0.1	.	0.2	0.1	.
400	i	1.9	1.4	0.9	0.3	0.6	0.5	0.6	0.4	1.5	1.1	2.2	1.9	6.9	1.4
	p	10.8	6.8	8.6	5.5	5.4	4.4	5.1	4.4	17.3	10.5	40.2	19.0	61.9	21.6
401	i	.	.	.	0.1	0.0	.	0.0	0.0	0.1	0.2	0.5	0.2	.	.
	p	.	.	0.1	0.1	0.0	.	0.0	0.0	0.2	0.3	0.7	0.3	0.1	0.1
402	i	0.1	.	0.0	0.0	0.5	0.2	1.7	0.1	2.9	0.1
	p	.	.	0.1	0.2	0.2	0.0	0.5	0.3	5.9	1.4	30.8	5.6	39.9	9.5
403	i	4.5	2.7	4.1	2.2	1.4	1.8	1.0	1.5	1.1	1.6	2.6	1.4	2.8	1.8
	p	18.6	10.5	15.6	10.8	7.1	7.4	5.5	7.6	9.7	10.2	23.4	14.8	24.9	14.7
404	i	0.9	0.6	1.8	2.0	2.9	4.1	2.6	3.3	0.9	1.2	1.0	1.1	0.6	1.0
	p	1.7	2.2	7.6	6.6	10.4	9.7	7.7	8.7	3.8	4.9	2.7	3.2	3.1	2.6
405	i	0.7	0.5	0.2	0.3	0.7	0.5	0.8	0.6	1.0	0.3	0.6	0.5	0.5	0.5
	p	1.3	0.5	0.6	0.6	1.8	0.7	1.5	1.2	2.3	1.6	4.9	1.7	3.7	0.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
406	pain/tenderness of skin	i	834	0.1	0.0	0.1	0.1	0.1	0.0	0.1	0.0	.	0.1
		p	1586	0.1	0.1	0.1	0.1	0.1	0.2	0.1	.	.	0.2
407	pruritus, skin itching	i	12621	0.9	0.7	1.0	0.7	0.6	0.5	0.9	0.9	1.6	1.2
		p	28063	1.9	1.3	2.5	0.9	0.7	0.8	1.6	2.8	3.6	5.9
408	warts	i	73978	5.0	4.5	5.6	2.7	15.2	7.0	3.1	2.4	2.2	2.2
		p	99882	6.8	6.0	7.6	3.2	21.2	9.1	4.2	3.2	3.3	3.0
409	condyloma acuminata	i	5112	0.3	0.5	0.2	0.1	0.2	0.7	0.5	0.2	0.0	.
		p	5166	0.4	0.5	0.2	0.1	0.2	0.6	0.5	0.2	0.0	.
410	local swelling/papul/lump	i	7071	0.5	0.4	0.6	0.5	0.5	0.6	0.6	0.4	0.4	0.2
		p	9410	0.6	0.5	0.8	0.6	0.7	0.7	0.7	0.5	0.4	0.7
411	multiple swelling/papul/lump	i	1680	0.1	0.1	0.2	0.3	0.0	0.1	0.1	0.2	0.1	0.1
		p	3065	0.2	0.2	0.3	0.5	0.3	0.3	0.2	0.2	0.1	0.1
412	local redness/erythema	i	9498	0.6	0.5	0.8	1.7	0.8	0.6	0.5	0.5	0.8	0.6
		p	16733	1.1	0.9	1.4	2.3	1.0	1.0	0.9	1.2	1.2	1.4
413	general redness/erythema	i	2917	0.2	0.2	0.2	0.8	0.3	0.2	0.1	0.1	0.2	.
		p	6051	0.4	0.3	0.5	1.0	0.3	0.3	0.3	0.5	0.4	0.5
414	changes in skin color NEC	i	1964	0.1	0.1	0.1	0.1	0.2	0.1	0.1	0.1	0.3	.
		p	2680	0.2	0.1	0.2	0.1	0.2	0.2	0.2	0.2	0.5	0.0
415	local infection finger/toe	i	22399	1.5	1.6	1.5	1.3	2.1	1.5	1.6	1.4	0.8	1.3
		p	23753	1.6	1.6	1.6	1.6	2.2	1.5	1.7	1.5	0.9	1.5
416	boil/carbuncle/cellulitis	i	28261	1.9	2.1	1.7	1.3	1.3	2.8	2.2	1.8	1.3	1.2
		p	34636	2.4	2.5	2.2	1.5	1.7	3.1	2.9	2.2	1.5	1.5
417	local infection skin NEC	i	17679	1.2	1.2	1.2	1.6	1.0	1.3	1.2	1.1	1.4	1.3
		p	24033	1.6	1.5	1.7	2.2	1.3	1.5	1.6	1.6	1.9	2.6
418	insect bite	i	16605	1.1	0.9	1.4	1.9	1.6	1.5	0.9	0.8	0.9	0.9
		p	17969	1.2	0.9	1.5	1.9	1.6	1.7	1.0	0.8	1.0	0.9
419	animal/human bite	i	8158	0.6	0.6	0.5	0.6	0.6	0.6	0.6	0.8	0.1	0.0
		p	9045	0.6	0.7	0.6	0.6	0.6	0.6	0.6	0.9	0.2	0.1
420	burns/scalds	i	11661	0.8	0.8	0.8	1.4	0.6	1.3	0.7	0.6	0.7	0.5
		p	16469	1.1	1.1	1.1	2.1	0.8	1.8	1.0	1.0	0.8	0.7
421	foreign body in skin	i	6889	0.5	0.5	0.4	0.6	0.5	0.6	0.6	0.4	0.2	0.2
		p	7920	0.5	0.6	0.4	0.6	0.5	0.6	0.7	0.4	0.3	0.2
422	bruise/contusion intact skin	i	29953	2.0	1.6	2.5	1.6	1.7	2.8	1.5	2.0	2.7	3.1
		p	33812	2.3	1.8	2.8	1.6	1.9	2.9	1.7	2.4	3.2	4.0
423	abrasion/scratch/blister	i	13784	0.9	1.0	0.9	0.6	1.7	1.6	0.5	0.5	1.0	1.7
		p	14985	1.0	1.1	0.9	0.7	1.8	1.6	0.5	0.6	1.1	2.1
424	laceration/cut	i	54115	3.7	5.1	2.3	6.9	5.4	4.9	3.2	2.2	1.8	2.8
		p	59890	4.1	5.5	2.7	7.1	5.8	5.3	3.8	2.5	2.2	3.2
425	injury to skin NEC	i	27703	1.9	2.3	1.5	4.0	2.6	2.1	1.6	1.1	1.7	1.9
		p	42007	2.9	3.4	2.3	5.1	3.6	3.1	2.3	2.1	3.1	4.2
426	corns/callosities	i	12557	0.9	0.7	1.0	0.3	0.6	0.7	0.9	0.8	1.1	2.2
		p	15616	1.1	0.8	1.3	0.3	0.7	0.8	1.1	1.1	1.5	2.7
427	sympt./complt. of skin texture	i	8041	0.5	0.4	0.7	0.8	0.5	0.4	0.6	0.4	0.6	1.2
		p	19511	1.3	1.0	1.7	2.2	0.8	0.8	1.1	0.9	1.9	5.3
428	sympt./complt. nails	i	2420	0.2	0.2	0.1	0.1	0.3	0.2	0.2	0.1	0.2	0.0
		p	3908	0.3	0.2	0.3	0.2	0.4	0.2	0.3	0.2	0.4	0.3
429	losing hair/baldness	i	3319	0.2	0.2	0.3	.	0.1	0.3	0.4	0.1	0.2	0.2
		p	7763	0.5	0.3	0.7	.	0.1	0.4	0.8	0.5	0.9	0.7
430	sympt./complt. hair NEC	i	1969	0.1	0.0	0.2	0.0	0.0	0.2	0.2	0.1	.	0.0
		p	5288	0.4	0.2	0.5	0.1	0.0	0.4	0.5	0.4	0.4	0.2

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
406	i	.	0.2	0.0	0.1	0.1	0.0	0.0	0.1	.	0.1	.	.	0.0	0.2
	p	.	0.2	0.0	0.1	0.1	0.0	0.1	0.2	0.1	0.1	.	.	0.1	0.3
407	i	0.5	0.9	0.6	0.6	0.5	0.6	0.6	1.2	0.8	1.1	1.1	1.9	1.2	1.2
	p	0.7	1.1	0.8	0.6	0.7	1.0	0.9	2.2	2.2	3.4	2.6	4.4	4.1	6.8
408	i	2.5	2.9	13.6	16.8	6.8	7.2	2.3	4.0	2.1	2.7	1.6	2.6	1.9	2.4
	p	2.7	3.8	19.2	23.2	8.3	10.0	3.1	5.3	2.7	3.8	3.0	3.6	2.3	3.3
409	i	0.2	0.0	0.2	0.2	0.9	0.5	0.8	0.2	0.1	0.2	0.1	.	.	.
	p	0.2	0.0	0.2	0.2	0.9	0.4	0.8	0.2	0.1	0.2	0.1	.	.	.
410	i	0.4	0.6	0.5	0.5	0.4	0.8	0.4	0.7	0.3	0.4	0.2	0.5	0.1	0.2
	p	0.4	0.8	0.8	0.6	0.5	0.9	0.5	0.9	0.5	0.6	0.2	0.5	0.4	0.9
411	i	0.2	0.4	.	0.1	.	0.2	0.0	0.2	0.1	0.2	.	0.1	0.2	.
	p	0.6	0.4	0.3	0.3	0.2	0.4	0.1	0.3	0.1	0.2	.	0.2	0.2	.
412	i	1.5	1.9	0.5	1.0	0.3	0.9	0.3	0.7	0.4	0.6	0.6	0.9	0.4	0.7
	p	2.5	2.2	0.6	1.4	0.6	1.3	0.5	1.4	1.2	1.3	1.0	1.3	1.8	1.3
413	i	0.5	1.0	0.2	0.5	0.2	0.2	0.1	0.1	0.0	0.3	0.4	.	.	.
	p	0.6	1.3	0.2	0.5	0.3	0.3	0.3	0.3	0.3	0.7	0.4	0.4	0.6	0.5
414	i	.	0.1	0.4	0.1	0.1	0.1	0.0	0.1	0.1	0.2	0.5	0.2	.	.
	p	.	0.1	0.4	0.1	0.1	0.2	0.1	0.3	0.1	0.2	0.5	0.5	0.0	.
415	i	1.4	1.2	2.3	1.9	1.8	1.2	1.5	1.7	1.6	1.3	0.2	1.3	1.2	1.4
	p	1.5	1.7	2.4	2.0	1.8	1.1	1.5	1.9	1.6	1.4	0.2	1.5	1.6	1.4
416	i	1.8	0.7	1.6	0.9	3.2	2.4	2.2	2.2	2.0	1.5	1.0	1.4	0.9	1.4
	p	1.8	1.3	2.1	1.2	3.5	2.7	2.8	2.9	2.2	2.1	1.4	1.6	1.2	1.7
417	i	1.8	1.4	1.1	0.9	1.2	1.4	1.0	1.3	1.4	0.8	0.9	1.8	1.7	1.1
	p	2.3	2.0	1.3	1.2	1.5	1.6	1.4	1.8	1.8	1.4	0.9	2.6	2.3	2.7
418	i	2.0	1.8	1.2	2.0	1.3	1.8	0.7	1.2	0.5	1.0	0.2	1.4	0.4	1.1
	p	2.0	1.8	1.2	2.1	1.3	2.1	0.7	1.3	0.5	1.2	0.5	1.5	0.4	1.1
419	i	0.8	0.4	0.5	0.6	0.8	0.4	0.5	0.7	1.1	0.5	0.2	.	0.1	0.0
	p	0.8	0.4	0.6	0.7	0.8	0.4	0.5	0.8	1.2	0.6	0.3	0.2	0.3	0.0
420	i	1.5	1.2	0.7	0.4	1.2	1.4	0.8	0.6	0.3	0.9	0.6	0.7	0.6	0.4
	p	2.5	1.7	0.9	0.6	1.5	2.1	1.1	0.8	0.6	1.4	0.6	0.9	1.1	0.5
421	i	0.7	0.6	0.5	0.5	0.7	0.4	0.6	0.5	0.4	0.4	0.2	0.2	.	0.2
	p	0.7	0.6	0.6	0.5	0.8	0.4	0.8	0.5	0.4	0.4	0.2	0.3	.	0.2
422	i	1.8	1.4	1.8	1.6	2.2	3.4	1.4	1.7	1.2	2.9	2.2	3.1	1.0	4.2
	p	1.8	1.4	2.0	1.9	2.3	3.6	1.6	1.9	1.3	3.4	2.7	3.6	2.1	5.1
423	i	0.5	0.7	2.2	1.2	2.0	1.2	0.6	0.4	0.4	0.6	0.6	1.2	0.9	2.1
	p	0.5	1.0	2.3	1.2	2.0	1.2	0.6	0.4	0.4	0.8	0.8	1.4	2.1	2.1
424	i	8.7	5.0	8.0	2.8	7.3	2.5	4.3	2.1	2.8	1.6	2.5	1.3	1.7	3.3
	p	8.7	5.5	8.5	3.0	7.8	2.7	5.0	2.5	3.2	1.7	2.7	1.9	1.7	4.1
425	i	4.9	3.1	3.6	1.6	2.3	1.9	1.9	1.3	1.3	0.9	1.7	1.8	2.2	1.7
	p	6.5	3.7	4.9	2.2	3.6	2.5	2.7	1.8	2.6	1.6	2.8	3.3	3.5	4.6
426	i	0.2	0.4	0.5	0.8	0.3	1.1	0.8	1.0	0.6	0.9	1.5	0.9	1.5	2.6
	p	0.2	0.4	0.5	0.8	0.4	1.2	1.0	1.2	0.8	1.4	1.7	1.4	2.0	3.1
427	i	1.1	0.5	0.6	0.4	0.1	0.7	0.4	0.8	0.2	0.5	0.5	0.7	0.9	1.3
	p	2.1	2.3	1.0	0.6	0.6	1.1	0.9	1.3	0.6	1.2	1.4	2.4	3.0	6.5
428	i	0.1	0.2	0.4	0.2	0.3	0.1	0.2	0.2	0.0	0.2	.	0.3	0.0	0.0
	p	0.1	0.3	0.4	0.3	0.3	0.1	0.3	0.2	0.2	0.2	.	0.8	0.0	0.5
429	i	.	.	0.1	.	0.1	0.5	0.4	0.5	.	0.1	.	0.4	.	0.3
	p	.	.	0.1	.	0.4	0.5	0.7	0.8	0.2	0.9	.	1.6	0.1	1.0
430	i	.	0.0	0.0	0.0	0.0	0.5	0.1	0.4	0.1	0.1	.	.	.	0.0
	p	.	0.2	0.0	0.0	0.0	0.8	0.3	0.7	0.3	0.5	0.5	0.3	0.0	0.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
431	fear of skin disease	i	389	0.0	0.0	0.0	.	0.0	0.0	0.0	0.0	0.1	.
		p	849	0.1	0.0	0.1	.	0.0	0.1	0.1	0.1	0.1	.
432	disability/impairment of skin	i	0
		p	125	0.0	0.0	0.0	.	.	.	0.0	0.0	.	0.1
433	sympt./complt. of skin NEC	i	4226	0.3	0.2	0.4	0.6	0.2	0.4	0.3	0.3	0.1	0.2
		p	9419	0.6	0.6	0.7	0.8	0.3	0.6	0.7	0.6	0.8	0.7
434	herpes zoster	i	14617	1.0	0.8	1.2	0.2	0.8	0.7	0.7	1.3	2.0	2.4
		p	19029	1.3	1.0	1.6	0.2	0.8	0.7	0.9	1.7	3.2	4.1
435	herpes (simplex) lip/mouth	i	10565	0.7	0.3	1.1	0.2	0.6	1.2	0.8	0.6	0.7	0.2
		p	19915	1.4	0.7	2.0	0.3	1.0	2.2	1.5	1.4	0.8	0.3
436	herpes simplex genitalis	i	1701	0.1	0.1	0.1	.	0.0	0.3	0.1	0.1	.	.
		p	2471	0.2	0.1	0.2	.	0.0	0.3	0.2	0.2	.	.
437	herpes simplex NEC	i	2873	0.2	0.1	0.3	0.5	0.2	0.3	0.2	0.1	0.0	0.1
		p	4824	0.3	0.2	0.5	0.5	0.2	0.4	0.3	0.4	0.3	0.2
438	scabies/other ascariasis	i	2353	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	.	0.1
		p	3174	0.2	0.2	0.2	0.1	0.3	0.3	0.3	0.1	0.1	0.1
439	pediculosis/skininfestation	i	5307	0.4	0.4	0.4	0.4	0.7	0.8	0.3	0.1	.	0.2
		p	6780	0.5	0.5	0.5	0.5	1.1	0.9	0.3	0.1	0.0	0.2
440	dermatophytosis/dermatomycosis	i	84885	5.8	6.4	5.2	4.9	5.7	7.0	6.7	5.3	3.2	3.3
		p	123444	8.4	9.2	7.7	6.0	6.9	9.1	9.4	9.2	6.4	6.6
441	moniliasis/candidiasis (excl. GU)	i	12989	0.9	0.9	0.8	3.8	0.5	0.6	0.7	0.7	1.1	0.8
		p	19224	1.3	1.4	1.2	4.9	0.5	1.0	0.9	1.4	1.9	1.6
442	erysipelas	i	11164	0.8	0.7	0.9	0.1	0.2	0.4	0.4	1.3	1.7	2.5
		p	14689	1.0	0.8	1.2	0.1	0.2	0.5	0.6	1.8	2.3	3.2
443	mollusca contagiosa	i	10726	0.7	0.8	0.7	2.3	4.4	0.1	0.0	0.0	.	.
		p	11947	0.8	0.8	0.8	2.5	4.9	0.2	0.0	0.0	.	.
444	infection skin NEC	i	8808	0.6	0.6	0.6	1.1	0.6	0.5	0.7	0.5	0.4	0.5
		p	11970	0.8	0.8	0.9	1.3	0.9	0.6	0.9	0.6	0.9	1.0
445	malign. neoplasm skin	i	3582	0.2	0.3	0.2	.	.	0.0	0.1	0.5	0.8	0.8
		p	8892	0.6	0.6	0.6	.	.	0.1	0.3	0.9	2.2	2.7
446	lipoma	i	12344	0.8	1.1	0.6	0.1	0.2	0.6	1.3	0.9	1.3	0.3
		p	14566	1.0	1.2	0.8	0.1	0.3	0.7	1.5	1.1	1.5	0.5
447	benign. neoplasm skin NEC	i	21247	1.4	1.2	1.7	0.7	1.1	1.2	2.0	1.6	1.1	0.7
		p	25230	1.7	1.4	2.0	0.9	1.1	1.4	2.3	1.8	1.7	1.1
448	unspecified neoplasm skin	i	1427	0.1	0.1	0.1	.	0.0	0.1	0.1	0.1	0.2	0.3
		p	2006	0.1	0.2	0.1	.	0.1	0.1	0.1	0.1	0.2	0.4
449	hemangioma/lymfangioma	i	2246	0.2	0.1	0.2	0.5	0.2	0.0	0.1	0.3	0.1	.
		p	2488	0.2	0.1	0.2	0.5	0.2	0.0	0.1	0.3	0.1	.
450	nevus/mole	i	11790	0.8	0.5	1.1	0.4	1.0	1.1	1.1	0.4	0.4	0.2
		p	15283	1.0	0.6	1.5	0.4	1.4	1.3	1.4	0.7	0.5	0.4
451	cong. anom. of skin NEC	i	260	0.0	0.0	0.0	0.0
		p	443	0.0	0.0	0.0	0.1
452	impetigo	i	21724	1.5	1.4	1.5	7.4	3.7	1.3	0.7	0.4	0.3	0.4
		p	22357	1.5	1.5	1.6	7.6	3.7	1.3	0.8	0.4	0.3	0.4
453	pilonidal cyst/fistula	i	1323	0.1	0.1	0.1	.	.	0.1	0.1	0.1	.	0.0
		p	2316	0.2	0.2	0.1	.	.	0.2	0.2	0.2	0.0	0.4
454	seborrheic dermatitis	i	15261	1.0	1.2	0.9	1.4	0.7	1.0	1.4	0.9	0.7	0.4
		p	23454	1.6	1.7	1.5	1.6	0.8	1.4	2.3	1.4	1.3	1.1
455	atopic dermatitis/eczema	i	16862	1.1	1.0	1.3	6.0	2.0	1.2	0.6	0.4	0.2	0.7
		p	31750	2.2	1.9	2.4	12.2	3.3	2.1	1.2	0.8	0.5	1.2

S SKIN

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
431	i	.	.	.	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.3	.	.	.
	p	.	.	.	0.0	0.1	0.1	0.0	0.1	0.1	0.1	0.3	.	.	.
432	i
	p	0.0	.	.	0.0	.	.	.	0.2
433	i	0.5	0.6	0.0	0.3	0.1	0.7	0.4	0.2	0.1	0.4	0.1	0.2	0.5	.
	p	0.9	0.8	0.1	0.4	0.2	1.0	0.8	0.7	0.4	0.8	0.9	0.7	1.1	0.6
434	i	.	0.4	0.7	0.8	0.5	0.8	0.7	0.8	0.8	1.8	1.8	2.2	2.9	2.1
	p	.	0.4	0.7	0.9	0.5	0.9	0.8	1.0	1.1	2.2	3.0	3.4	4.1	4.0
435	i	0.1	0.3	0.3	0.9	0.4	2.0	0.4	1.2	0.3	0.9	0.4	0.9	0.2	0.2
	p	0.1	0.5	0.5	1.6	0.8	3.7	0.9	2.2	0.7	2.0	0.6	1.0	0.2	0.4
436	i	.	.	.	0.0	0.2	0.3	0.1	0.2	0.1	0.2
	p	.	.	.	0.0	0.2	0.4	0.1	0.4	0.2	0.2
437	i	0.7	0.3	0.1	0.3	0.1	0.5	0.1	0.3	0.0	0.2	0.0	0.0	.	0.1
	p	0.7	0.3	0.1	0.3	0.2	0.6	0.2	0.5	0.0	0.7	0.0	0.5	.	0.3
438	i	0.2	0.1	0.2	0.2	0.1	0.3	0.2	0.2	0.1	0.1	.	.	0.2	.
	p	0.2	0.1	0.2	0.4	0.2	0.3	0.2	0.3	0.2	0.1	0.2	.	0.2	.
439	i	0.6	0.1	0.2	1.2	0.9	0.7	0.4	0.2	0.0	0.1	.	.	0.0	0.3
	p	0.6	0.3	0.5	1.8	1.0	0.7	0.5	0.2	0.2	0.1	.	0.0	0.0	0.3
440	i	4.1	5.8	5.6	5.8	7.8	6.1	7.6	5.7	6.0	4.5	3.4	3.0	3.4	3.2
	p	5.3	6.8	7.0	6.7	9.9	8.3	10.5	8.2	10.0	8.3	6.1	6.6	8.1	5.9
441	i	4.1	3.5	0.7	0.3	0.7	0.4	0.7	0.7	0.6	0.8	1.1	1.1	0.8	0.9
	p	5.0	4.7	0.7	0.3	1.2	0.7	1.0	0.8	1.3	1.4	1.5	2.2	1.6	1.5
442	i	.	0.1	0.2	0.1	0.5	0.4	0.6	0.3	1.2	1.5	1.4	2.0	1.0	3.3
	p	.	0.1	0.2	0.1	0.5	0.5	0.7	0.5	1.4	2.2	1.6	2.8	1.2	4.3
443	i	2.2	2.4	4.4	4.4	0.2	0.0	0.0	0.0	0.1
	p	2.5	2.5	4.6	5.1	0.2	0.1	0.0	0.0	0.1
444	i	0.7	1.4	0.8	0.5	0.4	0.7	0.7	0.7	0.4	0.5	0.4	0.5	0.6	0.4
	p	1.0	1.6	0.9	0.8	0.5	0.8	1.0	0.8	0.5	0.6	0.9	0.9	1.2	0.9
445	i	0.0	0.1	0.1	0.2	0.8	0.2	1.2	0.5	1.0	0.7
	p	0.0	0.1	0.3	0.4	1.1	0.7	2.4	2.0	3.5	2.3
446	i	0.2	0.0	0.1	0.2	0.8	0.4	1.8	0.8	1.0	0.8	1.4	1.3	0.5	0.2
	p	0.2	0.0	0.2	0.4	0.9	0.6	2.0	0.9	1.2	1.0	1.6	1.5	0.5	0.5
447	i	0.4	1.0	0.9	1.2	1.0	1.3	1.4	2.7	1.5	1.7	1.1	1.0	1.1	0.4
	p	0.4	1.4	0.9	1.3	1.2	1.6	1.7	3.1	1.7	1.9	1.6	1.8	1.8	0.7
448	i	.	.	0.0	.	0.2	0.0	0.1	0.1	0.1	0.0	0.4	0.0	0.6	0.2
	p	.	.	0.1	.	0.2	0.1	0.1	0.2	0.2	0.0	0.3	0.2	0.6	0.3
449	i	0.7	0.3	0.2	0.2	0.0	0.0	0.1	0.1	0.2	0.4	.	0.1	.	.
	p	0.7	0.3	0.2	0.2	0.0	0.0	0.1	0.2	0.2	0.4	.	0.1	.	.
450	i	0.4	0.3	0.8	1.3	0.5	1.7	0.6	1.5	0.2	0.6	0.1	0.7	0.1	0.3
	p	0.4	0.5	0.9	1.8	0.6	2.1	0.8	2.0	0.4	0.9	0.1	0.8	0.1	0.5
451	i	0.0
	p	0.2
452	i	6.1	8.7	4.2	3.3	1.1	1.5	0.7	0.7	0.3	0.4	0.1	0.4	0.5	0.3
	p	6.4	8.8	4.1	3.3	1.2	1.5	0.7	0.9	0.3	0.4	0.1	0.4	0.5	0.3
453	i	0.2	0.0	0.2	0.1	0.1	0.1	.	.	.	0.0
	p	0.2	0.1	0.3	0.2	0.2	0.1	.	0.0	0.0	0.6
454	i	1.1	1.7	0.7	0.6	1.1	0.9	1.5	1.3	1.1	0.7	1.0	0.4	0.2	0.6
	p	1.3	2.0	0.8	0.8	1.5	1.3	2.5	2.1	1.5	1.2	1.6	1.1	0.7	1.3
455	i	5.6	6.4	1.7	2.3	0.7	1.7	0.5	0.8	0.4	0.3	0.3	0.1	0.3	0.9
	p	12.6	11.8	2.8	3.7	1.4	2.8	0.8	1.7	0.7	0.9	0.7	0.3	0.5	1.5

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
456	contact dermatitis	i	18681	1.3	0.8	1.8	0.9	1.0	2.0	1.5	0.9	0.7	0.9
		p	27597	1.9	1.3	2.5	1.3	1.2	2.8	2.2	1.6	1.1	1.7
457	eczema NEC	i	109004	7.4	6.5	8.3	10.5	6.8	7.8	7.4	7.1	6.6	7.3
		p	232076	15.8	13.6	17.9	17.0	10.8	14.7	14.7	17.6	20.7	23.6
458	diaper rash	i	7217	0.5	0.5	0.5	8.1	.	.	0.0	0.0	.	0.0
		p	11035	0.8	0.7	0.8	12.1	0.0	0.0	0.0	0.0	.	0.0
459	pityriasis rosea	i	5626	0.4	0.3	0.4	0.1	0.5	0.7	0.3	0.4	0.1	0.1
		p	7376	0.5	0.5	0.5	0.1	0.5	0.9	0.5	0.5	0.2	0.2
460	psoriasis w/wo arthropathy	i	6651	0.5	0.5	0.4	0.2	0.3	0.4	0.6	0.5	0.6	0.5
		p	27904	1.9	2.0	1.8	0.3	0.4	1.2	1.7	3.0	4.3	2.9
461	pompholyx/dis. sweat glands	i	5932	0.4	0.3	0.5	0.1	0.2	0.8	0.5	0.2	0.1	0.0
		p	7289	0.5	0.3	0.7	0.1	0.3	1.0	0.7	0.3	0.1	0.0
462	sebaceous cyst	i	35350	2.4	2.6	2.2	0.2	0.7	3.1	3.0	2.7	2.1	2.1
		p	41375	2.8	3.1	2.6	0.2	0.8	3.5	3.6	3.2	2.5	2.7
463	ingrown toenail/dis. of nail NEC	i	13981	1.0	1.1	0.8	0.3	1.3	1.9	0.6	0.7	0.7	0.9
		p	18523	1.3	1.4	1.2	0.4	1.5	2.5	0.9	1.0	0.9	1.4
464	alopecia/folliculitis/dis. of hair NEC	i	13317	0.9	0.9	0.9	0.7	0.6	0.9	1.3	0.9	0.1	0.5
		p	18378	1.3	1.3	1.2	0.7	0.9	1.3	1.7	1.3	0.3	1.1
465	acne vulgaris	i	24230	1.7	1.5	1.8	0.3	1.6	5.7	1.3	0.2	0.1	0.1
		p	83884	5.7	5.6	5.9	0.3	4.3	22.2	3.9	0.7	0.1	0.3
466	chron. ulcer skin, incl. varicose	i	5410	0.4	0.3	0.4	0.2	0.0	0.3	0.2	0.3	0.7	2.2
		p	16996	1.2	0.9	1.4	0.4	0.1	0.3	0.3	1.0	3.1	10.1
467	urticaria	i	17634	1.2	1.1	1.3	3.1	1.8	1.5	0.9	0.7	0.7	1.0
		p	20868	1.4	1.3	1.5	3.3	1.9	1.7	1.2	0.9	0.9	1.3
468	seborrheic warts	i	3466	0.2	0.1	0.4	0.1	0.1	0.1	0.1	0.4	0.7	0.8
		p	4270	0.3	0.2	0.4	0.1	0.1	0.1	0.2	0.5	0.8	1.0
469	acne rosacea	i	3768	0.3	0.2	0.3	0.3	0.2	0.1	0.2	0.4	0.4	0.4
		p	5889	0.4	0.3	0.5	0.3	0.3	0.2	0.3	0.8	0.5	0.5
470	dis. skin/subcut. tissue NEC	i	21089	1.4	1.2	1.7	1.3	1.0	1.8	1.3	1.4	1.2	2.6
		p	36547	2.5	2.0	3.0	2.0	1.2	2.7	2.5	2.7	2.5	4.6

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
456	i	1.1	0.7	0.8	1.2	0.6	3.4	1.0	2.1	0.7	1.1	0.2	1.0	0.6	1.1
	p	1.5	1.1	1.1	1.3	1.2	4.4	1.3	3.0	1.5	1.6	0.5	1.6	1.4	1.9
457	i	10.1	10.9	5.6	8.0	5.5	10.1	6.0	8.8	7.0	7.1	7.0	6.3	9.2	6.3
	p	17.0	17.0	9.7	11.9	9.9	19.7	12.3	17.1	15.6	19.5	21.3	20.3	27.4	21.6
458	i	7.4	8.8	0.0	.	0.0	.	.	.	0.0
	p	11.1	13.3	0.1	.	0.1	.	.	0.0	.	0.0	.	.	.	0.0
459	i	0.0	0.2	0.5	0.4	0.6	0.9	0.2	0.5	0.5	0.3	0.1	.	.	0.2
	p	0.1	0.2	0.7	0.4	0.7	1.2	0.3	0.7	0.7	0.3	0.1	0.2	.	0.3
460	i	0.4	0.0	0.3	0.2	0.4	0.3	0.7	0.4	0.5	0.5	0.2	0.9	1.1	0.3
	p	0.6	0.0	0.6	0.3	1.1	1.3	2.0	1.5	3.1	3.0	4.6	4.1	2.7	3.0
461	i	0.2	0.1	0.1	0.4	0.4	1.3	0.4	0.6	0.2	0.3	.	0.2	0.0	.
	p	0.2	0.1	0.1	0.4	0.4	1.5	0.5	0.9	0.2	0.4	.	0.2	0.0	.
462	i	0.1	0.3	0.8	0.5	3.8	2.5	3.0	3.1	3.0	2.4	2.6	1.7	2.7	1.7
	p	0.2	0.5	0.8	0.7	4.2	2.9	3.5	3.6	3.6	2.8	3.0	2.0	3.5	2.4
463	i	0.4	0.2	1.4	1.2	2.7	1.1	0.7	0.5	0.5	0.9	0.4	0.9	0.4	1.2
	p	0.7	0.2	1.6	1.3	3.3	1.6	1.0	0.9	0.7	1.3	0.5	1.2	1.2	1.4
464	i	0.5	0.9	0.4	0.8	0.9	1.0	1.4	1.3	1.0	0.8	0.0	0.1	0.3	0.6
	p	0.5	0.9	0.8	1.0	1.3	1.3	1.8	1.6	1.6	1.1	0.3	0.4	0.4	1.5
465	i	0.5	0.1	1.3	1.9	5.8	5.6	0.8	1.8	0.2	0.2	.	0.1	0.2	0.0
	p	0.5	0.1	2.5	6.1	24.1	20.2	2.6	5.3	0.7	0.7	.	0.2	0.2	0.4
466	i	0.3	0.1	0.0	0.1	0.4	0.2	0.2	0.2	0.3	0.3	0.5	0.9	2.0	2.4
	p	0.4	0.4	0.1	0.1	0.4	0.3	0.2	0.4	1.3	0.7	3.1	3.2	6.2	12.2
467	i	4.1	2.0	2.3	1.3	1.0	1.9	0.5	1.4	0.7	0.7	0.5	0.8	1.9	0.5
	p	4.1	2.4	2.3	1.5	1.2	2.1	0.8	1.7	0.8	1.0	0.6	1.1	2.1	0.8
468	i	.	0.1	.	0.1	0.0	0.1	0.0	0.2	0.2	0.6	0.8	0.6	0.1	1.1
	p	.	0.1	0.1	0.1	0.0	0.1	0.1	0.2	0.3	0.7	0.8	0.8	0.1	1.4
469	i	0.4	0.2	0.1	0.3	0.0	0.1	0.2	0.3	0.5	0.3	0.4	0.4	0.1	0.6
	p	0.4	0.2	0.2	0.4	0.1	0.3	0.2	0.4	0.6	0.9	0.5	0.5	0.3	0.7
470	i	1.8	0.9	1.1	0.9	1.2	2.5	1.0	1.6	1.0	1.7	1.2	1.3	2.1	2.8
	p	2.3	1.7	1.5	1.0	1.8	3.7	1.9	3.1	2.0	3.3	2.5	2.4	3.8	5.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
471	excessive thirst	i	593	0.0	0.0	0.0	0.2	0.0	.	0.1	0.0	0.1	.
		p	694	0.0	0.0	0.1	0.2	0.0	.	0.1	0.0	0.1	.
472	excessive appetite, excl. boulimia	i	300	0.0	0.0	0.0	0.2	0.0	0.0	0.0	.	.	.
		p	1198	0.1	0.0	0.1	0.2	0.0	0.0	0.2	0.0	.	.
473	loss of appetite, excl. anorexia nerv.	i	1027	0.1	0.1	0.1	0.3	0.0	0.1	0.0	0.1	0.0	0.1
		p	2148	0.1	0.1	0.2	0.5	0.1	0.1	0.0	0.1	0.4	0.5
474	feeding problem infant, excl. P11	i	1948	0.1	0.2	0.1	1.7	0.2
		p	3227	0.2	0.3	0.2	2.6	0.4
475	feeding problem adult, excl. T06	i	1779	0.1	0.1	0.1	.	.	0.2	0.1	0.0	.	0.3
		p	5049	0.3	0.2	0.5	.	.	0.3	0.2	0.4	0.6	1.4
476	feeding problem NEC	i	428	0.0	0.0	0.0	0.1	0.0	0.0	0.0	0.0	.	.
		p	1429	0.1	0.1	0.1	0.2	0.0	0.1	0.1	0.1	0.2	0.0
477	weight gain, excl. T82	i	1050	0.1	0.0	0.1	.	.	0.1	0.1	0.1	.	.
		p	2071	0.1	0.0	0.2	.	.	0.3	0.1	0.3	0.1	0.0
478	weight loss, excl. T06	i	1699	0.1	0.1	0.1	.	.	0.2	0.1	0.0	0.1	0.4
		p	3840	0.3	0.2	0.3	.	0.1	0.3	0.3	0.2	0.5	0.9
479	weight problem NEC	i	2037	0.1	0.1	0.2	.	0.0	0.2	0.2	0.2	0.1	.
		p	5822	0.4	0.1	0.7	0.1	0.0	0.4	0.5	0.6	0.6	0.2
480	failure to thrive	i	1965	0.1	0.2	0.1	0.7	0.6	0.1
		p	4270	0.3	0.4	0.2	1.4	1.3	0.1
481	dehydration	i	347	0.0	0.0	0.0	0.0	.	.	0.0	0.0	.	0.3
		p	975	0.1	0.1	0.1	0.1	.	0.0	0.0	0.0	.	0.8
482	thyroid lump/mass	i	455	0.0	0.0	0.1	.	.	.	0.1	0.0	0.1	.
		p	455	0.0	0.0	0.1	.	.	.	0.1	0.0	0.1	.
483	fear of endocr. metab. nutr. dis.	i	1188	0.1	0.0	0.1	0.0	0.0	0.1	0.1	0.1	0.0	0.1
		p	1699	0.1	0.0	0.2	0.0	0.0	0.1	0.1	0.2	0.3	0.1
484	disability/impairment end. gl.	i	0
		p	52	0.0	.	0.0	0.0	.	.
485	sympt./complt. end./met./nutr. NEC	i	2449	0.2	0.1	0.2	0.1	0.6	0.3	0.0	0.0	0.1	.
		p	3811	0.3	0.2	0.3	0.2	1.0	0.5	0.1	0.0	0.1	.
486	infectious dis. endocrine glands	i	0
		p	10	0.0	.	0.0	0.0
487	malign. neopl. thyroid	i	159	0.0	.	0.0	0.0	0.0	.
		p	837	0.1	0.0	0.1	.	.	0.0	0.0	0.1	0.1	0.3
488	benign. neopl. thyroid	i	9	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	51	0.0	.	0.0	.	.	.	0.0	0.0	.	.
489	unspecified neopl. endocr. NEC	i	331	0.0	0.0	0.0	0.1	.	.
		p	916	0.1	0.0	0.1	.	.	.	0.1	0.2	0.1	0.0
490	injury thyroid	i	0
		p	93	0.0	0.0	0.0	0.0	.	0.1
491	thyroglossal duct(cyst)	i	238	0.0	.	0.0	.	.	0.0	0.0	0.1	.	.
		p	416	0.0	0.0	0.1	.	.	0.0	0.0	0.1	.	0.0
492	cong. anom. endocr. NEC	i	53	0.0	0.0	.	0.1
		p	445	0.0	0.0	0.0	0.1	0.1	0.0	0.0	.	.	.
493	goiter/module w/o thyrotox	i	784	0.1	0.0	0.1	.	0.0	0.0	0.1	0.1	0.1	.
		p	3350	0.2	0.1	0.4	.	0.0	0.0	0.1	0.4	0.8	0.4
494	obesity/overweight	i	20265	1.4	0.6	2.1	.	0.7	1.5	1.9	1.7	1.0	0.6
		p	84663	5.8	2.7	8.8	0.1	1.2	3.9	6.4	10.5	7.9	4.9
495	hyperthyroidism/thyrotoxicosis	i	2210	0.2	0.0	0.3	.	0.0	0.1	0.2	0.2	0.1	0.1
		p	18829	1.3	0.3	2.2	0.2	0.0	0.3	1.0	1.8	3.2	5.8

T ENDOCRINE/METABOLIC/NUTRITIONAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
471	i	0.1	0.2	.	0.1	.	.	0.0	0.1	0.0	.	0.2	0.1	.	.
	p	0.1	0.2	.	0.1	.	.	0.0	0.1	0.0	.	0.2	0.1	.	.
472	i	0.2	0.2	.	0.1	.	0.0	.	0.0
	p	0.2	0.2	0.0	0.1	.	0.0	0.0	0.3	.	0.1
473	i	0.5	0.2	0.0	.	.	0.2	.	0.0	0.1	0.1	.	0.0	0.3	0.0
	p	0.6	0.4	0.1	.	.	0.2	0.0	0.1	0.1	0.1	0.3	0.5	0.5	0.4
474	i	1.8	1.5	0.3	0.1
	p	2.8	2.3	0.5	0.2
475	i	0.2	0.1	0.2	0.1	0.1	0.0	.	.	0.4	0.3
	p	0.2	0.3	0.2	0.3	0.3	0.5	0.7	0.6	0.4	1.9
476	i	0.3	.	.	0.0	.	0.1	0.0	.	0.1
	p	0.5	.	.	0.1	.	0.2	0.1	0.1	0.1	0.1	0.0	0.3	.	0.0
477	i	0.3	0.0	0.1	0.1	0.1
	p	0.5	0.0	0.2	0.1	0.4	.	0.2	0.1	.
478	i	0.3	0.1	0.1	0.2	0.0	0.0	0.2	0.1	0.9	0.2
	p	.	.	0.1	.	0.3	0.3	0.2	0.4	0.1	0.3	0.5	0.5	1.2	0.8
479	i	.	.	.	0.0	0.1	0.3	0.1	0.4	0.1	0.3	.	0.1	.	.
	p	.	0.2	.	0.0	0.1	0.7	0.2	0.7	0.2	1.0	0.1	1.0	.	0.3
480	i	1.0	0.5	0.7	0.4	0.1	0.0
	p	1.8	1.0	1.6	1.0	0.2	0.1
481	i	0.1	0.0	0.0	0.1	.	.	.	0.1	0.4
	p	0.2	0.0	.	.	.	0.0	0.1	0.0	0.1	.	.	.	0.2	1.1
482	i	0.0	0.1	.	0.0	.	0.2	.	.
	p	0.0	0.1	.	0.0	.	0.2	.	.
483	i	.	0.0	0.0	.	0.0	0.1	0.0	0.2	0.1	0.2	0.1	0.0	.	0.2
	p	.	0.1	0.0	.	0.0	0.1	0.0	0.2	0.1	0.3	0.1	0.5	.	0.2
484	i
	p	0.0
485	i	0.2	0.1	0.4	0.9	0.3	0.3	.	0.1	.	0.0	0.1	0.0	.	.
	p	0.2	0.2	0.6	1.4	0.6	0.4	0.0	0.1	0.0	0.0	0.1	0.1	.	.
486	i
	p	0.0
487	i	0.1	.	0.1	.	.
	p	0.0	0.0	0.0	0.1	0.0	0.2	0.0	0.2	0.1	0.4
488	i	0.0
	p	0.0	.	0.0
489	i	0.1	0.1
	p	0.0	0.1	0.2	0.2	.	0.2	.	0.0
490	i
	p	0.0	0.2
491	i	0.0	.	0.0	.	0.1
	p	0.0	.	0.1	.	0.2	.	.	0.0	.
492	i	0.1
	p	0.1	0.1	.	0.2	0.1	.	0.0	0.1
493	i	.	.	.	0.1	.	0.0	.	0.1	0.0	0.1	0.1	0.2	.	.
	p	.	.	.	0.1	0.0	0.1	0.0	0.3	0.3	0.6	0.1	1.4	.	0.7
494	i	.	.	0.4	1.0	0.6	2.4	0.7	3.1	0.6	2.7	0.9	1.1	0.7	0.5
	p	0.0	0.3	0.5	1.9	1.5	6.4	3.0	9.8	5.2	15.6	3.3	11.4	2.4	6.2
495	i	.	.	0.0	.	0.1	0.2	0.0	0.4	.	0.3	.	0.2	.	0.2
	p	0.3	0.1	0.0	.	0.2	0.4	0.2	1.9	0.6	3.0	1.1	4.9	1.1	8.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
496	hypothyroidism/myxedema	i	1068	0.1	0.0	0.1	.	.	0.1	0.1	0.1	0.1	0.0
		p	21860	1.5	0.5	2.5	0.3	0.1	0.2	0.8	2.8	4.6	5.5
497	hypoglycemia	i	1925	0.1	0.1	0.2	.	0.0	0.0	0.1	0.2	0.3	0.7
		p	3489	0.2	0.2	0.3	.	0.0	0.1	0.2	0.3	0.4	1.2
498	renal glucosuria	i	100	0.0	0.0	0.0	.	.	0.0	0.0	.	0.0	.
		p	289	0.0	0.0	0.0	.	.	0.0	0.0	0.0	0.0	.
499	diabetes mellitus	i	13087	0.9	0.8	1.0	0.0	0.1	0.3	0.4	1.4	3.4	3.2
		p	175879	12.0	9.3	14.6	0.3	1.0	2.3	3.7	16.9	49.1	63.5
500	vitamin defic./nutr. dis. NEC	i	1690	0.1	0.1	0.1	0.0	0.1	0.1	0.1	0.1	0.3	0.6
		p	20239	1.4	0.9	1.8	0.5	0.3	0.2	0.8	1.7	4.9	6.7
501	gout	i	6605	0.5	0.6	0.3	.	.	0.1	0.2	0.8	1.6	1.9
		p	17617	1.2	1.7	0.7	.	.	0.1	0.6	2.0	3.9	5.8
502	lipid metabolism dis.	i	2201	0.2	0.2	0.1	.	0.1	0.0	0.2	0.3	0.2	.
		p	23130	1.6	1.8	1.3	.	0.1	0.2	1.3	3.9	4.0	1.3
503	dis. thyroid NEC	i	757	0.1	0.0	0.1	.	0.0	.	0.0	0.1	0.2	.
		p	5944	0.4	0.1	0.7	0.0	0.4	0.0	0.2	0.7	1.5	0.9
504	dis. endocr. gland NEC	i	1274	0.1	0.0	0.2	.	0.0	0.1	0.2	0.0	.	0.0
		p	7899	0.5	0.2	0.8	0.0	0.1	0.6	0.6	0.8	0.5	0.6
505	metabolic disease NEC	i	1486	0.1	0.1	0.1	0.4	0.0	0.1	0.1	0.0	0.2	0.2
		p	13078	0.9	0.6	1.1	1.3	0.4	0.3	0.5	1.0	2.5	3.5

T ENDOCRINE/METABOLIC/NUTRITIONAL

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
496	i	0.1	0.0	0.2	0.0	0.2	.	0.1	.	0.0
	p	0.0	0.6	.	0.2	0.1	0.3	0.1	1.4	0.8	4.7	3.2	5.6	1.8	7.4
497	i	.	.	.	0.0	.	0.1	0.1	0.1	0.2	0.1	0.2	0.5	0.2	0.9
	p	.	.	.	0.0	0.1	0.1	0.2	0.2	0.3	0.2	0.2	0.5	0.3	1.6
498	i	0.1	0.0	.	.	.	0.0	.	.	.
	p	0.1	0.0	0.0	.	0.0	0.1	.	.	.
499	i	.	0.0	0.0	0.1	0.2	0.4	0.5	0.4	1.1	1.6	3.1	3.7	3.7	2.9
	p	0.4	0.1	1.2	0.8	2.4	2.1	4.3	3.1	14.4	19.4	39.4	56.5	50.8	70.2
500	i	.	0.1	0.0	0.1	0.1	0.0	0.1	0.1	0.0	0.2	0.6	0.1	0.7	0.5
	p	0.5	0.5	0.1	0.5	0.3	0.1	0.6	1.0	1.2	2.2	3.0	6.3	5.7	7.2
501	i	0.0	0.1	0.3	0.1	1.2	0.5	2.8	0.7	2.6	1.6
	p	0.0	0.1	1.2	0.1	3.4	0.7	5.8	2.4	7.4	4.9
502	i	.	.	.	0.1	0.0	.	0.4	0.0	0.2	0.3	0.1	0.3	.	.
	p	.	.	0.0	0.1	0.1	0.2	1.8	0.7	4.5	3.3	3.8	4.2	1.8	1.0
503	i	.	.	.	0.1	.	.	.	0.1	0.1	0.1	0.3	0.2	.	.
	p	.	0.0	0.1	0.7	.	0.0	.	0.4	0.2	1.2	0.7	2.0	.	1.4
504	i	.	.	0.0	0.0	.	0.3	.	0.3	0.0	0.1	.	.	0.0	.
	p	0.0	0.0	0.1	0.1	0.4	0.9	0.2	1.0	0.3	1.3	0.2	0.7	0.3	0.7
505	i	0.6	0.2	0.0	0.0	.	0.1	0.1	0.1	0.0	0.0	0.3	0.1	.	0.4
	p	1.0	1.5	0.2	0.6	0.4	0.2	0.5	0.5	0.6	1.4	1.9	3.0	2.1	4.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
506	complaints urination	i	6117	0.4	0.5	0.3	0.3	0.4	0.3	0.3	0.5	0.9	0.9
		p	10600	0.7	0.8	0.6	0.4	0.6	0.5	0.4	0.8	2.1	2.2
507	incontinence (excl. P12)	i	4672	0.3	0.1	0.5	.	0.0	0.1	0.2	0.5	1.0	1.5
		p	54669	3.7	1.6	5.8	0.2	0.3	0.5	1.0	2.8	9.2	38.6
508	anuria/oliguria	i	44	0.0	0.0	0.0	.	.	.	0.0	.	.	0.0
		p	44	0.0	0.0	0.0	.	.	.	0.0	.	.	0.0
509	complaints of urine	i	2742	0.2	0.2	0.1	0.1	0.1	0.0	0.2	0.3	0.5	0.5
		p	4430	0.3	0.4	0.2	0.1	0.1	0.1	0.3	0.4	0.9	1.0
510	sympt./complt. bladder NEC	i	1043	0.1	0.0	0.1	0.0	0.1	0.0	0.0	0.1	0.0	0.2
		p	3399	0.2	0.2	0.3	0.0	0.1	0.1	0.1	0.2	0.8	1.1
511	sympt./complt. kidney	i	224	0.0	0.0	0.0	.	0.0	.	0.0	0.0	.	.
		p	620	0.0	0.0	0.1	.	0.0	0.0	0.0	0.0	0.2	.
512	fear of urinary diseases	i	460	0.0	0.0	0.0	.	.	0.0	0.0	0.1	.	.
		p	781	0.1	0.0	0.1	.	.	0.0	0.1	0.1	0.2	.
513	disability/impairment urinary syst.	i	12	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	32	0.0	0.0	0.0	.	.	.	0.0	.	.	.
514	sympt./complt. urin. syst. NEC	i	209	0.0	0.0	0.0	0.1	0.0	.	.	0.0	.	.
		p	1198	0.1	0.1	0.1	0.1	0.0	.	0.1	0.1	0.3	0.2
515	acute pyelonephritis/pyelitis	i	4246	0.3	0.1	0.5	.	0.1	0.4	0.2	0.4	0.5	0.6
		p	5237	0.4	0.2	0.5	.	0.1	0.5	0.3	0.4	0.8	0.8
516	cystitis/urinary inf. NEC	i	170350	11.6	3.7	19.3	8.6	6.7	9.8	10.2	12.9	19.7	25.1
		p	189740	12.9	4.3	21.3	9.7	7.3	10.3	10.9	14.6	24.3	29.0
517	non-specific urethritis	i	4846	0.3	0.5	0.1	0.1	0.1	0.5	0.4	0.4	0.2	0.3
		p	5654	0.4	0.6	0.2	0.1	0.1	0.6	0.5	0.4	0.3	0.3
518	malign. neoplasm kidney	i	108	0.0	0.0	0.0	.	.	.	0.0	.	0.1	.
		p	2592	0.2	0.1	0.2	0.1	0.0	0.0	0.1	0.3	0.7	0.5
519	malign. neoplasm bladder	i	643	0.0	0.1	0.0	0.3	0.5
		p	3987	0.3	0.5	0.1	.	.	.	0.0	0.4	1.3	1.8
520	malign. neoplasm urinary tract NEC	i	297	0.0	0.0	0.0	.	0.0	.	0.0	0.0	0.0	0.1
		p	835	0.1	0.1	0.1	.	0.0	.	0.0	0.1	0.2	0.4
521	benign. neoplasm urinary tract	i	186	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	0.1
		p	513	0.0	0.0	0.0	.	.	.	0.0	0.0	0.1	0.3
522	unspecified neopl. urin. tract	i	213	0.0	0.0	0.0	0.0	0.0	0.2
		p	1007	0.1	0.1	0.0	.	.	.	0.0	0.1	0.4	0.3
523	injury urinary tract	i	169	0.0	0.0	0.0	.	0.0	0.0	.	0.0	.	.
		p	188	0.0	0.0	0.0	.	0.0	0.0	.	0.0	.	.
524	cong. anom. urinary tract	i	97	0.0	0.0	0.0	0.0	0.0	.	0.0	0.0	.	.
		p	1232	0.1	0.1	0.1	0.2	0.2	0.1	0.1	0.1	0.0	.
525	glomerulonephritis/nephrosis	i	555	0.0	0.0	0.0	.	0.1	.	0.0	0.0	0.1	0.1
		p	3895	0.3	0.3	0.3	0.1	0.1	0.1	0.1	0.5	0.5	0.7
526	orthostatic albuminuria	i	126	0.0	0.0	0.0	.	.	0.0	.	0.0	.	.
		p	190	0.0	0.0	0.0	.	.	0.0	0.0	0.0	.	0.1
527	urin. calculus, all types/sites	i	8419	0.6	0.8	0.4	.	0.1	0.3	0.6	1.1	1.0	0.4
		p	16766	1.1	1.5	0.8	.	0.1	0.4	1.4	2.2	1.7	0.8
528	abnormal urine test/findings NOS	i	1295	0.1	0.1	0.1	.	0.1	0.0	0.1	0.1	0.0	0.5
		p	3436	0.2	0.2	0.3	0.1	0.2	0.2	0.2	0.2	0.2	0.9
529	dis. urinary system NEC	i	4921	0.3	0.2	0.4	0.1	0.1	0.1	0.2	0.5	0.6	1.6
		p	24304	1.7	1.4	1.9	0.5	0.3	0.6	0.7	2.2	4.2	9.4

U UROLOGY

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
506	i	0.2	0.4	0.4	0.5	0.2	0.3	0.2	0.4	0.8	0.1	1.9	0.1	1.7	0.5
	p	0.3	0.4	0.5	0.6	0.4	0.6	0.2	0.6	1.2	0.4	3.7	0.9	3.7	1.4
507	i	.	.	0.0	0.0	0.0	0.1	0.1	0.3	0.2	0.8	0.6	1.3	0.7	2.0
	p	0.1	0.2	0.2	0.5	0.3	0.8	0.6	1.5	1.0	4.6	4.4	12.9	23.0	46.9
508	i	0.0	0.1	.
	p	0.0	0.1	.
509	i	.	0.1	0.1	.	.	0.1	0.2	0.2	0.3	0.2	1.0	0.1	0.7	0.3
	p	.	0.1	0.1	0.0	0.0	0.1	0.3	0.2	0.6	0.2	1.6	0.3	1.2	0.8
510	i	.	0.1	0.0	0.1	.	0.0	0.0	0.1	0.1	0.1	.	0.0	.	0.4
	p	.	0.1	0.0	0.1	.	0.2	0.0	0.2	0.2	0.2	1.0	0.8	1.3	1.0
511	i	.	.	.	0.1	.	.	0.0	0.0	.	0.1
	p	.	.	.	0.1	.	0.1	0.0	0.1	0.0	0.1	0.2	0.1	.	.
512	i	0.0	0.0	0.0	0.0	0.1	0.0
	p	0.0	0.0	0.0	0.1	0.1	0.0	.	0.4	.	.
513	i	0.0
	p	0.0	0.0
514	i	0.2	.	.	0.0	0.1	0.0
	p	0.2	.	.	0.0	.	.	0.1	0.1	0.1	0.1	0.3	0.2	.	0.3
515	i	.	.	0.1	0.1	0.0	0.8	0.1	0.4	0.3	0.4	0.1	0.8	.	1.0
	p	.	.	0.1	0.1	0.0	0.9	0.2	0.4	0.3	0.4	0.4	1.1	0.2	1.0
516	i	4.5	12.8	2.4	11.1	1.3	18.6	2.6	18.2	4.9	20.7	8.4	28.5	13.2	31.4
	p	4.9	14.7	2.6	12.1	1.3	19.7	2.8	19.3	5.7	23.4	11.1	34.4	16.3	35.7
517	i	0.2	.	0.3	.	0.7	0.3	0.7	0.1	0.5	0.2	0.2	0.2	0.3	0.2
	p	0.2	.	0.2	.	0.8	0.4	0.8	0.1	0.6	0.3	0.3	0.2	0.5	0.2
518	i	0.0	0.2	.	.
	p	0.1	.	.	0.1	.	0.1	0.1	0.1	0.4	0.3	0.4	0.8	0.5	0.5
519	i	0.4	0.1	1.4	0.0
	p	0.0	.	0.7	0.1	2.6	0.3	4.0	0.6
520	i	.	.	.	0.1	.	.	.	0.0	.	0.0	.	0.1	0.2	.
	p	.	.	.	0.1	.	.	.	0.0	0.0	0.1	0.3	0.1	0.7	0.2
521	i	0.0	.	0.0	.	0.1	0.3	.
	p	0.0	0.1	0.0	0.1	0.2	0.6	0.1
522	i	0.0	0.0	0.0	.	.	0.3
	p	0.0	.	0.1	0.0	0.7	0.2	0.3	0.4
523	i	.	.	.	0.1	0.1	0.0	.	.	0.0
	p	.	.	.	0.1	0.1	0.0	.	.	0.0	0.0
524	i	0.1	.	.	0.0	.	.	0.0	.	0.0
	p	0.4	.	0.1	0.2	0.1	0.1	0.0	0.1	0.0	0.1	.	0.0	.	.
525	i	.	.	0.1	.	.	.	0.0	0.1	0.0	0.0	0.2	.	.	0.2
	p	0.1	.	0.1	0.1	0.1	0.0	0.1	0.2	0.6	0.5	0.8	0.3	0.4	0.9
526	i	0.1	0.0
	p	0.1	.	0.0	.	.	0.0	.	.	.	0.1
527	i	.	.	0.2	0.1	0.3	0.3	0.7	0.5	1.6	0.6	1.8	0.3	0.4	0.4
	p	.	.	0.2	0.1	0.5	0.4	1.7	1.1	2.9	1.5	3.0	0.8	1.0	0.7
528	i	.	.	.	0.2	0.0	0.0	0.1	0.1	0.1	0.1	.	0.1	0.5	0.5
	p	0.1	0.0	0.0	0.4	0.2	0.2	0.2	0.3	0.3	0.1	0.2	0.3	0.6	1.1
529	i	.	0.2	0.1	0.1	0.1	0.2	0.2	0.4	0.5	0.4	0.5	0.7	0.9	1.9
	p	0.5	0.5	0.4	0.3	0.4	0.9	0.5	1.0	2.2	2.2	3.8	4.5	11.4	8.4

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
530	question of pregnancy (excl. W02)	i	3410	0.2	.	0.5	.	.	0.4	0.5	0.0	.	.
		p	3968	0.3	.	0.5	.	.	0.5	0.6	0.0	.	.
531	fear of being pregnant	i	2044	0.1	.	0.3	.	.	0.5	0.1	0.1	.	.
		p	3052	0.2	.	0.4	.	.	0.6	0.2	0.1	.	.
532	ante-partum bleeding	i	1395	0.1	.	0.2	.	0.1	0.2	0.2	.	.	.
		p	1691	0.1	.	0.2	.	0.1	0.3	0.2	.	.	.
533	vaginal discharge in pregn.	i	1167	0.1	.	0.2	.	.	0.1	0.2	.	.	.
		p	1148	0.1	.	0.2	.	.	0.1	0.2	.	.	.
534	vomiting/nausea of pregn.	i	622	0.0	.	0.1	.	.	0.1	0.1	.	.	.
		p	1280	0.1	.	0.2	.	.	0.2	0.2	.	.	.
535	wanted pregnancy	i	3887	0.3	.	0.5	.	.	0.4	0.6	.	.	.
		p	7396	0.5	.	0.9	.	.	0.6	1.3	0.0	.	.
536	morning-after pill	i	2070	0.1	.	0.3	.	0.1	0.5	0.1	0.0	.	.
		p	3418	0.2	.	0.5	.	0.1	0.7	0.3	0.1	.	.
537	fam. plan./oral contraceptive	i	42773	2.9	.	5.7	.	0.2	9.9	3.6	0.3	.	.
		p	516121	35.2	.	68.7	.	1.1	98.7	53.1	7.2	.	.
538	fam. plan./IUD	i	6135	0.4	.	0.8	.	.	0.3	1.1	0.1	.	.
		p	22261	1.5	.	3.0	.	.	1.0	4.0	0.3	.	.
539	fam. plan./sterilization ♀	i	3328	0.2	.	0.4	.	.	0.1	0.6	0.0	.	.
		p	5457	0.4	.	0.7	.	.	0.2	1.1	0.1	.	.
540	fam. plan. NEC ♀	i	1889	0.1	.	0.2	.	0.1	0.2	0.3	0.0	.	.
		p	10310	0.7	.	1.4	.	0.1	0.7	1.5	0.4	.	.
541	sub-/infertility ♀	i	3749	0.3	.	0.5	.	.	0.4	0.6	.	.	.
		p	11592	0.8	.	1.5	.	.	0.8	2.0	0.1	.	.
542	heavy post-partum bleeding	i	450	0.0	.	0.1	.	.	0.1	0.1	.	.	.
		p	688	0.0	.	0.1	.	.	0.1	0.1	.	.	.
543	complt. post partum NEC	i	789	0.1	.	0.1	.	.	0.0	0.2	.	.	.
		p	1303	0.1	.	0.2	.	.	0.1	0.3	.	.	.
544	sympt./complt. lactation	i	872	0.1	.	0.1	.	.	0.1	0.1	.	.	.
		p	2151	0.1	.	0.3	.	.	0.1	0.4	.	.	.
545	sympt./complt. breast in pregn. NEC	i	745	0.1	.	0.1	.	.	0.1	0.1	.	.	.
		p	1163	0.1	.	0.2	.	.	0.1	0.2	.	.	.
546	starting delivery	i	387	0.0	.	0.1	.	.	0.0	0.1	.	.	.
		p	643	0.0	.	0.1	.	.	0.1	0.1	.	.	.
547	fear of complic of pregn./deliv.	i	901	0.1	.	0.1	.	.	0.1	0.1	0.0	.	.
		p	1495	0.1	.	0.2	.	.	0.1	0.3	0.0	.	.
548	disability/impairment Pr/D/Pu/Fp	i	0	0.0	.	0.0
		p	12	0.0	.	0.0	.	.	.	0.0	.	.	.
549	sympt./complt. Pr/D/Pu/Fp NEC	i	1410	0.1	.	0.2	.	.	0.2	0.2	.	.	.
		p	2489	0.2	.	0.3	.	.	0.2	0.4	0.0	.	.
550	infect. dis. urinary tr. pregn./puerp.	i	2871	0.2	.	0.4	.	.	0.3	0.5	.	.	.
		p	3200	0.2	.	0.4	.	.	0.3	0.5	.	.	.
551	infect. dis. pregn. NEC	i	2300	0.2	.	0.3	.	.	0.4	0.3	.	.	.
		p	2641	0.2	.	0.4	.	.	0.4	0.3	.	.	.
552	malign. neoplasm pregn.	i	0	0.0	.	0.0
		p	3	0.0	.	0.0	.	.	.	0.0	.	.	.
553	benign. neoplasm pregn.	i	56	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	56	0.0	.	0.0	.	.	.	0.0	.	.	.
554	injuries complicating pregn.	i	0	0.0	.	0.0
		p	0	0.0	.	0.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
555	cong. anom. of mother/complic. pregn.	i	144	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	376	0.0	.	0.0	.	.	.	0.1	.	.	.
556	non-obst. cond. complic. pregn. NEC	i	2181	0.1	.	0.3	.	.	0.2	0.4	.	.	.
		p	3266	0.2	.	0.4	.	.	0.2	0.6	.	.	.
557	pregnancy: confirmed	i	46480	3.2	.	6.2	.	0.0	5.1	7.3	0.1	.	.
		p	69407	4.7	.	9.3	.	0.1	7.2	11.0	0.1	.	.
558	unwanted pregnancy: confirmed	i	2388	0.2	.	0.3	.	.	0.3	0.3	0.0	.	.
		p	3003	0.2	.	0.4	.	.	0.4	0.4	0.0	.	.
559	ectopic pregnancy	i	911	0.1	.	0.1	.	.	0.1	0.1	.	.	.
		p	1284	0.1	.	0.2	.	.	0.1	0.2	.	.	.
560	toxemia/(pre)eclampsia	i	473	0.0	.	0.1	.	.	0.0	0.1	.	.	.
		p	1491	0.1	.	0.2	.	.	0.2	0.2	.	.	.
561	abortion, imminens/spontan.	i	5414	0.4	.	0.7	.	0.0	0.5	0.9	.	.	.
		p	7600	0.5	.	1.0	.	0.0	0.7	1.2	0.0	.	.
562	abortion, (il)legally induced	i	1259	0.1	.	0.2	.	0.1	0.2	0.2	.	.	.
		p	2399	0.2	.	0.3	.	0.1	0.4	0.3	.	.	.
563	pregnancy, high risk	i	4966	0.3	.	0.7	.	0.0	0.4	0.8	.	.	.
		p	12438	0.8	.	1.7	.	0.0	0.9	2.2	.	.	.
564	normal delivery live born(s)	i	9241	0.6	.	1.2	.	.	0.9	1.5	.	.	.
		p	0
565	normal delivery dead born(s)	i	128	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	0
566	complicated delivery live born	i	4027	0.3	.	0.5	.	.	0.4	0.6	0.0	.	.
		p	0
567	complicated delivery dead born	i	284	0.0	.	0.0	.	.	.	0.1	.	.	.
		p	0
568	normal puerperium	i	19759	1.3	.	2.6	.	0.1	1.2	3.5	.	.	.
		p	26363	1.8	.	3.5	.	0.1	1.8	4.6	.	.	.
569	mastitis puerperalis	i	2646	0.2	.	0.4	.	.	0.2	0.5	.	.	.
		p	2871	0.2	.	0.4	.	.	0.2	0.5	.	.	.
570	dis. of breast puerperium NEC	i	1539	0.1	.	0.2	.	.	0.2	0.2	.	.	.
		p	2056	0.1	.	0.3	.	.	0.2	0.3	.	.	.
571	complic. puerperium NEC	i	3191	0.2	.	0.4	.	.	0.3	0.5	.	.	.
		p	4323	0.3	.	0.6	.	.	0.3	0.8	.	.	.
572	diseases Pr/D/Pu NEC	i	2052	0.1	.	0.3	.	.	0.2	0.3	.	.	.
		p	3495	0.2	.	0.4	.	.	0.2	0.6	.	.	.

W PREGNANCY, CHILD BEARING, FAMILY PLANNING

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
555	i	0.1
	p	0.1
556	i	0.4	.	0.8
	p	0.3	.	1.3
557	i	10.2	.	14.8	.	0.1
	p	.	.	.	0.1	.	14.5	.	22.3	.	0.1
558	i	0.7	.	0.7	.	0.0
	p	0.9	.	0.8	.	0.0
559	i	0.2	.	0.3
	p	0.2	.	0.5
560	i	0.1	.	0.2
	p	0.3	.	0.5
561	i	.	.	.	0.0	.	1.0	.	1.8
	p	.	.	.	0.0	.	1.4	.	2.5
562	i	.	.	.	0.1	.	0.3	.	0.3
	p	.	.	.	0.1	.	0.9	.	0.5	.	0.0
563	i	.	.	.	0.0	.	0.9	.	1.7
	p	.	.	.	0.0	.	1.8	.	4.5
564	i	1.9	.	3.0
	p
565	i	0.1
	p
566	i	0.7	.	1.2	.	0.0
	p
567	i	0.1
	p
568	i	.	.	.	0.1	.	2.5	.	7.0
	p	.	.	.	0.1	.	3.6	.	9.3
569	i	.	.	.	0.0	.	0.3	.	1.0
	p	.	.	.	0.0	.	0.3	.	1.1
570	i	0.5	.	0.4
	p	0.5	.	0.7
571	i	0.6	.	1.1
	p	0.7	.	1.5
572	i	0.3	.	0.7
	p	0.5	.	1.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
573	genital pain ♀	i	214	0.0	.	0.0	.	.	0.0	0.0	.	.	.
		p	327	0.0	.	0.0	.	.	0.0	0.1	.	.	0.0
574	menstruation absent/scanty	i	5929	0.4	.	0.8	.	0.0	1.2	0.6	0.0	.	.
		p	9607	0.7	.	1.3	.	0.1	1.9	1.0	0.1	.	.
575	menstruation excessive	i	14227	1.0	.	1.9	.	0.2	1.2	1.6	1.1	.	.
		p	23868	1.6	.	3.2	.	0.3	1.8	2.8	1.9	.	.
576	menstrual pain	i	12100	0.8	.	1.6	.	0.7	2.2	1.0	0.1	.	.
		p	27158	1.9	.	3.6	.	1.2	4.8	2.3	0.6	.	.
577	sympt./complt. menstruation NEC	i	15493	1.1	.	2.1	.	0.1	2.3	1.7	0.5	.	.
		p	23381	1.6	.	3.1	.	0.2	3.2	2.6	1.0	.	.
578	premenstrual symptoms	i	1938	0.1	.	0.3	.	.	0.2	0.2	0.1	.	.
		p	3837	0.3	.	0.5	.	0.0	0.3	0.5	0.2	.	.
579	sympt./complt. menopausal	i	14721	1.0	.	2.0	.	.	0.5	4.0	0.2	0.0	
		p	45565	3.1	.	6.1	.	.	1.1	12.8	2.1	0.2	
580	postmenopausal bleeding	i	1016	0.1	.	0.1	.	.	0.0	0.3	0.1	0.2	
		p	1767	0.1	.	0.2	.	.	0.0	0.4	0.3	0.4	
581	painful intercourse ♀	i	2523	0.2	.	0.3	.	.	0.5	0.2	0.1	0.0	
		p	3940	0.3	.	0.5	.	.	0.7	0.3	0.2	0.0	
582	postcoital bleeding	i	1238	0.1	.	0.2	.	.	0.1	0.2	0.0	.	
		p	1284	0.1	.	0.2	.	.	0.1	0.2	0.0	.	
583	vaginal discharge	i	11699	0.8	.	1.6	0.1	0.1	1.4	1.3	0.5	0.3	0.1
		p	15507	1.1	.	2.1	0.1	0.1	2.3	1.6	0.6	0.3	0.2
584	sympt./complt. vagina NEC	i	3417	0.2	.	0.4	0.2	0.2	0.3	0.3	0.3	0.1	0.1
		p	6285	0.4	.	0.8	0.3	0.2	0.4	0.4	0.6	0.4	0.4
585	sympt./complt. vulva	i	2782	0.2	.	0.4	0.0	0.1	0.4	0.1	0.2	0.1	0.0
		p	3918	0.3	.	0.5	0.0	0.1	0.5	0.2	0.3	0.4	0.2
586	sympt./complt. pelvis ♀	i	144	0.0	.	0.0	0.0	.	0.0	0.0	0.0	0.0	
		p	411	0.0	.	0.1	0.0	.	0.0	0.1	0.0	0.0	
587	lump/mass mamma ♀	i	2112	0.1	.	0.3	.	.	0.1	0.2	0.2	0.2	
		p	2812	0.2	.	0.4	.	.	0.2	0.3	0.2	0.2	
588	pain mamma ♀	i	4883	0.3	.	0.7	.	.	0.3	0.6	0.3	0.2	0.1
		p	6459	0.4	.	0.9	.	.	0.4	0.8	0.4	0.3	0.1
589	sympt./complt. mamma NEC ♀	i	14642	1.0	.	2.0	0.1	0.3	1.1	1.8	0.8	0.3	0.2
		p	19287	1.3	.	2.6	0.1	0.4	1.4	2.3	1.1	0.5	0.5
590	postponement of menstruation	i	14232	1.0	.	1.9	.	0.5	1.6	1.5	0.7	.	.
		p	17098	1.2	.	2.3	.	0.6	2.0	1.8	0.9	.	.
591	fear genital/mamma dis. ♀	i	3557	0.2	.	0.5	.	0.1	0.1	0.3	0.4	0.2	0.2
		p	6999	0.5	.	0.9	.	0.1	0.2	0.7	0.8	0.3	0.4
592	disab./impairm. gen. syst. ♀	i	10	0.0	.	0.0	.	.	0.0	0.0	.	.	.
		p	10	0.0	.	0.0	.	.	0.0	0.0	.	.	.
593	sympt./complt. gen. syst. ♀	i	665	0.0	.	0.1	0.0	0.0	0.0	0.1	.	.	.
		p	1174	0.1	.	0.2	0.0	0.0	0.1	0.2	0.0	.	.
594	syphilis (genit. and NOS) ♀	i	42	0.0	.	0.0	.	.	0.0	0.0	.	.	.
		p	201	0.0	.	0.0	.	.	0.1	0.0	.	.	.
595	gonorrhoea (genit. and NOS) ♀	i	67	0.0	.	0.0	.	.	0.0	0.0	.	.	.
		p	453	0.0	.	0.1	.	.	0.1	0.1	.	.	.
596	venereal dis. NEC ♀	i	2061	0.1	.	0.3	.	.	0.3	0.2	0.0	.	.
		p	2863	0.2	.	0.4	.	.	0.5	0.3	0.0	.	.
597	urogenital candidiasis ♀	i	36813	2.5	.	4.9	0.5	0.2	4.4	4.3	1.4	0.4	0.8
		p	47156	3.2	.	6.3	0.9	0.2	5.3	5.4	1.8	1.0	1.3

X FEMALE GENITAL SYSTEM (INCLUDING BREAST)

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
573	i	0.0	.	0.1
	p	0.0	.	0.1	0.0
574	i	.	.	.	0.0	.	2.4	.	1.3	.	0.1
	p	.	.	.	0.1	.	3.8	.	2.0	.	0.2
575	i	.	.	.	0.4	.	2.5	.	3.2	.	2.3
	p	.	.	.	0.6	.	3.7	.	5.7	.	3.8
576	i	.	.	.	1.5	.	4.5	.	2.1	.	0.3
	p	.	.	.	2.5	.	9.8	.	4.8	.	1.2
577	i	.	.	.	0.2	.	4.7	.	3.5	.	1.1
	p	.	.	.	0.5	.	6.5	.	5.2	.	2.0
578	i	0.4	.	0.5	.	0.2
	p	.	.	.	0.0	.	0.6	.	1.1	.	0.4
579	i	1.1	.	8.0	.	0.4	.	0.0
	p	2.2	.	25.2	.	3.8	.	0.4
580	i	0.0	.	0.5	.	0.2	.	0.2
	p	0.0	.	0.8	.	0.5	.	0.6
581	i	1.1	.	0.4	.	0.2	.	.	.	0.1
	p	1.4	.	0.7	.	0.4	.	.	.	0.1
582	i	0.2	.	0.4	.	0.0
	p	0.2	.	0.4	.	0.0
583	i	.	0.2	.	0.2	.	2.9	.	2.6	.	0.9	.	0.5	.	0.2
	p	.	0.2	.	0.2	.	4.6	.	3.2	.	1.2	.	0.5	.	0.3
584	i	.	0.3	.	0.3	.	0.6	.	0.6	.	0.4	.	0.2	.	0.2
	p	.	0.6	.	0.3	.	0.8	.	0.9	.	1.2	.	0.7	.	0.6
585	i	.	0.0	.	0.2	.	0.9	.	0.3	.	0.5	.	0.1	.	0.0
	p	.	0.0	.	0.2	.	0.9	.	0.5	.	0.6	.	0.7	.	0.3
586	i	.	0.0	0.0	.	0.0	.	0.0	.	.
	p	.	0.0	.	.	.	0.0	.	0.1	.	0.0	.	0.0	.	0.1
587	i	0.3	.	0.5	.	0.3	.	0.3	.	.
	p	0.4	.	0.7	.	0.3	.	0.3	.	.
588	i	0.6	.	1.3	.	0.6	.	0.4	.	0.2
	p	0.8	.	1.7	.	0.7	.	0.6	.	0.2
589	i	.	0.2	.	0.6	.	2.2	.	3.7	.	1.6	.	0.5	.	0.4
	p	.	0.2	.	0.7	.	2.8	.	4.8	.	2.2	.	0.8	.	0.7
590	i	.	.	.	1.0	.	3.3	.	3.1	.	1.3
	p	.	.	.	1.1	.	4.0	.	3.7	.	1.7
591	i	.	.	.	0.1	.	0.3	.	0.6	.	0.9	.	0.2	.	0.3
	p	.	.	.	0.1	.	0.5	.	1.5	.	1.6	.	0.3	.	0.6
592	i	0.0
	p	0.0
593	i	.	0.1	.	0.0	.	0.0	.	0.2
	p	.	0.1	.	0.0	.	0.1	.	0.4
594	i	0.0	.	0.0
	p	0.1	.	0.0
595	i	0.0	.	0.0
	p	0.1	.	0.1
596	i	0.7	.	0.5	.	0.1
	p	1.0	.	0.6	.	0.1
597	i	.	1.1	.	0.3	.	8.9	.	8.8	.	2.7	.	0.6	.	1.2
	p	.	1.9	.	0.5	.	10.7	.	11.1	.	3.6	.	1.8	.	2.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
598	urogenital trichomoniasis ♀	i	3866	0.3	.	0.5	.	.	0.5	0.3	0.3	.	.
		p	4557	0.3	.	0.6	.	.	0.7	0.4	0.3	.	.
599	pelvic inflammatory disease	i	4426	0.3	.	0.6	.	.	0.5	0.5	0.2	0.1	.
		p	6729	0.5	.	0.9	.	0.0	0.7	0.9	0.2	0.2	0.2
600	carcinoma in situ cervix	i	238	0.0	.	0.0	.	.	.	0.0	0.0	.	0.0
		p	1150	0.1	.	0.2	.	.	.	0.2	0.0	0.1	0.0
601	cervix carcinoma NEC	i	192	0.0	.	0.0	.	.	.	0.0	0.0	.	0.1
		p	1834	0.1	.	0.2	.	.	.	0.1	0.3	0.4	0.2
602	mal. neopl. mamma ♀	i	2849	0.2	.	0.4	.	.	0.0	0.1	0.5	0.5	0.9
		p	23595	1.6	.	3.2	.	.	0.0	0.7	3.2	4.6	7.7
603	mal. neopl. genit. syst. ♀	i	988	0.1	.	0.1	.	.	.	0.0	0.2	0.4	0.0
		p	6137	0.4	.	0.8	.	.	.	0.1	0.9	1.6	1.5
604	fibroid/myoma (uterus/cervix)	i	3711	0.3	.	0.5	.	.	.	0.6	0.3	0.0	0.1
		p	10318	0.7	.	1.4	.	0.0	.	1.4	1.1	0.2	0.5
605	ben. neopl. mamma ♀	i	3074	0.2	.	0.4	.	0.1	0.3	0.3	0.1	.	.
		p	4713	0.3	.	0.6	.	0.1	0.5	0.5	0.2	0.1	0.1
606	ben. neopl. genit. syst. ♀	i	2465	0.2	.	0.3	.	0.0	0.2	0.4	0.1	0.0	.
		p	4379	0.3	.	0.6	.	0.0	0.2	0.6	0.2	0.4	.
607	unspec. neopl. genit. syst. ♀	i	973	0.1	.	0.1	.	.	0.0	0.1	0.1	0.1	.
		p	2076	0.1	.	0.3	.	.	0.1	0.2	0.3	0.2	.
608	injury genit. syst./mamma ♀	i	346	0.0	.	0.0	0.1	0.0	0.0	.	0.0	.	.
		p	945	0.1	.	0.1	0.1	0.0	0.1	0.1	0.0	.	.
609	cong. anom. genit. syst./mamma ♀	i	24	0.0	.	0.0	0.0
		p	562	0.0	.	0.1	0.1	0.0	0.1	0.0	0.0	.	.
610	vaginitis/vulvitis NOS	i	13668	0.9	.	1.8	0.2	0.5	1.5	1.4	0.7	0.2	0.4
		p	20678	1.4	.	2.7	0.2	0.5	1.9	2.0	1.2	1.0	1.4
611	cervicitis/cerv. dis. NEC	i	2877	0.2	.	0.4	.	.	0.2	0.4	0.1	0.1	.
		p	5145	0.4	.	0.7	.	.	0.4	0.7	0.2	0.1	.
612	abnormal PAP-smear	i	675	0.0	.	0.1	.	.	.	0.1	0.1	0.0	.
		p	4074	0.3	.	0.5	.	.	0.1	0.6	0.4	0.1	.
613	uterovaginal prolapse	i	7823	0.5	.	1.1	.	.	0.1	0.5	0.9	1.3	1.6
		p	32130	2.2	.	4.3	.	.	0.1	1.2	3.2	6.7	12.5
614	cystic breast (chronic)	i	4619	0.3	.	0.6	.	0.1	0.2	0.6	0.3	0.0	.
		p	6671	0.5	.	0.9	.	0.2	0.3	0.8	0.5	0.1	.
615	premenstrual tension syndrome	i	2698	0.2	.	0.4	.	.	0.2	0.4	0.1	.	.
		p	4379	0.3	.	0.6	.	.	0.2	0.7	0.1	.	.
616	gardnerella infection	i	2635	0.2	.	0.4	.	.	0.4	0.3	0.1	.	.
		p	4284	0.3	.	0.6	.	.	0.6	0.5	0.3	.	.
617	infection/abscess gl. Bartholini	i	1427	0.1	.	0.2	.	.	0.0	0.2	0.1	.	.
		p	1830	0.1	.	0.2	.	.	0.0	0.3	0.1	.	0.1
618	endometriosis	i	1349	0.1	.	0.2	.	.	0.1	0.2	0.0	.	.
		p	3466	0.2	.	0.5	.	.	0.2	0.5	0.2	0.1	.
619	atrophial vaginae/vulvae	i	2769	0.2	.	0.4	.	.	.	0.0	0.5	0.6	0.7
		p	7405	0.5	.	1.0	.	.	0.0	0.0	1.1	1.7	2.1
620	dis. genital syst./mamma NEC ♀	i	6047	0.4	.	0.8	.	0.3	0.6	0.7	0.1	0.2	.
		p	10556	0.7	.	1.4	0.1	0.4	0.7	1.3	0.4	0.3	0.7

X FEMALE GENITAL SYSTEM (INCLUDING BREAST)

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
598	i	1.1	.	0.7	.	0.6
	p	1.4	.	0.8	.	0.6
599	i	1.1	.	1.1	.	0.3	.	0.2	.	.
	p	.	.	.	0.0	.	1.4	.	1.7	.	0.5	.	0.3	.	0.3
600	i	0.1	.	0.0	.	.	.	0.1
	p	0.4	.	0.1	.	0.3	.	0.1
601	i	0.0	.	0.1	.	.	.	0.1
	p	0.1	.	0.6	.	0.7	.	0.4
602	i	0.0	.	0.3	.	1.1	.	0.9	.	1.8
	p	0.1	.	1.4	.	6.3	.	8.1	.	11.8
603	i	0.0	.	0.3	.	0.6	.	0.0
	p	0.3	.	1.8	.	2.9	.	2.2
604	i	1.2	.	0.6	.	0.1	.	0.1
	p	0.0	.	2.8	.	2.2	.	0.3	.	0.8
605	i	.	.	.	0.1	.	0.7	.	0.7	.	0.3
	p	.	.	.	0.1	.	1.1	.	1.1	.	0.4	.	0.2	.	0.2
606	i	.	.	.	0.1	.	0.3	.	0.8	.	0.1	.	0.0	.	.
	p	.	.	.	0.1	.	0.4	.	1.2	.	0.5	.	0.7	.	.
607	i	0.1	.	0.2	.	0.3	.	0.1	.	.
	p	0.2	.	0.3	.	0.5	.	0.3	.	.
608	i	.	0.2	.	0.0	.	0.1	.	.	.	0.1
	p	.	0.2	.	0.0	.	0.3	.	0.2	.	0.1
609	i	.	0.0
	p	.	0.2	.	0.0	.	0.2	.	0.1	.	0.0
610	i	.	0.3	.	1.0	.	3.1	.	2.8	.	1.3	.	0.4	.	0.5
	p	.	0.3	.	1.0	.	3.8	.	4.0	.	2.3	.	1.7	.	2.1
611	i	0.5	.	0.8	.	0.2	.	0.1	.	.
	p	0.8	.	1.5	.	0.4	.	0.2	.	.
612	i	0.2	.	0.2	.	0.1	.	.
	p	0.2	.	1.2	.	0.8	.	0.1	.	.
613	i	0.1	.	1.1	.	1.8	.	2.2	.	2.5
	p	0.1	.	2.5	.	6.4	.	11.9	.	19.1
614	i	.	.	.	0.1	.	0.4	.	1.2	.	0.7	.	0.0	.	.
	p	.	.	.	0.2	.	0.5	.	1.7	.	1.1	.	0.2	.	.
615	i	0.3	.	0.9	.	0.1
	p	0.5	.	1.5	.	0.2
616	i	0.8	.	0.6	.	0.2
	p	1.1	.	0.9	.	0.5
617	i	0.1	.	0.4	.	0.3
	p	0.1	.	0.5	.	0.3	.	.	.	0.2
618	i	0.1	.	0.5	.	0.1
	p	0.4	.	1.1	.	0.3	.	0.1	.	.
619	i	0.0	.	1.0	.	1.1	.	1.0
	p	0.1	.	0.1	.	2.2	.	3.1	.	3.2
620	i	.	.	.	0.7	.	1.1	.	1.5	.	0.3	.	0.4	.	.
	p	.	0.2	.	0.8	.	1.5	.	2.6	.	0.7	.	0.6	.	1.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
621	sympt./complt. gen. syst. ♂	i	3550	0.2	0.5	.	0.4	0.3	0.2	0.2	0.3	0.1	0.1
		p	4781	0.3	0.7	.	0.4	0.4	0.2	0.3	0.4	0.2	0.3
622	discharge penis/urethra	i	244	0.0	0.0	.	.	.	0.0	.	.	0.1	.
		p	244	0.0	0.0	.	.	.	0.0	.	.	0.1	.
623	sympt./complt. prostate	i	494	0.0	0.1	.	.	.	0.0	0.0	0.1	0.3	.
		p	1242	0.1	0.2	.	.	.	0.0	0.2	0.2	0.2	0.7
624	sympt./complt. potency	i	961	0.1	0.1	.	.	0.0	0.0	0.2	0.0	.	.
		p	2182	0.1	0.3	.	.	0.0	0.1	0.4	0.5	.	.
625	sympt./complt. sexual dysf. ♂	i	498	0.0	0.1	.	.	.	0.1	0.0	0.1	.	.
		p	1000	0.1	0.1	.	.	0.0	0.1	0.1	0.2	0.0	.
626	sub-/infertility ♂	i	930	0.1	0.1	.	.	.	0.2
		p	2208	0.2	0.3	.	.	0.0	0.4
627	fam. plan. ♂	i	4979	0.3	0.7	.	.	0.1	1.0	0.1	.	.	.
		p	5968	0.4	0.8	.	.	0.1	1.1	0.1	.	.	.
628	sympt./complt. mamma ♂	i	905	0.1	0.1	.	.	0.2	0.2	.	0.0	0.1	.
		p	921	0.1	0.1	.	.	0.2	0.2	.	0.0	0.1	.
629	fear dis. gen. syst./mamma ♂	i	1244	0.1	0.2	.	.	0.1	0.1	0.1	0.1	0.2	.
		p	1753	0.1	0.2	.	.	0.2	0.2	0.1	0.2	0.2	.
630	disab./impairm. gen. syst./mamma ♂	i	0
		p	69	0.0	0.0	0.1
631	sympt./complt. gen. syst./mamma NEC ♂	i	742	0.1	0.1	.	.	0.0	0.1	0.1	.	.	.
		p	832	0.1	0.1	.	.	0.0	0.1	0.1	.	.	.
632	syphilis ♂	i	207	0.0	0.0	.	.	.	0.0
		p	310	0.0	0.0	.	.	.	0.0	0.0	0.0	.	.
633	gonorrhoea ♂	i	1731	0.1	0.2	.	.	0.5	0.1	0.0	.	0.1	.
		p	1787	0.1	0.2	.	.	0.4	0.1	0.0	0.0	0.0	0.1
634	venereal disease NEC ♂	i	653	0.0	0.1	.	.	0.0	0.1
		p	906	0.1	0.1	.	.	0.0	0.1	0.0	.	.	.
635	urogenital trichomoniasis ♂	i	413	0.0	0.1	.	.	0.0	0.0	0.0	0.0	0.0	.
		p	761	0.1	0.1	.	.	0.0	0.0	0.1	0.1	0.0	.
636	prostatitis/seminal vesiculitis	i	4236	0.3	0.6	.	.	0.1	0.3	0.5	0.4	0.7	.
		p	7468	0.5	1.0	.	.	0.2	0.5	1.0	0.9	1.1	.
637	orchitis/epididymitis	i	4865	0.3	0.7	.	.	0.1	0.3	0.6	0.3	0.3	0.2
		p	5752	0.4	0.8	.	.	0.2	0.3	0.6	0.4	0.4	0.4
638	infection gen. syst. NEC ♂	i	9160	0.6	1.3	.	1.4	0.5	0.5	0.7	0.6	0.3	0.4
		p	10942	0.7	1.5	.	1.7	0.5	0.7	0.9	0.7	0.3	0.7
639	malign. neopl. prostate	i	840	0.1	0.1	0.1	0.3	0.5	.
		p	6260	0.4	0.9	0.2	2.2	4.2	.
640	malign. neopl. testis	i	232	0.0	0.0	.	.	0.0	0.0	.	0.0	.	.
		p	918	0.1	0.1	.	.	0.0	0.0	0.1	0.0	0.3	0.0
641	malign. neopl. gen. syst. NEC ♂	i	13	0.0	0.0	0.0	.	.
		p	276	0.0	0.0	.	.	.	0.0	0.0	0.0	0.0	0.3
642	benign. neopl. gen. syst. ♂	i	196	0.0	0.0	.	.	0.0	0.0	.	.	0.1	.
		p	301	0.0	0.0	.	.	0.0	0.0	0.0	0.0	0.0	0.1
643	unspecif. neopl. gen. syst. ♂	i	380	0.0	0.1	.	.	0.0	0.0	0.0	0.1	0.1	.
		p	402	0.0	0.1	.	.	0.0	0.0	0.0	0.0	0.1	0.1
644	unspecif. neopl. mamma ♂	i	440	0.0	0.1	.	.	0.1	0.0	0.0	0.1	.	.
		p	475	0.0	0.1	.	.	0.1	0.0	0.0	0.0	0.1	.
645	injury gen. syst./mamma ♂	i	1169	0.1	0.2	.	0.1	0.2	0.1	0.0	0.1	0.1	.
		p	1205	0.1	0.2	.	0.1	0.2	0.1	0.0	0.1	0.1	.

Y MALE GENITAL SYSTEM (INCLUDING BREAST)

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
621	i	0.8	.	0.7	.	0.4	.	0.4	.	0.6	.	0.2	.	0.4	.
	p	0.8	.	0.9	.	0.5	.	0.6	.	0.8	.	0.4	.	0.9	.
622	i	0.1	.	.	.	0.2	.	.	.
	p	0.1	.	.	.	0.2	.	.	.
623	i	0.0	.	0.1	.	0.2	.	0.9	.
	p	0.0	.	0.3	.	0.4	.	1.9	.
624	i	0.1	.	0.1	.	0.4	.	0.0	.	.	.
	p	0.1	.	0.1	.	0.8	.	1.2	.	.	.
625	i	0.1	.	0.0	.	0.2	.	.	.
	p	0.1	.	0.2	.	0.2	.	0.5	.	0.0	.
626	i	0.4
	p	0.1	.	0.8
627	i	0.1	.	1.9	.	0.1
	p	0.2	.	2.2	.	0.3
628	i	.	.	0.3	.	0.3	.	.	.	0.1	.	0.2	.	.	.
	p	.	.	0.3	.	0.3	.	.	.	0.1	.	0.2	.	.	.
629	i	0.2	.	0.3	.	0.1	.	0.4	.	.	.
	p	0.3	.	0.3	.	0.2	.	0.5	.	.	.
630	i
	p	0.3	.
631	i	.	.	0.1	.	0.1	.	0.2
	p	.	.	0.1	.	0.1	.	0.3
632	i	0.1
	p	0.1	.	0.1	.	0.0	.	.	.
633	i	0.9	.	0.2	.	0.1	.	.	.	0.2	.
	p	0.8	.	0.2	.	0.1	.	0.0	.	0.2	.
634	i	0.1	.	0.2
	p	0.1	.	0.3	.	0.1
635	i	.	.	0.1	.	0.1	.	0.1	.	0.0	.	0.1	.	.	.
	p	.	.	0.1	.	0.1	.	0.1	.	0.2	.	0.1	.	.	.
636	i	0.2	.	0.6	.	1.1	.	1.0	.	2.0	.
	p	0.4	.	0.9	.	2.1	.	2.0	.	3.1	.
637	i	.	.	0.2	.	0.5	.	1.1	.	0.6	.	0.8	.	0.6	.
	p	.	.	0.3	.	0.6	.	1.2	.	0.8	.	1.0	.	1.1	.
638	i	2.8	.	0.9	.	1.0	.	1.4	.	1.3	.	0.7	.	1.1	.
	p	3.2	.	1.0	.	1.3	.	1.7	.	1.4	.	0.8	.	1.9	.
639	i	0.1	.	0.7	.	1.3	.
	p	0.5	.	4.9	.	11.9	.
640	i	0.1	.	0.0	.	.	.	0.1	.	.	.
	p	.	.	0.0	.	0.1	.	0.1	.	0.1	.	0.8	.	0.1	.
641	i	0.0	.	.	.
	p	0.0	.	0.0	.	0.0	.	0.9	.
642	i	0.0	.	0.0	0.3	.
	p	0.0	.	0.0	.	0.1	.	0.1	.	0.3	.
643	i	0.1	.	0.0	.	0.1	.	0.1	.	0.3	.
	p	0.1	.	0.0	.	0.1	.	0.2	.	0.3	.
644	i	.	.	0.3	.	0.0	.	0.0	.	0.0	.	0.2	.	.	.
	p	.	.	0.3	.	0.0	.	0.0	.	0.0	.	0.2	.	.	.
645	i	0.2	.	0.3	.	0.3	.	0.1	.	0.1	.	0.2	.	.	.
	p	0.2	.	0.3	.	0.3	.	0.1	.	0.1	.	0.2	.	.	.

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
646	phimosis/redund. prepuce	i	2689	0.2	0.4	.	0.7	0.6	0.1	0.1	0.1	.	0.1
		p	3901	0.3	0.5	.	1.5	0.8	0.1	0.1	0.1	0.0	0.1
647	hypospadias	i	0
		p	208	0.0	0.0	.	0.2	0.0
648	undesc. testicle/cryptorchism	i	1596	0.1	0.2	.	0.4	0.5	0.1	0.0	.	.	.
		p	2420	0.2	0.3	.	0.5	0.9	0.1	0.0	.	.	.
649	cong. anom. gen. syst./mamma NEC ♂	i	260	0.0	0.0	.	.	0.0
		p	370	0.0	0.1	.	.	0.0	0.0	0.0	0.0	.	.
650	benign prostatic hypertrophy	i	2080	0.1	0.3	0.0	0.2	0.8	0.9
		p	6691	0.5	0.9	0.0	0.5	2.6	3.1
651	hydrocele	i	1526	0.1	0.2	.	0.2	0.1	0.0	0.1	0.2	0.1	0.1
		p	1896	0.1	0.3	.	0.3	0.1	0.0	0.1	0.2	0.1	0.2
652	dis. gen. syst./mamma NEC ♂	i	4024	0.3	0.6	.	0.1	0.7	0.4	0.1	0.2	0.1	0.2
		p	5926	0.4	0.8	.	0.2	0.7	0.6	0.2	0.3	0.7	0.3

Y MALE GENITAL SYSTEM (INCLUDING BREAST)

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
646	i	1.5	.	1.2	.	0.2	.	0.2	.	0.1	.	.	.	0.2	.
	p	2.9	.	1.7	.	0.2	.	0.2	.	0.2	.	0.1	.	0.2	.
647	i
	p	0.3	.	0.1
648	i	0.8	.	1.0	.	0.2	.	0.0
	p	1.0	.	1.8	.	0.2	.	0.0
649	i	.	.	0.1
	p	.	.	0.1	.	0.0	.	0.1	.	0.1
650	i	0.0	.	0.3	.	1.9	.	2.6	.
	p	0.0	.	1.0	.	6.1	.	9.0	.
651	i	0.5	.	0.2	.	0.0	.	0.2	.	0.4	.	0.1	.	0.3	.
	p	0.7	.	0.2	.	0.1	.	0.2	.	0.4	.	0.3	.	0.4	.
652	i	0.2	.	1.3	.	0.9	.	0.3	.	0.4	.	0.3	.	0.5	.
	p	0.3	.	1.5	.	1.2	.	0.4	.	0.6	.	1.7	.	0.9	.

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
653	poverty/financial probl.	i	1699	0.1	0.2	0.1	.	.	0.1	0.1	0.3	0.0	0.3
		p	3934	0.3	0.3	0.2	.	.	0.1	0.4	0.5	0.1	0.3
654	problem food/water	i	39	0.0	.	0.0	.	.	.	0.0	.	.	.
		p	89	0.0	.	0.0	.	.	.	0.0	.	.	0.1
655	probl. housing/neighborh.	i	3645	0.2	0.2	0.3	0.0	.	0.3	0.2	0.3	0.6	0.5
		p	10716	0.7	0.5	1.0	0.2	0.1	0.8	0.7	0.7	1.6	2.1
656	probl. house moving	i	1580	0.1	0.1	0.1	0.1	0.1	0.2	0.0	0.1	0.2	0.5
		p	7980	0.5	0.4	0.7	0.1	0.1	0.6	0.5	0.5	0.9	1.8
657	fear of having a social probl.	i	191	0.0	0.0	0.0	.	.	.	0.0	0.0	.	.
		p	534	0.0	0.0	0.0	.	.	0.0	0.0	0.1	.	.
658	probl. working conditions	i	2935	0.2	0.3	0.1	.	.	0.3	0.2	0.4	.	.
		p	7956	0.5	0.7	0.4	.	.	0.6	0.9	0.7	0.3	.
659	probl. job/employment NEC	i	9011	0.6	0.7	0.5	.	.	0.7	1.0	0.9	.	.
		p	20422	1.4	1.6	1.1	.	.	1.5	2.2	2.2	0.2	.
660	probl. w. being unemployed	i	638	0.0	0.1	0.0	.	.	0.1	0.1	0.0	.	.
		p	2119	0.1	0.2	0.1	.	.	0.1	0.3	0.2	.	.
661	probl. w. education	i	3507	0.2	0.2	0.3	.	0.9	0.7	0.0	0.0	.	.
		p	7183	0.5	0.4	0.6	.	1.3	1.4	0.2	0.1	.	.
662	probl. soc. insurance/welfare	i	1778	0.1	0.2	0.1	0.0	.	0.0	0.1	0.3	0.1	0.1
		p	5379	0.4	0.5	0.2	0.0	0.0	0.1	0.5	0.9	0.2	0.1
663	probl. legal/police	i	130	0.0	0.0	0.0	.	.	0.0	0.0	.	.	.
		p	906	0.1	0.1	0.0	.	.	0.0	0.1	0.1	.	.
664	probl. health care syst./access	i	2705	0.2	0.2	0.2	0.3	0.0	0.1	0.1	0.3	0.2	0.5
		p	6088	0.4	0.3	0.5	0.5	0.0	0.2	0.4	0.6	0.9	0.6
665	probl. soc./cultural syst.	i	1463	0.1	0.2	0.0	0.1	0.1	0.2	0.1	0.1	0.0	.
		p	6113	0.4	0.5	0.3	0.1	0.1	0.6	0.7	0.4	0.0	.
666	relation problems partners	i	13535	0.9	0.7	1.2	.	0.0	0.6	1.5	1.1	0.9	0.7
		p	28576	1.9	1.3	2.6	.	0.1	1.0	2.9	3.1	1.9	1.8
667	problems w. behaving partner	i	950	0.1	0.0	0.1	.	.	.	0.1	0.1	0.1	.
		p	3125	0.2	0.1	0.3	.	.	0.1	0.3	0.4	0.2	0.1
668	probl. w. partner being ill	i	6600	0.4	0.4	0.5	.	.	0.1	0.4	0.9	1.2	1.2
		p	22276	1.5	1.1	1.9	.	0.0	0.1	1.1	2.9	4.5	4.8
669	probl. loss of partner	i	9818	0.7	0.5	0.9	.	0.1	0.2	0.5	1.3	1.3	2.3
		p	39445	2.7	1.4	3.9	.	0.1	0.4	1.7	4.5	8.5	10.1
670	relation probl. with child	i	9171	0.6	0.3	1.0	0.8	0.4	0.2	0.7	1.0	0.7	0.6
		p	16677	1.1	0.5	1.7	2.0	0.8	0.3	1.1	1.8	1.3	0.9
671	probl. behaving child	i	2012	0.1	0.1	0.2	.	0.1	.	0.2	0.2	0.1	.
		p	3415	0.2	0.1	0.4	.	0.2	0.0	0.4	0.3	0.1	.
672	probl. with child being ill	i	1563	0.1	0.1	0.1	0.0	.	.	0.1	0.1	0.4	0.2
		p	6218	0.4	0.2	0.6	0.0	0.1	0.1	0.6	0.7	0.7	0.5
673	loss or death of child	i	1640	0.1	0.0	0.2	0.0	.	0.1	0.1	0.1	0.1	0.4
		p	4355	0.3	0.2	0.4	0.1	.	0.2	0.4	0.4	0.4	0.8
674	relation probl. parent/other family	i	9089	0.6	0.4	0.9	0.3	0.2	0.7	0.6	0.9	0.7	0.3
		p	21014	1.4	0.9	1.9	0.7	0.9	1.6	1.5	1.8	1.7	1.3
675	probl. w. behaving parent/other family	i	586	0.0	0.0	0.0	0.1	0.1	.	0.0	0.1	0.1	0.1
		p	1013	0.1	0.0	0.1	0.1	0.1	.	0.1	0.1	0.1	0.1
676	probl. w. parent/other family being ill	i	3128	0.2	0.1	0.3	0.1	.	0.1	0.2	0.4	0.3	0.4
		p	9232	0.6	0.3	1.0	0.3	0.2	0.5	0.7	1.0	0.6	0.6
677	loss parent/family	i	5505	0.4	0.2	0.5	0.1	0.1	0.2	0.4	0.5	0.5	0.9
		p	14109	1.0	0.6	1.3	0.6	0.2	0.8	1.2	1.2	0.8	1.6

Z SOCIAL PROBLEMS

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
653	i	0.0	0.1	0.2	0.1	0.5	0.0	.	0.0	0.2	0.4
	p	0.0	0.1	0.5	0.3	0.7	0.3	.	0.2	0.2	0.4
654	i	0.0
	p	0.0	0.1
655	i	0.1	.	.	.	0.4	0.2	0.1	0.3	0.3	0.3	0.4	0.7	0.7	0.4
	p	0.1	0.2	0.1	0.0	0.6	1.0	0.4	1.0	0.6	0.8	0.9	2.1	1.6	2.4
656	i	0.2	.	0.2	.	0.2	0.1	0.0	0.0	0.1	0.0	0.3	0.1	0.6	0.4
	p	0.3	.	0.2	0.0	0.3	0.9	0.4	0.6	0.5	0.6	0.5	1.2	1.5	2.0
657	i	0.0	0.0	0.1	0.0
	p	0.0	0.1	0.0	0.0	0.1	0.0
658	i	0.2	0.4	0.4	0.1	0.5	0.2
	p	0.5	0.8	1.1	0.7	1.0	0.3	0.5	0.0	.	.
659	i	0.4	1.0	1.4	0.7	1.2	0.5
	p	1.2	1.8	2.7	1.6	2.7	1.6	0.2	0.3	.	.
660	i	0.1	0.0	0.2	0.0	0.0
	p	0.1	0.2	0.3	0.2	0.2	0.2
661	i	.	.	1.1	0.6	0.3	1.0	0.0	0.0	.	0.0
	p	.	.	1.5	1.2	0.7	2.1	0.2	0.2	0.1	0.1
662	i	.	0.0	.	.	0.0	.	0.2	0.1	0.3	0.3	0.2	0.0	0.0	0.1
	p	.	0.0	.	0.1	0.2	0.0	0.7	0.3	1.2	0.5	0.3	0.1	0.0	0.1
663	i	0.0	.	0.0	0.0
	p	0.1	0.0	0.2	0.0	0.1	0.0
664	i	0.4	0.2	.	0.0	0.2	0.0	0.1	0.1	0.1	0.5	0.5	0.0	0.2	0.6
	p	0.7	0.3	0.1	0.0	0.3	0.0	0.4	0.4	0.2	1.0	0.8	0.9	0.2	0.8
665	i	0.2	.	0.2	.	0.3	0.1	0.1	0.1	0.2	0.0	0.1	.	.	.
	p	0.3	0.0	0.2	0.0	0.7	0.4	0.6	0.7	0.7	0.2	0.1	.	.	.
666	i	.	.	.	0.1	0.1	1.1	1.0	2.1	1.0	1.2	0.8	0.9	1.0	0.6
	p	.	.	.	0.2	0.5	1.6	1.8	4.1	2.1	3.9	1.8	2.0	1.8	1.7
667	i	0.0	0.2	0.1	0.1	0.1	0.1	.	.
	p	0.1	0.1	0.7	0.3	0.5	0.2	0.2	0.1	0.1
668	i	0.1	0.1	0.3	0.4	0.7	1.0	1.1	1.3	1.8	0.8
	p	.	.	.	0.0	0.1	0.2	0.7	1.4	2.0	3.8	2.7	5.8	6.6	3.8
669	i	.	.	.	0.1	.	0.3	0.4	0.6	0.9	1.6	0.8	1.7	2.5	2.3
	p	.	.	0.0	0.3	0.2	0.6	1.1	2.3	2.4	6.6	4.4	11.7	7.6	11.5
670	i	0.6	1.0	0.2	0.6	0.2	0.1	0.1	1.2	0.6	1.5	0.1	1.1	0.1	0.8
	p	1.4	2.5	0.5	1.1	0.4	0.3	0.3	2.0	1.0	2.6	0.2	2.1	0.1	1.3
671	i	.	.	.	0.2	.	.	0.1	0.3	0.2	0.2	.	0.1	.	.
	p	.	.	.	0.3	.	0.1	0.1	0.6	0.2	0.4	0.1	0.1	.	.
672	i	0.0	0.0	0.2	0.1	0.2	0.5	0.3	0.2	0.2
	p	0.1	.	0.2	.	.	0.2	0.1	1.1	0.5	0.9	0.8	0.6	0.3	0.6
673	i	.	0.0	.	.	0.0	0.3	0.0	0.2	0.1	0.2	0.2	.	0.2	0.5
	p	0.0	0.1	.	.	0.1	0.3	0.2	0.6	0.2	0.5	0.3	0.5	0.6	0.8
674	i	0.3	0.3	0.2	0.2	0.4	1.0	0.5	0.8	0.4	1.3	0.1	1.2	0.0	0.5
	p	0.4	1.0	0.9	0.9	1.1	2.1	0.8	2.1	1.2	2.4	0.9	2.3	0.6	1.6
675	i	0.1	.	0.1	0.0	.	.	0.0	0.0	.	0.1	.	0.1	.	0.2
	p	0.1	.	0.1	0.1	.	.	0.1	0.1	.	0.1	.	0.1	.	0.2
676	i	.	0.2	.	.	0.0	0.2	0.1	0.3	0.1	0.8	0.4	0.3	0.2	0.6
	p	0.1	0.4	0.1	0.4	0.3	0.7	0.4	1.1	0.3	1.6	0.5	0.7	0.2	0.8
677	i	.	0.2	0.1	0.1	0.2	0.2	0.2	0.6	0.1	0.9	0.3	0.7	0.7	1.0
	p	0.9	0.2	0.1	0.3	0.8	0.9	0.8	1.5	0.4	2.1	0.6	1.0	0.9	1.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
totals and according to sex and age**

no.	episode		totals		sex		age						
			abs.	/1000 pers.	male	female	0-4	5-14	15-24	25-44	45-64	65-74	≥75
678	loss or death of friend	i	352	0.0	0.0	0.0	.	.	0.1	0.0	0.0	0.1	0.1
		p	1413	0.1	0.1	0.1	.	.	0.1	0.1	0.1	0.4	0.1
679	relation probl. w. friend	i	787	0.1	0.0	0.1	.	.	0.0	0.1	0.1	0.2	0.0
		p	1140	0.1	0.0	0.1	.	0.0	0.0	0.1	0.1	0.2	0.1
680	probl. assault/harmfull events	i	1950	0.1	0.1	0.2	0.1	0.2	0.2	0.1	0.1	0.1	.
		p	4394	0.3	0.1	0.5	0.1	0.2	0.5	0.4	0.1	0.3	0.1
681	probl. with being ill	i	525	0.0	0.0	0.0	.	.	.	0.1	0.1	0.0	0.0
		p	2720	0.2	0.1	0.2	.	.	0.0	0.1	0.3	1.0	0.1
682	social handicap	i	747	0.1	0.0	0.1	.	.	0.1	0.1	0.0	.	0.1
		p	2460	0.2	0.1	0.2	0.0	0.1	0.1	0.2	0.2	0.0	0.5
683	social isolation	i	2483	0.2	0.1	0.2	.	.	.	0.1	0.1	0.7	1.6
		p	14286	1.0	0.5	1.5	.	.	0.1	0.2	0.8	3.3	9.7
684	social problems	i	3683	0.3	0.2	0.3	0.1	0.1	0.2	0.2	0.4	0.2	0.8
		p	13608	0.9	0.6	1.2	0.7	0.2	0.7	1.0	1.0	1.5	2.2

Z SOCIAL PROBLEMS

no.		0-4 male	0-4 female	5-14 male	5-14 female	15-24 male	15-24 female	25-44 male	25-44 female	45-64 male	45-64 female	65-74 male	65-74 female	≥75 male	≥75 female
678	i	0.1	.	0.0	.	.	0.0	0.0	0.1	0.1	0.1
	p	0.1	0.1	0.1	0.1	0.0	0.2	0.0	0.6	0.1	0.1
679	i	0.0	0.0	0.1	0.1	0.0	0.4	.	.	0.1
	p	.	.	.	0.0	.	0.1	0.0	0.2	0.1	0.1	0.4	0.0	.	0.1
680	i	0.0	0.2	0.1	0.2	0.0	0.4	0.0	0.2	0.1	0.1	.	0.2	.	.
	p	0.0	0.2	0.2	0.2	0.1	1.0	0.1	0.7	0.1	0.2	.	0.5	0.4	.
681	i	0.0	0.1	0.1	.	0.0	.	.	0.1
	p	0.0	0.0	0.2	0.4	0.2	0.7	1.2	.	0.1
682	i	0.1	0.1	0.1	0.1	0.0	0.1
	p	0.0	.	.	0.1	0.1	0.1	0.2	0.3	0.2	0.2	0.1	0.0	0.1	0.7
683	i	0.0	0.1	0.1	0.0	0.4	1.0	2.1	1.4
	p	0.0	0.1	0.2	0.3	0.6	0.9	1.4	4.6	5.4	11.9
684	i	0.1	0.1	0.1	.	0.2	0.1	0.1	0.4	0.5	0.3	0.0	0.3	0.6	0.8
	p	0.4	0.9	0.4	0.0	0.4	1.0	0.6	1.4	0.8	1.1	0.3	2.5	2.4	2.2

**A.4 BASIC TABLE B: MORBIDITY IN GENERAL PRACTICE BY URBANIZATION
DEGREE, REGION, SEASON AND INSURANCE MODE**

(not standardized for age and sex)

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
001	pain: generalized/unspec.	i	0.3	0.2	0.3	0.3	0.4	0.3	0.2
		p	5.6	2.6	2.4	2.9	9.9	3.0	2.2
002	chills/fever	i	0.8	0.6	0.4	0.4	0.7	0.5	0.8
		p	1.0	0.8	0.6	0.6	0.8	0.7	1.0
003	general weak-/tiredness	i	1.6	2.6	2.8	2.6	2.2	2.4	2.3
		p	4.5	5.6	5.5	6.0	4.7	5.7	4.4
004	fainting (syncope)	i	0.9	0.9	0.9	0.5	1.0	0.7	1.1
		p	1.1	1.1	1.1	0.7	1.3	0.9	1.4
005	symptoms infant	i	0.1	0.2	0.2	0.3	0.2	0.2	0.1
		p	0.2	0.2	0.2	0.3	0.2	0.2	0.2
006	allergy NEC	i	3.2	2.5	2.4	3.3	2.6	3.2	2.0
		p	6.5	5.3	5.1	9.4	5.8	6.8	4.3
007	feeling/behaving old/senile	i	0.2	0.2	0.1	0.6	0.2	0.2	0.1
		p	1.2	1.1	1.4	4.3	1.9	1.6	1.3
008	fear of disease/death	i	0.4	0.4	0.6	0.5	0.5	0.5	0.6
		p	1.1	1.2	1.4	1.4	1.3	1.1	1.4
009	disability/impairment NOS	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.2	0.1	0.2	0.1	0.1	0.1	0.3
010	general sympt./complt. NEC	i	0.8	1.2	1.6	1.0	0.9	1.1	1.5
		p	1.9	2.0	2.6	1.9	2.1	1.8	2.8
011	tuberculosis, general, excl. lung	i	0.0	.	0.0	.	.	0.0	0.0
		p	0.1	0.0	0.1	0.1	0.1	0.1	0.1
012	measles	i	0.3	0.2	0.2	.	0.0	0.2	0.2
		p	0.3	0.2	0.2	.	0.1	0.2	0.2
013	chickenpox	i	0.6	0.6	0.5	0.7	0.7	0.5	0.8
		p	0.6	0.7	0.6	0.8	0.8	0.6	0.9
014	malaria	i	0.0	0.0	0.0	0.1	0.0	0.0	.
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
015	rubella	i	0.8	1.3	0.5	0.6	0.3	1.1	0.6
		p	0.9	1.3	0.6	0.6	0.3	1.2	0.6
016	infectious mononucleosis	i	0.6	0.3	0.4	0.6	0.7	0.4	0.4
		p	0.9	0.7	0.7	0.8	0.9	0.8	0.7
017	viral dis. with exanthems NEC	i	0.5	0.6	0.5	0.3	0.6	0.4	0.6
		p	0.5	0.6	0.6	0.4	0.7	0.5	0.7
018	viral dis. NOS/NEC	i	5.3	7.2	6.0	3.1	7.8	5.6	5.6
		p	5.5	7.7	6.4	3.3	8.1	6.0	6.0
019	infectious dis. NOS/NEC	i	0.1	0.3	0.2	0.1	0.2	0.2	0.2
		p	0.3	0.5	0.5	0.2	0.5	0.4	0.4
020	carcinomatosis (prim. site unkn)	i	0.1	0.0	0.2	0.1	0.1	0.1	0.0
		p	0.3	0.2	0.3	0.3	0.3	0.3	0.3
021	accident/injury NOS	i	0.9	0.6	0.8	0.7	0.9	0.7	0.8
		p	1.4	1.0	1.3	1.4	1.6	1.1	1.4
022	mult. trauma/internal injury	i	0.0	0.0	.	.	0.0	0.0	0.1
		p	0.0	0.0	0.0	.	0.0	0.0	0.1
023	late effects of trauma	i	0.2	0.1	0.2	0.3	0.1	0.1	0.3
		p	0.5	0.4	0.5	0.9	0.5	0.4	0.8
024	poisoning by/abuse of med. agents	i	0.1	0.1	0.1	0.1	0.0	0.1	0.1
		p	0.2	0.2	0.4	0.3	0.1	0.3	0.2
025	poisoning by benzodiazepines	i	.	0.0	.	.	0.0	0.0	.
		p	.	0.0	0.0	.	0.0	0.0	.

A GENERAL AND UNSPECIFIED

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
001	i	0.2	0.3	0.3	0.3	0.3	0.2
	p	2.6	5.5	3.2	2.9	4.4	2.3
002	i	0.4	0.6	0.8	0.6	0.5	0.6
	p	0.5	0.8	1.0	0.9	0.8	0.6
003	i	3.2	1.6	2.1	2.4	2.6	1.8
	p	6.0	4.9	4.3	6.0	5.8	4.4
004	i	0.7	0.6	1.3	0.7	1.0	0.5
	p	0.9	0.8	1.5	1.0	1.2	0.6
005	i	0.2	0.1	0.2	0.1	0.2	0.1
	p	0.2	0.2	0.2	0.2	0.2	0.1
006	i	2.5	4.4	1.8	2.6	2.9	2.6
	p	4.9	8.0	4.8	6.7	6.4	5.5
007	i	0.1	0.6	0.1	0.1	0.2	0.3
	p	1.1	2.7	1.4	1.1	1.4	1.6
008	i	0.4	0.5	0.6	0.5	0.5	0.4
	p	0.8	1.3	1.7	1.3	1.4	0.9
009	i	0.0	0.1	0.0	0.0	0.0	0.0
	p	0.1	0.2	0.1	0.2	0.2	0.1
010	i	0.8	0.9	1.6	1.4	1.3	0.8
	p	1.6	1.7	2.7	2.5	2.3	1.5
011	i	.	0.0	0.0	.	0.0	.
	p	0.0	0.1	0.1	0.1	0.1	0.0
012	i	0.3	0.2	0.1	0.2	0.1	0.2
	p	0.3	0.2	0.1	0.2	0.2	0.3
013	i	0.8	0.4	0.3	0.9	0.7	0.4
	p	0.9	0.4	0.3	1.0	0.7	0.5
014	i	.	.	0.0	0.0	0.0	0.0
	p	.	0.0	0.0	0.0	0.0	0.0
015	i	2.3	0.4	0.5	0.3	0.8	0.9
	p	2.4	0.4	0.5	0.5	0.9	0.9
016	i	0.4	0.3	0.7	0.5	0.5	0.4
	p	0.6	0.5	1.0	0.9	0.7	0.7
017	i	0.7	0.3	0.4	0.5	0.5	0.5
	p	0.8	0.4	0.4	0.7	0.5	0.6
018	i	4.2	4.1	6.1	9.0	5.8	5.4
	p	4.5	4.5	6.3	9.5	6.1	5.7
019	i	0.2	0.3	0.2	0.1	0.2	0.2
	p	0.3	0.5	0.5	0.2	0.4	0.3
020	i	0.0	0.2	0.1	0.1	0.1	0.0
	p	0.1	0.5	0.2	0.3	0.3	0.2
021	i	0.7	0.8	0.9	0.6	0.9	0.4
	p	1.2	1.1	1.5	1.0	1.3	0.9
022	i	0.0	0.0	0.0	0.0	0.0	.
	p	0.1	0.0	0.0	0.0	0.0	.
023	i	0.1	0.1	0.3	0.1	0.1	0.2
	p	0.3	0.4	0.8	0.6	0.5	0.5
024	i	0.0	0.1	0.1	0.1	0.1	0.0
	p	0.2	0.2	0.3	0.3	0.3	0.2
025	i	.	0.0	.	0.0	0.0	.
	p	.	0.0	0.0	0.0	0.0	0.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
026	poisoning by med. agents NOS	i	0.1	0.1	0.2	0.1	0.1	0.1	0.1
		p	0.2	0.2	0.3	0.2	0.2	0.2	0.1
027	adv. eff. med. agent proper dose: all.	i	0.7	0.7	0.7	0.6	0.3	0.7	0.6
		p	0.9	0.9	1.0	0.8	0.4	1.0	0.8
028	adv. eff. OAC	i	0.5	0.5	0.7	1.1	0.7	0.7	0.6
		p	0.8	0.7	0.9	1.6	0.9	0.9	0.8
029	adv. eff. med. agent NEC	i	1.1	1.1	1.6	1.7	1.4	1.3	1.1
		p	2.1	2.2	2.6	2.9	2.4	2.3	2.3
030	toxic eff. other substances	i	0.1	0.3	0.1	0.1	0.2	0.2	0.2
		p	0.2	0.5	0.3	0.2	0.2	0.3	0.3
031	complic. surg./med. treatm.: hematoma	i	0.0	0.1	0.0	.	0.0	0.0	0.1
		p	0.0	0.2	0.0	0.1	0.1	0.1	0.1
032	complic. surg./med. treatm.: wounds	i	0.1	0.2	0.2	0.1	0.2	0.1	0.2
		p	0.3	0.4	0.4	0.3	0.4	0.4	0.4
033	complic. surg./med. treatm.: NEC	i	0.0	0.0	.	0.1	0.0	0.0	0.0
		p	2.7	2.6	2.1	1.6	2.8	2.1	3.0
034	adv. eff. physical factors NEC	i	0.2	0.5	0.3	0.2	0.2	0.4	0.2
		p	0.7	0.7	0.6	0.6	0.5	0.8	0.5
035	adv. eff. prosthetic device	i	0.1	0.0	0.2	0.1	0.0	0.1	0.1
		p	0.2	0.1	0.4	0.1	0.2	0.2	0.2
036	mult.congenital syndromes	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.4	0.3	0.1	0.2	0.3	0.3
037	abnormal results investigation NEC	i	0.1	0.0	0.1	0.1	0.0	0.1	0.0
		p	0.1	0.3	0.2	0.2	0.1	0.2	0.1
038	toxoplasmosis	i	0.0	0.0	.	.	.	0.0	0.0
		p	0.0	0.0	0.1	0.1	.	0.0	0.1
039	premat./immature liveborn infant	i	0.0	0.0	0.1	.	0.0	0.0	0.0
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
040	all perinatal morbidity	i	0.1	0.1	0.0	0.1	0.1	0.1	0.1
		p	0.2	0.1	0.3	0.1	0.3	0.2	0.2
041	perinatal mortality	i	0.0	0.0	.
		p
042	death (excl. perinatal)	i	0.8	0.8	1.3	0.6	1.1	0.8	0.9
		p
043	no disease	i	9.8	11.5	10.4	12.0	10.4	9.5	14.2
		p	16.1	20.0	18.2	20.7	17.4	16.1	24.8
044	disease NEC	i	0.2	0.1	0.1	1.0	0.1	0.3	0.1
		p	0.4	0.2	0.5	1.5	0.2	0.6	0.3

A GENERAL AND UNSPECIFIED

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
026	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.1	0.2	0.2	0.3	0.2	0.1
027	i	0.5	0.5	1.0	0.7	0.6	0.7
	p	0.6	0.8	1.2	1.0	0.9	0.8
028	i	0.5	0.5	0.9	0.8	0.8	0.4
	p	0.7	0.7	1.1	1.0	1.1	0.5
029	i	1.7	1.2	1.3	0.9	1.5	1.0
	p	2.6	2.2	2.4	2.0	2.8	1.7
030	i	0.2	0.3	0.1	0.1	0.2	0.2
	p	0.5	0.4	0.2	0.2	0.4	0.2
<hr/>							
031	i	0.0	.	0.0	0.1	0.0	0.0
	p	0.2	0.0	0.0	0.1	0.1	0.0
032	i	0.2	0.1	0.2	0.2	0.2	0.1
	p	0.4	0.3	0.4	0.4	0.4	0.2
033	i	0.1	0.0	0.0	0.0	0.0	0.0
	p	1.6	2.8	2.4	2.9	2.5	2.0
034	i	0.5	0.4	0.2	0.2	0.3	0.3
	p	1.1	1.0	0.3	0.3	0.8	0.5
035	i	.	0.1	0.2	0.1	0.1	0.1
	p	0.0	0.3	0.4	0.1	0.3	0.2
<hr/>							
036	i	0.0	0.0	0.0	0.0	0.0	.
	p	0.4	0.1	0.3	0.2	0.3	0.2
037	i	0.0	0.1	0.1	0.0	0.1	0.1
	p	0.1	0.3	0.2	0.2	0.3	0.2
038	i	.	0.0	0.0	.	.	0.0
	p	0.0	0.1	0.0	0.0	0.0	0.1
039	i	.	0.0	0.0	0.0	0.0	.
	p	0.1	0.0	0.1	0.1	0.1	0.1
040	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.1	0.2	0.3	0.3	0.2	0.2
<hr/>							
041	i	.	0.0	.	.	0.0	.
	p
042	i	1.0	0.7	0.9	1.0	0.6	0.4
	p
043	i	13.8	10.4	11.1	7.8	11.4	9.1
	p	23.4	18.4	17.8	14.1	19.3	15.9
044	i	0.0	0.4	0.5	0.1	0.3	0.2
	p	0.1	0.7	0.9	0.1	0.5	0.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
045	enlarged lymph gland(s)	i	0.9	1.0	1.1	0.9	0.9	1.0	0.8
		p	1.1	1.2	1.4	1.2	1.2	1.3	1.0
046	sympt. lymph glands(s) NEC	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.1	.	0.1	0.0	0.1
047	fear of blood/lymph disease	i	0.2	0.2	0.2	0.5	0.2	0.2	0.1
		p	0.2	0.3	0.3	0.5	0.3	0.3	0.2
048	disability/impairment blood	i
		p
049	sympt./complt. blood NOS/NEC	i	.	0.0	0.0	.	0.0	.	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
050	acute lymphadenitis/-angitis	i	0.5	0.4	0.9	0.8	0.5	0.6	0.8
		p	0.5	0.5	1.0	0.7	0.6	0.6	0.9
051	chron. aspec. lymphadenitis	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.1	0.1	0.1	0.1	0.2	0.1	0.1
052	malignant lymphoma	i	0.0	0.0	.	0.0	0.0	0.0	0.0
		p	0.1	0.2	0.2	0.1	0.1	0.2	0.1
053	leukemia	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.3	0.3	0.2	0.3	0.3	0.3	0.2
054	malignant neopl. blood NEC	i	0.0	0.1	0.0	0.1	0.1	0.1	0.0
		p	0.3	0.2	0.3	0.2	0.4	0.3	0.2
055	benigne neopl. blood/bloodform. org.	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
056	ruptured spleen	i	.	0.0	0.0
		p	.	0.0	0.0
057	injuries blood/bloodform. org. NEC	i
		p	0.0	.	.	.	0.0	.	.
058	hered. haemolytic anaemias	i	0.0	.	0.0	.	0.0	.	0.0
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
059	cong. anom. blood/bloodform. org. NEC	i	.	.	.	0.0	.	0.0	.
		p	0.0	0.0	0.1	0.1	0.0	0.0	0.0
060	iron deficiency anemia	i	1.2	0.8	1.6	0.7	1.4	1.1	1.0
		p	6.7	3.3	4.1	3.5	6.6	4.5	4.1
061	pernic./other deficiency anemia	i	0.1	0.1	0.4	0.0	0.1	0.2	0.2
		p	2.4	1.4	2.7	1.5	2.7	2.1	1.6
062	anemia NOS/NEC	i	0.5	0.5	0.5	0.7	0.4	0.5	0.6
		p	2.3	1.5	1.7	2.3	1.6	2.0	1.9
063	hemophilia/coagulation disorders	i	0.1	0.1	0.0	0.2	0.1	0.1	0.0
		p	0.3	0.3	0.4	0.6	0.7	0.3	0.4
064	abnormal white cells	i	0.0	0.0
		p	0.1	0.0	0.0	0.1	0.0	0.0	0.1
065	unexplained abn. blood test	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.1	0.2	0.1	0.1	0.2	0.1	0.2
066	abn. blood test NOS	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.2	0.2	0.1	0.4	0.3	0.2	0.2
057	splenomegaly	i	.	.	0.1	.	.	0.0	.
		p	0.0	0.0	0.1	0.1	0.0	0.0	0.0
068	HIV-infection (AIDS/ARS)	i	.	0.0	0.0
		p	.	0.0	0.0	0.1	0.0	0.0	0.0
069	diseases blood/bloodform. org. NOS/NEC	i	0.1	0.1	0.0	.	0.1	0.0	0.1
		p	0.2	0.2	0.4	0.1	0.3	0.2	0.3

B BLOOD, BLOODFORM. ORG., LYMPHATICS, SPLEEN

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
045	i	0.9	0.8	1.2	1.0	1.0	0.9
	p	1.0	1.0	1.3	1.4	1.2	1.2
046	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.1
047	i	0.3	0.2	0.2	0.2	0.2	0.2
	p	0.4	0.2	0.2	0.3	0.3	0.2
048	i
	p
049	i	0.0	.	0.0	.	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
050	i	0.6	0.6	0.8	0.4	0.6	0.5
	p	0.6	0.6	0.9	0.5	0.7	0.6
051	i	0.2	0.1	0.1	0.0	0.1	0.1
	p	0.2	0.1	0.1	0.1	0.1	0.1
052	i	.	0.0	0.1	0.0	0.0	0.0
	p	0.1	0.1	0.2	0.2	0.1	0.1
053	i	0.0	.	0.0	0.1	0.0	0.0
	p	0.1	0.4	0.2	0.4	0.2	0.2
054	i	0.0	0.1	0.1	.	0.1	0.1
	p	0.2	0.3	0.3	0.2	0.3	0.2
055	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.0	0.0	0.0	.	0.0	0.0
056	i	.	.	.	0.0	0.0	.
	p	.	.	.	0.0	0.0	.
057	i
	p	.	0.0
058	i	0.0	.	.	0.0	0.0	0.0
	p	0.0	0.1	0.0	0.1	0.1	0.0
059	i	.	.	0.0	.	0.0	.
	p	0.0	0.0	0.1	0.0	0.0	0.0
060	i	1.2	1.3	0.7	1.1	1.1	1.3
	p	4.4	6.4	3.6	4.0	4.9	4.1
061	i	0.1	0.1	0.2	0.2	0.2	0.0
	p	2.0	2.4	1.9	1.8	2.3	1.4
062	i	0.4	0.5	0.6	0.5	0.4	0.6
	p	2.1	1.9	1.9	1.7	1.7	2.0
063	i	0.2	0.1	0.0	0.0	0.1	0.1
	p	0.5	0.4	0.3	0.4	0.4	0.3
064	i	.	.	.	0.0	0.0	.
	p	0.0	0.0	0.1	0.0	0.1	0.0
065	i	0.0	0.0	0.1	0.1	0.0	0.1
	p	0.1	0.1	0.2	0.2	0.2	0.1
066	i	0.1	0.0	0.0	0.0	0.0	0.0
	p	0.2	0.2	0.2	0.3	0.3	0.1
067	i	.	0.0	.	.	0.0	.
	p	0.0	0.1	.	0.0	0.0	0.0
068	i	.	.	0.0	.	0.0	.
	p	.	.	0.1	0.0	0.0	0.0
069	i	0.0	0.1	0.0	0.1	0.1	0.0
	p	0.2	0.3	0.3	0.2	0.3	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
070	generalized abd. pain/cramps	i	1.8	2.6	2.0	1.0	2.5	1.7	2.8
		p	3.3	3.9	3.1	1.5	3.4	2.8	4.3
071	stomach pain	i	0.7	0.8	0.6	1.0	0.7	0.7	0.9
		p	5.4	4.0	3.4	4.2	4.1	4.5	4.0
072	heartburn	i	0.3	0.3	0.5	0.0	0.5	0.3	0.5
		p	1.8	1.1	1.6	0.9	2.9	1.2	1.4
073	peri-anal itching	i	0.3	0.3	0.4	0.2	0.2	0.3	0.4
		p	0.4	0.5	0.8	0.3	0.3	0.5	0.7
074	localised abd. pain	i	0.5	0.5	0.8	0.7	0.5	0.5	0.7
		p	0.7	0.8	1.0	0.9	0.8	0.8	1.0
075	flatulence/belching	i	0.2	0.1	0.3	0.2	0.2	0.2	0.1
		p	0.3	0.2	0.4	0.2	0.2	0.3	0.2
076	nausea	i	0.2	0.4	0.2	0.1	0.2	0.2	0.3
		p	0.6	0.9	0.6	0.1	1.1	0.6	0.6
077	vomiting	i	0.3	0.2	0.1	0.3	0.2	0.2	0.3
		p	0.5	0.4	0.3	0.5	0.5	0.4	0.5
078	diarrhoea	i	1.4	1.1	1.1	1.0	1.1	1.2	1.2
		p	2.5	1.8	1.8	1.8	1.7	2.1	1.9
079	constipation	i	1.2	1.3	1.2	1.4	1.4	1.2	1.3
		p	7.0	5.6	5.6	5.0	7.0	6.0	5.6
080	jaundice	i	0.0	0.0	0.0	.	.	.	0.1
		p	0.0	0.0	0.0	.	0.0	0.0	0.1
081	blood vomiting	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.0	.	.	0.1	0.0	0.0
082	melena	i	0.1	0.1	0.0	0.2	0.1	0.1	0.1
		p	0.2	0.3	0.1	0.2	0.3	0.2	0.2
083	rectal bleeding	i	0.1	0.1	0.0	0.1	0.0	0.1	0.1
		p	0.1	0.2	0.1	0.1	0.1	0.1	0.2
084	fecal incontinence	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
085	defecation anomalies NEC	i	0.2	0.2	0.2	0.4	0.1	0.2	0.2
		p	0.2	0.2	0.3	0.4	0.3	0.3	0.2
086	sympt./complt. of teeth, gums	i	0.3	0.2	0.5	0.2	0.2	0.2	0.5
		p	1.1	0.4	0.7	0.4	1.0	0.6	0.8
087	sympt./complt. mouth, tongue, lip	i	0.4	0.5	0.2	0.3	0.4	0.4	0.5
		p	0.8	0.7	0.6	0.3	0.6	0.7	0.6
088	swallowing problems	i	0.2	0.5	0.5	0.2	0.4	0.3	0.4
		p	0.4	0.6	0.6	0.4	0.6	0.4	0.6
089	fear of digestive diseases	i	0.1	0.1	0.2	0.2	0.0	0.1	0.2
		p	0.1	0.1	0.5	0.5	0.1	0.3	0.3
090	digestive disability/impairment	i	0.0	0.0
		p	0.0	0.0
091	sympt./complt. digestive NEC	i	0.3	0.3	0.3	0.1	0.3	0.2	0.3
		p	0.5	0.4	0.6	0.2	0.6	0.4	0.5
092	infectious diarrhoea/dysentery	i	0.3	0.5	0.3	0.5	0.3	0.5	0.2
		p	0.4	0.6	0.4	0.8	0.4	0.6	0.2
093	mumps	i	1.3	0.8	0.5	0.5	0.3	1.0	0.6
		p	1.3	0.8	0.5	0.5	0.3	1.1	0.6
094	infectious hepatitis	i	0.0	0.0	0.1	.	.	0.0	0.0
		p	0.1	0.1	0.1	0.2	0.1	0.2	0.1

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
070	i	2.0	1.6	2.1	2.5	2.1	1.8
	p	2.9	3.0	3.3	3.8	3.4	2.7
071	i	0.9	0.9	0.5	0.7	0.9	0.4
	p	3.8	4.8	3.9	4.9	5.0	3.1
072	i	0.2	0.3	0.5	0.4	0.4	0.2
	p	1.1	1.4	1.4	1.7	1.6	1.0
073	i	0.4	0.3	0.3	0.3	0.3	0.3
	p	0.6	0.4	0.6	0.5	0.6	0.4
074	i	0.6	0.5	0.6	0.6	0.6	0.5
	p	0.9	0.8	0.9	0.8	0.9	0.7
075	i	0.2	0.1	0.3	0.2	0.2	0.1
	p	0.3	0.2	0.4	0.2	0.3	0.1
076	i	0.3	0.2	0.2	0.3	0.3	0.2
	p	0.6	0.8	0.5	0.7	0.7	0.6
077	i	0.3	0.2	0.2	0.2	0.2	0.1
	p	0.5	0.5	0.4	0.3	0.5	0.3
078	i	1.6	1.4	1.0	0.8	1.3	0.8
	p	2.1	2.6	1.7	1.5	2.3	1.5
079	i	1.0	1.5	1.1	1.3	1.4	0.9
	p	5.3	6.9	6.0	5.6	7.1	3.9
080	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	.	0.0	0.0	0.0	0.0
081	i	0.0	0.0	.	0.0	0.0	.
	p	0.0	0.0	.	0.0	0.0	.
082	i	0.0	0.1	0.1	0.2	0.1	0.1
	p	0.1	0.3	0.2	0.3	0.3	0.1
083	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.1	0.2	0.1	0.1	0.1	0.1
084	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.0
085	i	0.2	0.2	0.2	0.2	0.2	0.2
	p	0.2	0.3	0.3	0.4	0.3	0.3
086	i	0.4	0.2	0.4	0.2	0.4	0.1
	p	0.7	0.8	0.8	0.5	0.8	0.5
087	i	0.4	0.3	0.4	0.5	0.4	0.3
	p	0.7	0.5	0.7	0.7	0.7	0.5
088	i	0.4	0.3	0.4	0.3	0.4	0.3
	p	0.5	0.4	0.6	0.5	0.6	0.4
089	i	0.2	0.1	0.1	0.1	0.1	0.1
	p	0.3	0.1	0.4	0.2	0.3	0.2
090	i	.	.	0.0	.	0.0	.
	p	.	.	0.0	.	0.0	.
091	i	0.3	0.2	0.2	0.3	0.3	0.2
	p	0.5	0.5	0.4	0.4	0.5	0.3
092	i	0.6	0.4	0.3	0.3	0.4	0.3
	p	0.8	0.4	0.4	0.3	0.5	0.4
093	i	2.1	0.9	0.3	0.1	0.7	1.0
	p	2.1	0.9	0.4	0.1	0.7	1.1
094	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.2	0.1	0.1	0.1	0.1	0.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
095	parasites/worms	i	3.5	3.2	3.7	2.1	3.3	3.2	3.4
		p	3.8	3.3	3.9	2.1	3.4	3.3	3.8
096	other presumed infections	i	6.2	6.5	7.5	6.2	5.5	5.9	9.0
		p	6.6	6.9	7.8	6.7	5.8	6.2	9.3
097	malign. neopl. stomach	i	0.0	.	0.1	.	0.0	0.0	0.0
		p	0.3	0.2	0.3	0.1	0.3	0.2	0.3
098	malign. neopl. colon, rectum	i	0.1	0.0	0.1	0.2	0.1	0.1	0.1
		p	0.7	0.5	1.1	0.5	0.7	0.7	0.8
099	malign. neopl. colon, rectum	i	0.1	0.0	0.0	0.1	0.0	0.1	0.0
		p	0.3	0.3	0.2	0.4	0.2	0.3	0.3
100	malign. neopl. pancreas	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.1	0.0	0.2	0.2	0.1	0.1	0.1
101	malign. neopl. dig. syst. NEC	i	0.0	0.0	0.1	0.2	0.1	0.1	0.1
		p	0.4	0.4	0.4	1.0	0.4	0.5	0.3
102	benign. neoplasms dig. syst. NEC	i	0.1	0.0	0.0	.	0.0	0.0	0.1
		p	0.1	0.1	0.2	.	0.1	0.1	0.1
103	neoplasms dig. syst. NEC	i	0.1	0.1	0.0	.	0.0	0.0	0.1
		p	0.1	0.2	0.2	0.1	0.1	0.1	0.3
104	foreign body through orifice	i	0.1	0.1	0.1	0.1	0.1	0.1	0.0
		p	0.1	0.1	0.1	0.1	0.1	0.1	0.1
105	injuries dig. syst. NEC	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
106	cong. anom. digest. system	i	0.1	0.0	0.0	.	0.0	0.0	0.1
		p	0.2	0.1	0.1	0.2	0.2	0.1	0.2
107	disease of teeth/gums	i	0.8	0.7	1.6	0.7	0.8	0.8	1.2
		p	1.5	1.0	2.0	1.3	1.3	1.3	1.7
108	disease of mouth/tongue/lips	i	2.0	1.9	3.0	2.1	2.0	2.2	2.2
		p	2.8	2.7	3.8	2.3	2.6	3.0	2.9
109	disease of esophagus	i	0.4	0.5	1.0	0.8	0.3	0.6	0.6
		p	0.9	1.1	1.6	1.2	1.1	1.1	1.3
110	duodenal ulcer	i	0.5	0.5	1.3	0.6	0.6	0.7	0.5
		p	1.5	1.4	3.7	2.2	2.0	2.1	1.9
111	gastric ulcers NEC	i	0.3	0.2	0.1	0.2	0.2	0.2	0.2
		p	0.9	0.6	0.6	1.6	1.0	0.9	0.7
112	peptic ulcers NEC	i	0.1	0.1	0.2	0.1	0.1	0.1	0.1
		p	0.4	0.4	0.4	0.4	0.4	0.4	0.5
113	disord. stomach function	i	2.4	3.7	3.3	3.2	2.4	2.9	4.2
		p	4.4	5.4	5.3	5.7	5.1	4.4	6.7
114	appendicitis	i	0.4	0.3	0.6	0.4	0.4	0.4	0.5
		p	0.5	0.4	0.8	0.4	0.5	0.5	0.5
115	inguinal hernia	i	0.7	0.7	0.3	0.5	0.5	0.6	0.5
		p	0.8	1.0	0.5	0.5	0.8	0.8	0.8
116	hiatus (diaphr.) hernia	i	0.4	0.5	1.7	0.5	0.4	0.9	0.5
		p	1.4	1.8	3.6	1.6	1.5	2.2	1.8
117	abdominal hernia NEC	i	0.2	0.2	0.2	0.3	0.3	0.2	0.1
		p	0.3	0.2	0.3	0.4	0.3	0.3	0.2
118	diverticular dis. intestines	i	0.2	0.3	0.5	0.6	0.2	0.4	0.3
		p	0.6	0.8	1.3	1.0	0.6	0.9	0.9
119	irritable bowel syndrome	i	2.5	4.5	4.6	5.9	2.5	4.5	3.4
		p	3.8	6.9	7.3	8.4	4.2	6.8	5.1

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
095	i	3.3	2.8	3.4	3.7	3.4	3.0
	p	3.3	3.0	3.7	3.9	3.6	3.1
096	i	6.3	6.0	5.9	8.1	7.3	4.9
	p	6.8	6.3	6.3	8.5	7.6	5.3
097	i	0.0	0.1	0.1	.	0.1	0.0
	p	0.2	0.3	0.3	0.2	0.2	0.2
098	i	0.0	0.1	0.1	0.0	0.1	0.0
	p	0.6	0.9	0.7	0.7	0.7	0.7
099	i	.	0.1	0.1	0.0	0.0	0.1
	p	0.1	0.4	0.4	0.3	0.3	0.3
100	i	0.0	0.1	0.0	0.0	0.0	0.0
	p	0.1	0.2	0.1	0.1	0.1	0.0
101	i	0.0	0.1	0.1	0.0	0.1	0.1
	p	0.3	0.6	0.6	0.3	0.5	0.3
102	i	0.0	0.1	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.1
103	i ^x	0.1	0.0	0.0	0.1	0.1	0.0
	p	0.1	0.1	0.2	0.2	0.2	0.1
104	i	0.1	.	0.0	0.2	0.0	0.1
	p	0.1	0.0	0.1	0.3	0.1	0.2
105	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	.	0.0	0.0	0.0	0.0
106	i	0.0	0.0	0.1	0.1	0.0	0.0
	p	0.2	0.1	0.2	0.2	0.1	0.2
107	i	0.7	0.7	1.1	1.0	1.0	0.7
	p	1.2	1.2	1.6	1.8	1.6	1.1
108	i	2.3	2.4	2.4	1.8	2.4	1.9
	p	3.0	3.3	3.0	2.4	3.1	2.5
109	i	0.6	0.7	0.7	0.5	0.7	0.3
	p	1.0	1.0	1.3	1.2	1.4	0.6
110	i	0.5	0.6	0.9	0.7	0.7	0.6
	p	2.1	1.2	2.8	2.0	2.3	1.5
111	i	0.2	0.2	0.2	0.2	0.2	0.1
	p	0.8	1.0	0.9	0.6	0.9	0.5
112	i	0.1	0.0	0.1	0.2	0.1	0.0
	p	0.4	0.3	0.3	0.6	0.5	0.2
113	i	3.7	2.2	3.7	3.0	3.5	2.2
	p	5.4	3.7	5.4	5.8	5.8	3.5
114	i	0.4	0.6	0.5	0.2	0.4	0.4
	p	0.5	0.6	0.6	0.3	0.5	0.5
115	i	0.6	0.6	0.4	0.7	0.5	0.6
	p	0.9	0.8	0.6	0.9	0.8	0.8
116	i	0.4	1.4	0.6	0.5	0.7	0.8
	p	1.9	2.3	2.3	1.6	2.2	1.7
117	i	0.2	0.2	0.3	0.2	0.2	0.1
	p	0.2	0.3	0.4	0.2	0.3	0.2
118	i	0.3	0.4	0.5	0.2	0.4	0.4
	p	0.8	0.9	1.0	0.8	1.0	0.8
119	i	4.1	2.6	4.3	5.0	4.3	3.3
	p	6.5	4.2	6.3	7.6	6.7	5.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
120	chron. enteritis/colitis ulc.	i	0.2	0.2	0.2	0.1	0.3	0.2	0.3
		p	1.1	1.1	0.8	0.8	1.2	0.9	1.1
121	anal fissure/perianal abscess	i	0.6	0.9	0.8	0.7	0.7	0.8	0.7
		p	0.9	1.3	1.1	0.9	0.9	1.1	1.1
122	hepatomegaly	i	.	0.0	0.1	.	0.0	0.0	0.0
		p	0.0	0.0	0.1	.	0.0	0.0	0.1
123	cirrhosis/liver disease NEC	i	0.1	0.2	0.2	0.5	0.1	0.2	0.2
		p	0.4	0.7	0.7	1.6	0.6	0.7	0.7
124	diseases of gall-bladder/biliary duct	i	0.5	0.4	0.8	0.6	0.4	0.6	0.6
		p	1.4	1.2	1.5	1.6	1.4	1.3	1.4
125	diseases digestive system NEC	i	0.3	0.6	0.6	0.7	0.4	0.5	0.4
		p	1.2	1.3	1.1	1.2	1.3	1.2	1.2

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
120	i	0.3	0.2	0.2	0.2	0.2	0.2
	p	1.0	1.1	1.0	0.8	1.1	0.9
121	i	0.9	0.7	0.9	0.6	0.8	0.6
	p	1.2	1.0	1.3	0.9	1.2	0.8
122	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.1	0.0	0.0
123	i	0.2	0.1	0.2	0.3	0.2	0.1
	p	0.6	0.6	0.7	0.9	0.8	0.4
124	i	0.4	0.8	0.6	0.4	0.6	0.5
	p	0.9	1.5	1.3	1.6	1.5	1.1
<hr/>							
125	i	0.6	0.5	0.5	0.4	0.5	0.4
	p	1.2	1.2	1.4	1.0	1.3	0.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
126	complaints of vision	i	0.7	0.8	1.0	0.9	0.6	0.8	1.1
		p	2.4	2.3	2.2	3.2	1.1	2.1	3.7
127	fear of eye disease	i	0.0	0.0	0.0	.	0.1	0.0	0.0
		p	0.0	0.0	0.0	.	0.1	0.0	0.0
128	disability/impairment eye	i	.	0.0	.	.	0.0	.	.
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.1
129	sympt./complt. eye NEC	i	1.1	1.0	1.0	1.2	1.0	1.1	1.0
		p	2.8	2.1	2.1	2.4	2.3	2.5	2.0
130	infectious conjunctivitis	i	1.7	1.9	1.9	1.5	1.7	1.8	1.8
		p	1.9	2.0	2.0	1.8	1.8	2.0	1.9
131	allergic conjunctivitis	i	5.1	5.8	6.3	3.5	5.4	5.4	5.4
		p	5.7	6.5	7.0	3.6	5.9	6.1	6.0
132	blepharitis/stye/chalazion	i	1.2	1.5	1.4	0.9	1.2	1.2	1.5
		p	1.3	1.6	1.6	1.1	1.3	1.4	1.8
133	infection of eye NEC	i	0.5	0.5	0.5	0.5	0.5	0.5	0.5
		p	1.0	0.9	1.0	0.7	1.0	0.9	0.9
134	corneal ulcer	i	0.1	0.0	0.1	.	0.0	0.1	0.0
		p	0.1	0.0	0.1	0.1	0.0	0.1	0.0
135	neoplasms of eye/adnexa	i	0.1	0.0	.	.	0.1	0.0	0.0
		p	0.1	0.1	0.1	0.1	0.1	0.1	0.0
136	contusion/abrasion/black eye	i	0.6	0.7	0.6	0.7	0.5	0.7	0.7
		p	0.6	0.8	0.7	0.8	0.6	0.7	0.8
137	foreign body in eye	i	2.2	1.3	1.0	0.2	1.9	1.2	1.8
		p	2.5	1.5	1.0	0.4	2.0	1.4	1.9
138	injuries eye NEC	i	0.2	0.2	0.2	0.2	0.2	0.2	0.3
		p	0.3	0.2	0.3	0.3	0.3	0.3	0.4
139	blocked lacrimal duct of infants	i	0.1	0.0	.	.	0.1	0.0	0.1
		p	0.1	0.0	.	0.1	0.1	0.0	0.1
140	cong. anom. of eye NEC	i	0.1	0.0	.	.	0.0	0.0	0.0
		p	0.1	0.1	.	.	0.0	0.0	0.1
141	refractive errors	i	0.9	0.9	1.3	1.0	1.2	0.9	1.0
		p	4.0	4.6	5.1	3.6	5.9	4.1	4.6
142	cataract	i	0.2	0.2	0.2	0.2	0.2	0.2	0.2
		p	1.4	0.9	1.5	1.0	1.1	1.3	1.0
143	glaucoma	i	0.1	0.1	0.4	0.2	0.1	0.2	0.2
		p	2.1	0.9	1.2	0.6	2.0	1.4	0.9
144	blindness, all types	i	0.0	0.0	.	0.1	0.0	0.0	0.0
		p	0.3	0.2	0.3	0.3	0.2	0.2	0.3
145	strabismus	i	0.2	0.1	0.1	0.1	0.2	0.1	0.2
		p	0.6	0.4	0.4	0.2	0.5	0.4	0.6
146	retinopathy	i	0.2	0.2	0.0	0.3	0.0	0.2	0.1
		p	0.4	0.5	0.6	0.3	0.3	0.5	0.4
147	diseases of eye NEC	i	0.6	0.7	0.8	0.8	0.8	0.7	0.6
		p	1.1	1.2	1.2	2.0	1.4	1.2	1.3

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
126	i	0.7	0.5	1.2	0.9	0.9	0.5
	p	1.8	2.0	2.6	3.3	3.0	1.0
127	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
128	i	.	.	.	0.0	0.0	.
	p	0.0	0.1	0.0	0.0	0.0	0.0
129	i	1.1	0.9	1.2	1.0	1.2	0.8
	p	1.7	2.1	2.8	2.7	2.7	1.7
130	i	3.4	1.5	1.1	1.3	1.9	1.5
	p	3.6	1.6	1.2	1.4	2.0	1.6
131	i	4.8	5.3	5.9	5.7	5.6	4.7
	p	5.2	5.9	6.6	6.3	6.4	5.2
132	i	1.2	1.1	1.6	1.3	1.4	0.9
	p	1.4	1.3	1.7	1.5	1.6	1.1
133	i	0.4	0.7	0.5	0.4	0.5	0.5
	p	0.8	1.2	0.9	0.9	1.0	0.9
134	i	.	0.1	0.1	0.0	0.0	0.1
	p	0.0	0.1	0.1	0.1	0.0	0.1
135	i	0.0	0.1	0.0	.	0.0	0.0
	p	0.1	0.2	0.1	.	0.1	0.1
136	i	0.9	0.4	0.6	0.7	0.7	0.5
	p	1.0	0.4	0.7	0.8	0.8	0.7
137	i	1.1	1.7	1.5	1.3	1.6	1.0
	p	1.3	1.9	1.7	1.6	1.8	1.1
138	i	0.3	0.2	0.1	0.2	0.2	0.1
	p	0.4	0.3	0.2	0.3	0.3	0.2
139	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.1	0.1	0.1	0.0
140	i	0.0	0.1	0.0	.	0.0	0.0
	p	0.1	0.1	0.0	0.1	0.1	0.0
141	i	0.8	0.7	1.2	1.1	1.2	0.6
	p	5.0	3.8	5.0	3.8	6.0	1.3
142	i	0.2	0.3	0.1	0.1	0.2	0.1
	p	0.8	1.9	1.3	0.9	1.6	0.4
143	i	0.2	0.1	0.2	0.2	0.2	0.2
	p	1.1	1.4	1.5	1.4	1.6	0.9
144	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.1	0.3	0.3	0.2	0.2	0.2
145	i	0.2	0.1	0.2	0.1	0.2	0.0
	p	0.5	0.3	0.5	0.4	0.5	0.2
146	i	0.1	0.2	0.3	0.1	0.1	0.2
	p	0.5	0.5	0.6	0.4	0.5	0.3
147	i	0.8	0.6	0.7	0.6	0.8	0.5
	p	1.3	1.1	1.2	1.4	1.5	0.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
148	earache	i	0.2	0.2	0.3	0.3	0.2	0.2	0.4
		p	0.6	0.6	0.5	0.6	0.4	0.6	0.7
149	hearing complt.	i	0.2	0.2	0.1	0.1	0.2	0.2	0.2
		p	0.5	0.5	0.3	0.3	0.3	0.4	0.7
150	ringing/buzzing/tinnitus	i	0.2	0.2	0.2	0.2	0.2	0.2	0.2
		p	0.3	0.2	0.3	0.4	0.3	0.3	0.3
151	blood/discharge from ear	i	0.1	0.2	0.1	0.1	0.1	0.1	0.2
		p	0.2	0.4	0.1	0.1	0.2	0.2	0.3
152	fear of ear disease	i	0.0	0.0	.	.	0.0	.	0.0
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
153	disability/impairment of ear	i	.	0.0	0.0
		p	0.0	0.0	0.0
154	symp./complt. of ear NEC	i	0.2	0.3	0.3	0.3	0.3	0.3	0.3
		p	0.5	0.4	0.5	0.4	0.4	0.4	0.5
155	otitis externa	i	3.6	3.2	4.0	2.6	3.1	3.3	3.8
		p	5.2	4.1	5.0	3.9	4.7	4.4	5.1
156	acute otitis media/myringitis	i	5.6	7.4	7.0	6.9	5.8	6.8	6.8
		p	5.8	7.8	7.3	7.2	5.9	7.0	7.2
157	serous otitis media/glue ear	i	1.4	1.1	2.2	1.1	1.0	1.7	0.9
		p	1.8	1.8	2.9	1.7	1.4	2.4	1.4
158	eustachian tube dysfunction	i	2.1	2.6	2.3	1.4	2.7	2.1	2.3
		p	2.5	3.2	3.0	1.7	3.1	2.7	2.8
159	infections of ear NEC	i	0.3	0.4	0.4	0.2	0.3	0.3	0.6
		p	0.6	0.6	0.9	0.3	0.6	0.6	0.9
160	neoplasm of ear	i	0.0	0.0	.	0.1	.	0.0	.
		p	0.0	0.0	.	0.1	.	0.0	0.0
161	foreign body in ear	i	0.1	0.1	0.2	0.1	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.1	0.1	0.1	0.2
162	perforation tymp. membrane	i	0.1	0.1	0.3	0.1	0.2	0.1	0.2
		p	0.3	0.2	0.5	0.4	0.3	0.3	0.3
163	superficial injury of ear	i	0.1	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.0	.	0.0	0.0	0.1
164	injuries of ear NEC	i	0.0	0.1	0.1	0.1	0.0	0.1	0.1
		p	0.0	0.1	0.2	0.1	0.1	0.1	0.1
165	cong. anom. of ear	i	0.0	0.0	0.1	.	0.0	0.0	0.0
		p	0.0	0.0	0.1	.	0.0	0.1	0.0
166	ear wax (excessive)	i	8.0	8.2	9.0	8.2	9.3	8.2	8.2
		p	9.0	9.2	10.0	8.9	10.3	9.1	9.1
167	vertiginous syndromes	i	0.3	0.6	0.8	0.8	0.4	0.5	0.8
		p	0.9	1.1	1.4	1.3	1.2	1.0	1.5
168	otosclerosis	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.0	0.1	0.2	0.2	0.1	0.1	0.2
169	presbycusis	i	0.4	0.4	0.4	0.3	0.5	0.4	0.3
		p	0.5	0.6	0.5	0.7	0.6	0.6	0.5
170	acoustic trauma	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
171	deafness NEC	i	0.4	0.3	0.2	0.2	0.3	0.3	0.4
		p	0.8	0.6	0.6	0.9	0.7	0.7	0.8
172	disease of ear/mastoid NEC	i	0.2	0.2	0.4	0.4	0.2	0.3	0.2
		p	0.3	0.3	0.7	0.8	0.4	0.5	0.4

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
148	i	0.4	0.1	0.3	0.2	0.3	0.2
	p	0.8	0.4	0.6	0.6	0.6	0.5
149	i	0.1	0.1	0.2	0.3	0.2	0.2
	p	0.2	0.3	0.6	0.7	0.4	0.4
150	i	0.1	0.2	0.2	0.2	0.2	0.2
	p	0.3	0.4	0.3	0.2	0.2	0.4
151	i	0.1	0.1	0.1	0.2	0.1	0.1
	p	0.3	0.2	0.2	0.3	0.2	0.2
152	i	.	0.0	.	0.0	0.0	.
	p	.	0.0	0.0	0.0	0.0	.
153	i	.	.	0.0	.	0.0	.
	p	.	.	0.0	.	0.0	.
154	i	0.2	0.2	0.3	0.4	0.3	0.2
	p	0.4	0.3	0.6	0.5	0.5	0.3
155	i	2.9	4.1	4.0	2.7	3.4	3.4
	p	4.0	5.5	5.1	3.8	4.7	4.3
156	i	7.9	3.8	6.4	8.5	6.5	6.9
	p	8.3	4.1	6.6	8.9	6.8	7.1
157	i	1.1	1.0	1.6	2.0	1.4	1.5
	p	1.9	1.3	2.3	2.6	2.0	2.0
158	i	2.3	1.9	2.5	2.3	2.2	2.1
	p	2.6	2.4	3.1	2.9	2.8	2.5
159	i	0.5	0.2	0.3	0.3	0.4	0.2
	p	0.8	0.5	0.6	0.6	0.7	0.4
160	i	0.1	.	.	0.0	0.0	.
	p	0.1	.	0.0	0.0	0.0	0.0
161	i	0.2	0.1	0.2	0.1	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.1
162	i	0.2	0.1	0.2	0.1	0.1	0.2
	p	0.5	0.2	0.3	0.3	0.3	0.2
163	i	0.0	0.1	0.0	.	0.0	0.0
	p	0.0	0.1	0.0	0.1	0.1	0.0
164	i	0.0	0.0	0.2	.	0.1	0.1
	p	0.0	0.0	0.2	.	0.1	0.1
165	i	0.0	0.0	0.0	0.1	0.0	0.0
	p	0.0	0.0	0.1	0.1	0.0	0.0
166	i	8.3	10.0	7.0	7.9	8.2	8.2
	p	9.5	11.1	7.5	9.0	9.3	9.0
167	i	0.7	0.6	0.8	0.3	0.6	0.7
	p	1.1	1.2	1.4	0.9	1.1	1.2
168	i	0.0	0.0	0.1	0.1	0.0	0.1
	p	0.1	0.0	0.1	0.1	0.0	0.1
169	i	0.4	0.5	0.3	0.3	0.4	0.3
	p	0.7	0.7	0.4	0.5	0.6	0.4
170	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.1	0.0	0.1	0.2	0.1	0.1
171	i	0.3	0.4	0.3	0.3	0.3	0.3
	p	0.5	0.9	0.7	0.8	0.8	0.5
172	i	0.2	0.3	0.5	0.1	0.3	0.2
	p	0.4	0.4	0.7	0.3	0.5	0.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
173	complaints of heart	i	0.4	0.4	0.8	0.4	0.4	0.4	0.7
		p	1.2	1.1	1.7	0.8	1.1	1.2	1.3
174	edema of extremities	i	0.9	0.8	0.9	0.6	0.9	0.8	0.9
		p	6.6	3.2	5.3	2.4	6.5	4.2	5.1
175	fear of dis. circ. system	i	0.4	0.4	0.8	0.6	0.6	0.4	0.7
		p	0.7	0.8	1.3	0.9	1.1	0.8	1.1
176	disability/impairment circ. syst. NEC	i
		p	0.0	.	0.0	.	.	0.0	0.0
177	sympt./complt. circ. syst. NEC	i	0.2	0.1	0.2	0.2	0.2	0.1	0.2
		p	0.5	0.2	0.2	0.6	0.5	0.3	0.4
178	infections dis. circ. syst.	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.0	0.1	0.3	0.1	0.1	0.0
179	neoplasm circ. syst.	i	0.0	0.1	0.0	.	.	0.0	0.1
		p	0.0	0.1	0.0	0.1	0.0	0.0	0.1
180	injuries heart/circ. syst.	i	0.0	0.0	.	.	0.0	.	0.0
		p	0.0	0.0	0.0	.	0.0	.	0.1
181	cong. anom. heart/circ. syst.	i	0.0	0.0	.	.	0.1	0.0	.
		p	0.3	0.2	0.4	0.0	0.4	0.3	0.2
182	rheumatic fever/heart dis.	i	0.0	0.0	0.0	0.1	.	0.0	0.0
		p	0.1	0.1	0.0	0.4	0.1	0.1	0.0
183	acute myocardial infarction	i	0.4	0.4	0.6	0.4	0.4	0.4	0.6
		p
184	angina pectoris	i	1.2	1.0	1.5	2.0	1.2	1.2	1.5
		p	10.9	6.5	7.0	5.8	12.3	7.8	6.6
185	healed myocardial infarction	i
		p	5.3	4.8	5.1	3.0	8.1	4.1	5.4
186	chron. ischemic heart dis. NEC	i	0.2	0.2	0.2	0.1	0.2	0.1	0.2
		p	6.7	3.9	5.4	3.0	4.7	3.6	8.8
187	heart failure	i	1.0	0.7	0.8	1.3	0.8	0.9	0.8
		p	7.4	5.9	5.2	7.3	7.0	6.4	6.0
188	atrial fibrillation/flutter	i	0.2	0.2	0.3	0.2	0.2	0.2	0.1
		p	2.2	2.3	2.7	2.5	2.5	2.3	2.5
189	paroxysmal tachycardia	i	0.2	0.3	0.1	0.4	0.2	0.2	0.3
		p	0.5	0.7	0.7	0.7	0.7	0.6	0.7
190	ectopic beats, all types	i	0.2	0.2	0.2	0.2	0.2	0.2	0.1
		p	0.3	0.4	0.4	0.8	0.6	0.4	0.3
191	heart murmur NOS	i	0.1	0.0	0.1	0.2	0.0	0.1	0.0
		p	0.2	0.1	0.2	0.3	0.2	0.2	0.1
192	pulmonary heart dis.	i	0.0	.	.	.	0.0	0.0	.
		p	0.1	0.1	0.2	0.1	0.2	0.1	0.1
193	heart valve dis. NOS, non-rheum.	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.8	0.4	0.8	1.4	0.9	0.7	0.7
194	heart disease NEC	i	0.4	0.4	0.4	0.6	0.3	0.4	0.4
		p	3.2	2.5	2.9	2.4	4.6	2.4	2.9
195	elevated blood pressure w.o. hypertension	i	0.1	0.2	0.3	0.3	0.3	0.2	0.1
		p	0.5	1.9	1.7	2.0	1.0	1.8	0.6
196	uncomplicated hypertension	i	2.7	2.5	4.7	3.1	2.9	3.5	2.2
		p	48.6	41.7	44.0	45.5	52.4	44.3	43.1
197	hypertension with involv.targ. organs	i	0.3	0.3	0.5	0.7	0.3	0.5	0.3
		p	9.8	5.8	5.9	5.4	10.1	6.5	7.4

K CIRCULATORY

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
173	i	0.5	0.3	0.5	0.6	0.5	0.4
	p	1.1	1.0	1.3	1.4	1.3	1.0
174	i	0.9	1.4	0.6	0.4	1.0	0.6
	p	4.0	8.2	3.7	2.9	5.4	3.1
175	i	0.6	0.4	0.3	0.7	0.6	0.5
	p	1.0	0.8	0.8	1.0	1.0	0.9
176	i
	p	.	0.0	0.0	.	0.0	0.0
177	i	0.1	0.1	0.2	0.2	0.2	0.1
	p	0.2	0.3	0.4	0.5	0.4	0.2
178	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.2	0.1	0.0	0.1	0.1
179	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.0	0.1	0.1	0.0
180	i	.	0.0	.	.	.	0.0
	p	0.0	0.0	.	0.0	0.0	0.0
181	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.2	0.2	0.4	0.3	0.3	0.2
182	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.1
183	i	0.5	0.3	0.5	0.4	0.4	0.4
	p
184	i	0.9	1.2	1.7	1.3	1.3	1.0
	p	5.7	9.9	8.2	8.0	9.4	5.5
185	p	3.9	5.8	4.4	5.2	5.5	3.5
	i
186	i	0.1	0.2	0.2	0.0	0.1	0.2
	p	3.5	3.7	7.2	5.7	5.7	3.5
187	i	0.7	1.0	0.8	0.9	1.0	0.4
	p	5.7	6.6	8.4	4.8	7.2	4.1
188	i	0.1	0.3	0.3	0.1	0.2	0.3
	p	1.6	2.6	3.5	1.8	2.7	1.8
189	i	0.2	0.2	0.3	0.3	0.3	0.2
	p	0.8	0.6	0.7	0.6	0.7	0.6
190	i	0.1	0.3	0.2	0.1	0.2	0.1
	p	0.4	0.5	0.6	0.2	0.4	0.4
191	i	0.1	0.1	0.1	0.0	0.0	0.1
	p	0.2	0.2	0.2	0.1	0.2	0.2
192	i	0.0	.	0.0	0.0	0.0	.
	p	0.1	0.1	0.2	0.1	0.2	0.0
193	i	0.0	0.0	0.0	0.0	0.1	0.0
	p	0.7	0.5	1.1	0.6	0.8	0.5
194	i	0.3	0.7	0.3	0.3	0.4	0.4
	p	2.2	3.3	3.0	2.5	3.0	2.3
195	i	0.2	0.2	0.3	0.1	0.1	0.2
	p	1.8	1.0	1.8	1.1	1.3	1.3
196	i	2.8	5.0	2.2	2.5	3.0	3.5
	p	39.8	51.1	43.4	45.2	46.8	42.3
197	i	0.3	0.6	0.4	0.2	0.4	0.4
	p	5.3	8.2	8.5	6.3	8.4	5.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
198	(postural) hypotension	i	0.5	0.6	1.0	0.9	0.7	0.7	0.7
		p	0.9	1.0	1.6	1.1	1.0	1.0	1.3
199	transient cerebral ischemia	i	0.5	0.5	0.9	0.5	0.6	0.6	0.7
		p	1.6	1.5	1.9	1.7	1.9	1.6	1.7
200	cerebrovascular diseases	i	0.7	0.5	0.7	1.1	0.7	0.7	0.6
		p	3.0	2.9	3.4	4.1	3.2	3.1	3.3
201	atherosclerosis, excl. heart/brain	i	0.1	0.1	0.2	0.1	0.1	0.1	0.1
		p	1.1	0.8	1.2	0.2	0.9	0.7	1.4
202	arterial vascular diseases NEC	i	0.6	0.5	0.7	0.6	0.6	0.5	0.8
		p	3.9	2.5	3.1	2.7	3.2	2.6	4.6
203	pulmonary embolism	i	0.1	0.1	0.1	0.1	0.1	0.1	0.0
		p	0.4	0.2	0.6	0.1	0.5	0.3	0.3
204	(thrombo) phlebitis	i	0.4	0.6	0.4	0.3	0.4	0.4	0.5
		p	0.7	0.8	0.8	0.5	0.6	0.7	0.8
205	venous thrombosis	i	0.1	0.1	0.3	0.3	0.2	0.2	0.2
		p	1.4	0.7	1.2	1.2	1.4	1.1	0.8
206	various veins of legs	i	0.9	1.4	1.7	1.0	1.1	1.2	1.5
		p	3.3	3.5	4.5	2.7	3.6	3.4	4.0
207	various ulcer of skin	i	0.3	0.3	0.3	0.3	0.5	0.3	0.2
		p	1.4	1.1	1.0	1.1	1.4	1.2	0.9
208	hemorrhoids	i	1.4	1.8	1.8	1.7	1.3	1.6	2.0
		p	4.0	3.6	3.8	3.1	3.9	3.6	3.9
209	dis. circulatory system NEC	i	0.4	0.6	0.7	0.3	0.5	0.5	0.6
		p	1.9	1.8	2.0	1.0	2.4	1.7	1.9

K CIRCULATORY

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
198	i	0.5	1.2	0.6	0.5	0.7	0.6
	p	1.0	1.9	0.9	0.7	1.1	1.0
199	i	0.5	0.6	0.7	0.6	0.7	0.5
	p	1.2	2.1	1.9	1.4	1.8	1.3
200	i	0.5	1.1	0.5	0.7	0.6	0.5
	p	2.7	4.1	3.2	2.7	3.3	2.3
201	i	0.1	0.2	0.1	0.0	0.1	0.1
	p	0.8	1.0	1.0	0.8	1.1	0.5
202	i	0.6	0.6	0.5	0.6	0.7	0.4
	p	2.3	3.3	3.3	3.5	3.9	1.9
203	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	0.3	0.3	0.5	0.3	0.4	0.3
204	i	0.4	0.5	0.6	0.3	0.5	0.3
	p	0.6	0.8	0.8	0.6	0.8	0.6
205	i	0.3	0.2	0.2	0.1	0.2	0.2
	p	0.8	1.0	1.4	1.1	1.3	0.6
206	i	1.4	1.3	1.2	1.2	1.4	1.0
	p	3.4	3.9	3.9	3.2	4.2	2.4
207	i	0.2	0.3	0.4	0.3	0.3	0.1
	p	0.9	1.1	1.8	0.8	1.3	0.7
208	i	1.8	2.0	1.6	1.3	1.7	1.7
	p	3.8	3.9	3.6	3.6	3.9	3.5
209	i	0.5	0.4	0.8	0.5	0.5	0.5
	p	1.2	1.6	2.7	1.8	1.9	1.4

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
210	neck symp./complt., excl. headache	i	0.7	0.8	0.6	1.0	1.0	0.7	0.8
		p	1.2	1.2	1.3	1.5	1.8	1.2	1.2
211	back symp./complt.	i	1.4	1.4	1.2	1.3	1.6	1.2	1.5
		p	4.2	3.5	3.1	3.8	4.4	3.6	3.6
212	low back symp./complt. w/o radiation	i	0.2	0.5	0.4	0.1	0.2	0.4	0.3
		p	0.3	0.8	0.7	0.2	0.3	0.7	0.4
213	chest sympt./complt.	i	0.8	0.8	1.3	0.5	0.9	0.7	1.1
		p	1.2	1.0	1.6	0.7	1.2	1.0	1.6
214	flank sympt.complt.	i	0.1	0.1	0.1	.	0.0	0.1	0.1
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
215	axilla sympt./complt.	i	0.0	0.0	0.0	0.1	.	0.0	0.0
		p	0.0	0.0	0.0	0.1	.	0.0	0.0
216	jaw sympt./complt.	i	0.4	0.3	0.3	0.1	0.3	0.3	0.3
		p	0.7	0.4	0.3	0.3	0.4	0.5	0.5
217	shoulder sympt./complt.	i	0.4	0.3	0.3	0.2	0.3	0.3	0.5
		p	0.8	0.6	0.7	0.5	0.8	0.6	0.8
218	arm sympt./complt.	i	0.3	0.7	0.6	0.5	0.4	0.4	0.8
		p	0.6	1.2	0.9	0.8	0.6	0.8	1.5
219	elbow sympt./complt.	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.2	0.1	0.2	.	0.2	0.1	0.2
220	wrist sympt./complt.	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.2	0.1	0.2	.	0.2	0.1	0.3
221	hand/finger sympt./complt.	i	0.4	0.6	0.5	0.2	0.6	0.4	0.5
		p	0.7	0.9	0.7	0.4	0.8	0.7	0.8
222	hip sympt./complt.	i	0.2	0.2	0.2	0.2	0.2	0.2	0.3
		p	0.7	0.6	0.6	0.4	1.0	0.5	0.6
223	leg/thigh sympt./complt.	i	0.6	0.6	0.6	0.2	0.7	0.5	0.7
		p	1.5	1.4	1.3	1.0	1.6	1.2	1.6
224	knee sympt./complt.	i	0.8	0.8	0.7	0.4	1.2	0.6	0.8
		p	1.5	1.3	1.3	0.9	1.7	1.2	1.4
225	ankle sympt./complt.	i	0.2	0.2	0.1	0.3	0.2	0.2	0.1
		p	0.3	0.3	0.3	0.3	0.3	0.3	0.2
226	foot/toe sympt./complt.	i	0.6	0.8	0.7	0.4	0.6	0.7	0.7
		p	1.0	1.3	1.1	0.9	0.9	1.1	1.2
227	muscle pain/fibrositis	i	11.3	16.9	16.3	12.4	10.8	13.3	19.1
		p	15.3	22.2	21.1	17.7	16.5	17.6	24.5
228	(mult.) muscle sympt./complt. NEC	i	1.9	1.6	3.0	1.8	2.4	2.1	1.7
		p	2.7	2.4	4.1	3.1	3.5	3.0	2.7
229	(mult.) joints sympt./complt. NEC	i	0.8	0.5	0.8	0.3	1.1	0.6	0.6
		p	2.3	1.2	1.4	0.7	1.8	1.5	1.6
230	limbs sympt./complt. NOS	i	0.0	0.0	0.1	0.1	.	0.0	0.0
		p	0.0	0.0	0.1	0.1	.	0.1	0.0
231	fear of dis. musculoskeletal syst.	i	0.1	0.1	0.2	.	0.1	0.1	0.1
		p	0.1	0.1	0.3	0.1	0.2	0.1	0.2
232	disability/impairment musc. skel. syst.	i	0.5	1.1	1.0	1.3	1.1	1.0	0.8
		p	1.4	2.2	2.4	2.4	2.1	2.0	1.9
233	sympt./complt. musc. skel. syst.	i	0.1	0.1	0.2	0.0	0.1	0.1	0.2
		p	0.2	0.3	0.4	0.1	0.2	0.3	0.3
234	infections musc. skel. syst.	i	0.1	0.2	0.1	0.3	0.1	0.2	0.2
		p	0.2	0.3	0.3	0.4	0.2	0.3	0.3

L MUSCULOSKELETAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
210	i	0.4	0.7	0.6	1.3	0.9	0.5
	p	0.9	1.2	1.1	1.9	1.5	0.8
211	i	1.6	1.0	1.4	1.3	1.5	1.0
	p	3.4	3.5	3.4	4.4	4.3	2.5
212	i	0.9	0.2	0.2	0.2	0.4	0.3
	p	1.3	0.4	0.3	0.4	0.7	0.4
213	i	0.8	0.8	0.9	1.0	0.9	0.8
	p	1.3	0.9	1.2	1.2	1.2	0.9
214	i	0.1	0.0	0.1	0.1	0.1	0.0
	p	0.1	0.0	0.2	0.2	0.2	0.1
215	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
216	i	0.3	0.4	0.2	0.3	0.4	0.2
	p	0.4	0.7	0.4	0.4	0.6	0.3
217	i	0.3	0.3	0.3	0.3	0.4	0.2
	p	0.7	0.7	0.8	0.5	0.8	0.5
218	i	0.5	0.4	0.6	0.5	0.6	0.4
	p	0.8	0.6	1.1	1.1	1.1	0.6
219	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.1	0.1	0.2	0.2	0.2	0.1
220	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.2	0.1	0.1	0.2	0.1
221	i	0.6	0.3	0.5	0.5	0.5	0.5
	p	0.8	0.6	0.9	0.7	0.8	0.7
222	i	0.3	0.2	0.2	0.1	0.2	0.1
	p	0.7	0.7	0.5	0.4	0.7	0.4
223	i	0.5	0.5	0.6	0.6	0.7	0.4
	p	1.4	1.4	1.3	1.4	1.6	0.8
224	i	0.8	0.7	0.7	0.8	0.8	0.5
	p	1.3	1.4	1.4	1.2	1.5	0.8
225	i	0.2	0.1	0.2	0.1	0.2	0.1
	p	0.3	0.3	0.4	0.2	0.3	0.2
226	i	0.7	0.5	0.7	0.9	0.7	0.6
	p	1.0	0.9	1.1	1.5	1.2	0.8
227	i	15.5	9.6	16.1	16.5	16.7	10.0
	p	21.1	13.9	20.4	21.4	22.6	12.7
228	i	1.0	3.0	2.3	1.8	2.2	1.7
	p	2.0	4.3	3.1	2.5	3.5	2.1
229	i	0.4	0.4	0.6	1.2	0.7	0.6
	p	0.9	1.7	1.1	2.5	1.7	1.4
230	i	0.1	0.0	.	0.0	0.0	0.1
	p	0.1	0.0	0.0	0.0	0.0	0.1
231	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.2	0.0	0.2	0.2	0.1	0.1
232	i	0.6	0.8	0.9	1.4	0.9	0.9
	p	1.4	1.9	2.1	2.6	2.2	1.6
233	i	0.0	0.1	0.1	0.3	0.1	0.1
	p	0.3	0.1	0.3	0.4	0.3	0.2
234	i	0.2	0.2	0.0	0.3	0.2	0.1
	p	0.3	0.3	0.1	0.4	0.3	0.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
235	neoplasms musc. skel. syst.	i	0.0	0.2	0.1	0.2	0.1	0.1	0.1
		p	0.2	0.4	0.2	0.3	0.2	0.3	0.3
236	fracture: radius/ulna	i	0.4	0.4	0.4	0.3	0.5	0.3	0.4
		p	0.7	0.5	0.5	0.5	0.7	0.5	0.6
237	fracture: tibia/fibula	i	0.2	0.2	0.2	0.1	0.2	0.2	0.2
		p	0.4	0.4	0.5	0.3	0.3	0.4	0.5
238	fracture: hand/foot bones	i	0.9	0.8	0.7	0.8	0.8	0.8	0.9
		p	1.1	0.9	0.9	1.1	1.0	1.0	1.1
239	fracture: collum femoris	i	0.1	0.1	0.2	0.2	0.1	0.1	0.1
		p	0.2	0.2	0.3	0.4	0.3	0.2	0.3
240	fracture: femur NEC	i	0.1	0.1	0.0	0.0	0.0	0.1	0.1
		p	0.1	0.2	0.1	0.0	0.1	0.1	0.1
241	fracture: skull	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.1
242	fracture: facial bones	i	0.1	0.1	0.1	0.2	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.2	0.1	0.1	0.2
243	fracture: spine	i	0.1	0.0	0.1	0.1	0.0	0.1	0.1
		p	0.3	0.1	0.2	0.3	0.2	0.2	0.2
244	fracture: rib	i	0.1	0.1	0.3	0.1	0.1	0.2	0.1
		p	0.1	0.1	0.4	0.1	0.1	0.2	0.2
245	fracture: clavícula	i	0.1	0.2	0.0	0.1	0.1	0.1	0.1
		p	0.2	0.3	0.1	0.1	0.1	0.2	0.2
246	fracture: humerus	i	0.1	0.1	0.0	0.1	0.1	0.1	0.1
		p	0.2	0.1	0.1	0.3	0.2	0.2	0.1
247	fracture: pelvis	i	0.0	0.0	0.1	0.1	0.0	0.0	0.0
		p	0.0	0.1	0.1	0.1	0.0	0.1	0.0
248	fracture: NEC	i	0.8	0.6	0.3	0.5	0.7	0.6	0.4
		p	1.0	0.8	0.6	0.6	1.2	0.8	0.7
249	sprains/strains ankle(s)	i	3.2	3.5	3.6	1.8	3.4	2.9	4.0
		p	3.8	4.0	4.4	2.5	4.1	3.5	4.9
250	sprains/strains knee(s)	i	2.3	2.5	2.2	1.6	2.2	2.2	2.7
		p	2.5	3.1	2.6	1.9	2.6	2.5	3.2
251	sprains/strains shoulder/arm	i	1.4	1.7	1.4	0.9	1.7	1.3	1.7
		p	1.6	1.9	1.8	1.1	1.9	1.5	2.0
252	sprains/strains wrist/hand	i	2.8	3.3	2.4	2.0	2.2	2.7	3.3
		p	3.0	3.6	3.1	2.0	2.4	3.0	3.8
253	sprains/strains foot/toe	i	2.4	2.5	2.2	0.9	1.9	2.2	2.5
		p	2.7	2.7	2.5	1.0	2.1	2.4	2.7
254	sprains/strains spine, excl. cerv. spine	i	0.7	0.8	0.7	0.6	0.6	0.8	0.8
		p	0.9	0.9	0.9	0.9	0.7	0.9	1.0
255	sprains/strains cerv. spine	i	0.3	0.3	0.4	0.5	0.2	0.4	0.3
		p	0.5	0.4	0.6	0.6	0.3	0.5	0.4
256	whiplash calf	i	0.4	0.4	0.4	0.1	0.3	0.3	0.5
		p	0.4	0.5	0.5	0.1	0.3	0.4	0.6
257	sprains/strains musc. skel. syst. NEC	i	4.0	3.6	3.4	3.0	3.2	3.5	4.1
		p	4.6	4.0	4.1	3.8	3.7	4.0	4.8
258	(sub)luxation, dislocation	i	0.4	0.5	0.6	0.5	0.6	0.5	0.5
		p	0.6	0.6	0.7	0.8	0.8	0.6	0.7
259	cong. luxation of hip	i	0.1	0.1	0.1	0.1	0.2	0.1	0.1
		p	0.3	0.2	0.2	0.1	0.5	0.2	0.2

L MUSCULOSKELETAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
235	i	0.3	0.0	0.1	0.0	0.1	0.0
	p	0.4	0.2	0.3	0.2	0.4	0.2
236	i	0.4	0.3	0.5	0.3	0.4	0.4
	p	0.5	0.6	0.6	0.5	0.6	0.5
237	i	0.3	0.2	0.3	0.2	0.2	0.3
	p	0.5	0.4	0.5	0.3	0.4	0.4
238	i	0.8	1.2	0.7	0.7	0.7	0.8
	p	1.0	1.4	0.8	0.8	0.9	0.9
239	i	0.1	0.2	0.1	0.1	0.2	0.1
	p	0.2	0.4	0.3	0.2	0.3	0.2
240	i	0.0	0.2	0.1	0.0	0.1	0.1
	p	0.1	0.3	0.1	0.1	0.1	0.1
241	i	.	0.0	0.0	0.0	0.0	.
	p	0.0	0.0	0.0	0.0	0.0	0.0
242	i	0.1	0.0	0.2	0.0	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.1
243	i	0.2	0.1	0.0	0.1	0.1	0.1
	p	0.3	0.2	0.1	0.2	0.2	0.1
244	i	0.1	0.2	0.2	0.2	0.2	0.1
	p	0.1	0.2	0.2	0.2	0.2	0.2
245	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.2	0.1	0.2	0.2	0.1
246	i	0.0	0.1	0.1	0.0	0.0	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.1
247	i	0.0	0.0	0.1	.	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.0
248	i	0.4	0.8	0.5	0.5	0.6	0.6
	p	0.7	1.0	0.7	0.8	0.8	0.7
249	i	3.6	2.5	3.4	3.3	3.1	2.9
	p	4.4	3.1	4.2	3.9	3.9	3.4
250	i	2.2	2.0	2.1	2.8	2.4	1.9
	p	2.5	2.3	2.4	3.4	2.8	2.3
251	i	1.2	1.2	1.5	1.8	1.6	1.0
	p	1.4	1.5	1.7	2.1	1.9	1.2
252	i	2.6	2.6	3.0	3.0	3.0	2.1
	p	3.0	2.8	3.4	3.2	3.5	2.2
253	i	1.7	2.2	2.2	2.8	2.5	1.6
	p	1.9	2.4	2.4	3.0	2.7	1.7
254	i	0.6	1.0	0.8	0.6	0.9	0.5
	p	0.7	1.2	0.9	0.8	1.1	0.6
255	i	0.4	0.3	0.4	0.3	0.3	0.3
	p	0.5	0.6	0.5	0.3	0.5	0.4
256	i	0.4	0.3	0.3	0.5	0.4	0.4
	p	0.4	0.5	0.3	0.5	0.4	0.5
257	i	3.7	3.7	3.8	3.4	3.9	2.8
	p	4.3	4.2	4.4	3.8	4.5	3.1
258	i	0.5	0.5	0.5	0.5	0.5	0.4
	p	0.7	0.5	0.7	0.8	0.7	0.5
259	i	0.2	0.0	0.1	0.0	0.1	0.1
	p	0.4	0.2	0.2	0.1	0.2	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
260	cong. anom. musc. skel. syst. NEC	i	0.2	0.2	0.0	0.1	0.1	0.1	0.2
		p	0.5	0.3	0.3	0.4	0.3	0.4	0.4
261	syndromes cervical spine	i	1.5	1.0	1.2	1.9	1.3	1.2	1.6
		p	2.1	1.9	2.0	2.7	1.9	2.0	2.4
262	osteoarthritis of cervical spine	i	0.6	0.6	1.3	0.8	0.7	0.9	0.7
		p	1.4	1.2	2.5	1.6	1.4	1.7	1.3
263	osteoarthritis of spine NEC	i	0.6	0.7	0.6	0.4	0.5	0.6	0.6
		p	1.7	1.5	1.2	1.4	1.3	1.5	1.8
264	acquired deformities of spine	i	0.4	0.9	0.7	0.8	0.7	0.7	0.8
		p	1.2	1.8	1.6	1.8	1.7	1.5	1.6
265	back pain with radiating sympt.	i	0.4	0.5	0.4	0.5	0.5	0.5	0.4
		p	0.7	1.0	0.9	0.5	0.9	0.9	0.8
266	lumbago	i	6.7	6.8	7.1	4.3	5.8	6.2	7.9
		p	8.5	9.1	9.9	6.4	7.2	8.4	10.4
267	sciatica	i	1.0	1.0	1.1	1.4	1.5	1.1	0.8
		p	1.7	1.6	1.6	2.1	2.4	1.7	1.3
268	slipped lumbar disc	i	1.5	1.2	1.3	0.7	0.7	1.2	1.8
		p	3.4	3.0	2.9	3.0	2.3	2.9	4.0
269	ganglion joint/tendon	i	0.6	0.7	0.8	0.6	0.8	0.6	0.9
		p	0.7	0.8	0.9	0.8	0.8	0.7	1.1
270	rheumatoid arthritis	i	0.5	0.5	0.8	0.4	0.5	0.5	0.7
		p	4.2	2.2	2.7	3.0	3.2	2.9	3.3
271	spondylitis (ankylosing)	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.4	0.4	0.4	0.2	0.4	0.3	0.5
272	osteoarthritis hip	i	0.7	0.6	0.8	0.6	0.5	0.7	0.8
		p	2.0	1.6	2.1	1.7	1.6	1.7	2.3
273	osteoarthritis knee	i	1.1	0.9	1.2	1.6	1.3	1.0	1.3
		p	2.3	2.0	3.0	4.0	2.4	2.5	2.7
274	osteoarthritis NEC	i	0.8	0.9	1.4	1.6	0.8	1.1	1.0
		p	3.1	2.7	4.0	3.4	2.6	3.2	3.5
275	arthritis in allied conditions NEC	i	1.0	1.0	1.4	0.9	1.1	1.0	1.3
		p	1.9	1.8	3.0	2.2	2.3	2.0	2.4
276	shoulder syndrome	i	2.5	2.8	4.2	2.5	2.4	3.0	3.2
		p	3.6	4.2	5.9	3.5	3.4	4.3	4.8
277	dis. bursa, excl. shoulder	i	1.5	1.0	1.9	1.2	1.3	1.3	1.6
		p	1.9	1.3	2.1	1.5	1.7	1.6	1.9
278	epicondylitis lateralis	i	1.5	2.0	1.8	1.3	1.6	1.6	2.1
		p	2.1	2.7	2.4	1.9	2.0	2.2	2.8
279	tendinitis/synovitis	i	3.5	5.0	5.4	2.3	4.0	4.4	4.4
		p	4.2	6.1	6.8	3.3	4.7	5.4	5.5
280	osteochondrosis	i	0.3	0.3	0.2	0.2	0.4	0.2	0.3
		p	0.6	0.5	0.3	0.5	0.7	0.4	0.5
281	osteoporosis	i	0.2	0.2	0.3	0.2	0.2	0.2	0.1
		p	1.6	1.0	1.6	2.1	1.3	1.6	1.1
282	ac. traum. meniscus/lig. knee	i	1.3	1.2	1.1	1.3	1.2	1.2	1.2
		p	1.4	1.4	1.3	1.4	1.4	1.3	1.5
283	retropatellar chondropathy	i	0.8	1.0	1.2	1.1	0.9	1.0	1.0
		p	1.1	1.2	1.7	1.3	1.2	1.3	1.4
284	chron. internal knee derangement	i	0.2	0.3	0.3	0.3	0.2	0.2	0.5
		p	0.3	0.5	0.6	0.4	0.3	0.4	0.6

L MUSCULOSKELETAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
260	i	0.1	0.1	0.2	0.2	0.2	0.1
	p	0.4	0.4	0.4	0.4	0.4	0.3
261	i	0.8	1.7	1.6	1.1	1.5	1.0
	p	1.5	2.3	2.3	2.1	2.2	1.7
262	i	0.7	1.0	0.9	0.6	0.9	0.6
	p	1.7	1.5	1.9	1.3	1.9	1.1
263	i	0.7	0.5	0.8	0.3	0.7	0.5
	p	1.8	1.3	1.9	1.1	1.7	1.2
264	i	0.7	0.6	0.5	0.9	0.7	0.7
	p	1.5	1.3	1.6	1.8	1.6	1.4
265	i	0.5	0.3	0.5	0.5	0.5	0.4
	p	0.8	0.8	0.9	0.9	1.0	0.6
266	i	5.7	6.4	6.2	8.0	7.5	4.8
	p	7.7	8.3	8.5	10.7	10.3	6.1
267	i	1.0	1.1	1.4	0.7	1.1	0.9
	p	1.9	1.7	1.9	1.3	1.9	1.3
268	i	0.9	1.5	1.4	1.3	1.4	1.1
	p	2.6	2.9	3.3	3.7	3.3	2.6
269	i	0.5	0.6	1.1	0.5	0.7	0.6
	p	0.7	0.7	1.3	0.6	0.8	0.7
270	i	0.5	0.7	0.4	0.5	0.6	0.4
	p	3.1	4.0	2.6	2.5	3.5	2.3
271	i	0.1	0.0	0.1	0.2	0.1	0.0
	p	0.4	0.2	0.4	0.4	0.4	0.2
272	i	0.6	0.8	0.7	0.6	0.8	0.4
	p	1.6	1.9	1.9	1.9	2.1	1.4
273	i	1.2	1.0	1.0	1.2	1.3	0.7
	p	2.6	2.3	2.4	2.8	2.9	1.6
274	i	0.9	1.1	1.3	0.8	1.2	0.8
	p	3.1	2.9	4.3	2.5	3.6	2.4
275	i	1.3	1.1	1.2	0.8	1.2	0.8
	p	2.3	2.3	2.6	1.6	2.5	1.5
276	i	2.3	2.8	3.1	3.8	3.3	2.3
	p	3.5	3.9	4.6	5.3	4.7	3.6
277	i	1.5	1.7	1.3	1.1	1.4	1.4
	p	1.8	1.9	1.7	1.3	1.7	1.6
278	i	1.7	1.5	1.7	2.0	1.9	1.6
	p	2.2	2.3	2.2	2.6	2.6	2.1
279	i	4.4	4.2	5.1	3.7	4.6	3.7
	p	5.4	4.9	6.7	4.4	5.6	4.6
280	i	0.2	0.1	0.3	0.4	0.2	0.3
	p	0.6	0.3	0.6	0.5	0.5	0.5
281	i	0.2	0.3	0.2	0.1	0.2	0.2
	p	1.7	1.4	1.4	1.3	1.6	1.2
282	i	0.9	1.6	1.1	1.3	1.2	1.1
	p	1.1	1.7	1.2	1.5	1.5	1.2
283	i	0.7	0.5	1.4	1.2	1.1	0.7
	p	1.0	0.9	1.8	1.6	1.5	0.9
284	i	0.3	0.2	0.3	0.3	0.3	0.2
	p	0.5	0.2	0.5	0.5	0.5	0.3

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
285	acquired deformities of limbs	i	1.3	1.7	2.1	2.3	1.3	1.8	1.6
		p	1.9	2.5	3.2	3.5	1.9	2.6	2.7
286	dis. musc. skel. system NEC	i	1.3	1.5	1.6	1.0	1.4	1.3	1.6
		p	2.1	2.3	2.4	1.8	2.2	2.1	2.4

L MUSCULOSKELETAL

no.		season	summer	autumn	winter	type of insurance	
		spring				public	private
285	i	2.0	1.6	1.9	1.4	1.7	1.7
	p	2.9	2.3	2.9	2.3	2.7	2.2
286	i	1.0	1.3	2.0	1.3	1.4	1.2
	p	1.6	2.3	3.0	1.9	2.4	1.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
287	headache/facial pain	i	1.8	1.6	1.4	1.1	1.4	1.4	2.0
		p	6.8	4.8	4.9	3.2	5.0	4.9	6.3
288	sensibility disturbance	i	0.2	0.4	0.5	0.4	0.3	0.4	0.4
		p	0.7	0.7	1.0	0.7	0.6	0.8	0.8
289	restless legs syndrome	i	0.3	0.3	0.5	0.4	0.3	0.3	0.5
		p	1.2	0.8	1.2	0.8	1.3	0.9	1.1
290	convulsions	i	0.1	0.1	0.3	0.1	0.0	0.1	0.1
		p	0.2	0.1	0.3	0.2	0.1	0.2	0.2
291	symptoms sense/smell/taste	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.0	.	.	0.0	0.0	0.0
292	vertigo	i	1.9	1.6	1.2	1.5	1.5	1.6	1.6
		p	5.3	3.8	3.3	3.6	4.9	4.2	4.0
293	paralysis	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.6	0.5	0.3	0.3	0.5	0.4	0.6
294	sympt./complt. speech	i	0.1	0.2	0.3	0.1	0.1	0.2	0.2
		p	0.3	0.5	0.5	0.2	0.2	0.4	0.4
295	fear of neurological disease	i	0.0	0.1	0.1	.	0.0	0.0	0.0
		p	0.1	0.1	0.2	.	0.0	0.1	0.1
296	disability/impairment neurol. syst.	i	.	.	0.0	.	0.0	.	.
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
297	sympt./complt. neurol. syst. NEC	i	0.1	0.1	0.0	0.1	0.0	0.1	0.1
		p	0.2	0.2	0.2	0.1	0.2	0.2	0.2
298	poliomyelitis	i
		p	0.1	0.1	0.1	0.1	0.1	0.1	0.1
299	meningitis/encefalitis	i	0.0	0.0	0.0	0.1	0.1	0.0	0.0
		p	0.1	0.1	0.1	0.1	0.2	0.1	0.1
300	tetanus	i
		p	.	0.0	.	.	0.0	.	.
301	infections neur. syst. NEC	i	.	0.0	0.0	.	0.0	0.0	.
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
302	malignant neoplasm neur. syst.	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.2	0.1	0.2	0.2	0.2	0.1	0.3
303	benigne neopl. neur. syst.	i	0.0	0.0	.	0.2	0.0	0.0	0.1
		p	0.1	0.1	0.1	0.3	0.1	0.1	0.2
304	unspecified neopl. neur. syst.	i	0.0	0.0	0.1	0.1	0.0	0.0	0.0
		p	0.1	0.1	0.1	0.3	0.1	0.1	0.1
305	concussion: commotio cer.	i	0.6	0.6	0.5	0.4	0.7	0.5	0.7
		p	0.7	0.7	0.6	0.7	0.8	0.6	0.8
306	concussion: contusio cer.	i	0.1	0.1	0.1	0.1	0.2	0.1	0.1
		p	0.2	0.1	0.1	0.1	0.3	0.1	0.1
307	head injury NOS/NEC, w/o skull fract.	i	0.1	0.2	0.1	.	0.0	0.1	0.1
		p	0.1	0.2	0.1	.	0.1	0.1	0.1
308	injuries neur. syst. NEC	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.1	0.1	0.1	0.1	0.1	0.1	0.2
309	cong. anom. neur. syst.	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.3	0.2	0.3	0.2	0.2	0.2	0.3
310	multiple sclerosis	i	0.1	0.1	0.0	.	0.1	0.1	0.0
		p	0.5	0.6	0.4	0.2	0.6	0.5	0.3
311	Parkinsonism	i	0.1	0.1	0.1	0.3	0.2	0.1	0.1
		p	1.5	1.1	1.3	1.7	1.8	1.2	1.5

N NEUROLOGICAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
287	i	1.8	1.2	1.6	1.7	1.7	1.0
	p	5.6	4.5	5.4	5.6	6.1	3.4
288	i	0.5	0.2	0.4	0.3	0.3	0.3
	p	0.8	0.7	1.0	0.5	0.8	0.6
289	i	0.4	0.4	0.5	0.2	0.5	0.2
	p	0.8	1.1	1.3	0.7	1.2	0.7
290	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.2	0.1	0.3	0.1	0.2	0.2
291	i	.	.	0.0	0.0	0.0	0.0
	p	.	.	0.0	0.0	0.0	0.0
292	i	1.3	1.5	1.7	1.8	1.8	1.2
	p	3.6	4.4	4.4	4.4	5.0	2.8
293	i	0.1	0.2	0.1	0.1	0.1	0.1
	p	0.5	0.5	0.4	0.6	0.5	0.4
294	i	0.1	0.2	0.4	0.2	0.2	0.1
	p	0.3	0.3	0.7	0.4	0.5	0.2
295	i	0.1	0.0	.	0.1	0.0	0.1
	p	0.1	0.1	0.1	0.1	0.1	0.2
296	i	.	.	.	0.0	0.0	.
	p	.	0.0	.	0.1	0.0	0.0
297	i	0.1	0.1	0.0	0.1	0.0	0.1
	p	0.2	0.2	0.1	0.1	0.2	0.2
298	i
	p	0.1	0.1	0.1	0.1	0.1	0.1
299	i	0.0	0.1	0.0	0.1	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.1
300	i
	p	.	0.0	.	.	0.0	.
301	i	.	0.0	0.0	.	0.0	.
	p	0.0	0.0	0.0	0.0	0.0	0.0
302	i	0.0	0.0	0.1	0.0	0.0	0.0
	p	0.2	0.1	0.2	0.1	0.1	0.2
303	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.2	0.1	0.1	0.1
304	i	0.1	0.1	0.0	0.0	0.0	0.1
	p	0.1	0.2	0.1	0.0	0.1	0.1
305	i	0.5	0.4	0.6	0.6	0.5	0.6
	p	0.6	0.6	0.8	0.8	0.7	0.7
306	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.1	0.2	0.2	0.1	0.2	0.1
307	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	0.1	0.1	0.1	0.0	0.1	0.1
308	i	0.0	0.1	0.1	0.0	0.0	0.1
	p	0.1	0.1	0.1	0.1	0.1	0.1
309	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.4	0.2	0.2	0.2	0.3	0.2
310	i	0.1	0.1	0.1	0.0	0.0	0.1
	p	0.3	0.5	0.5	0.5	0.5	0.4
311	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.9	1.4	1.6	1.5	1.6	1.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
312	epilepsy, all types	i	0.3	0.3	0.3	0.3	0.2	0.3	0.4
		p	5.1	2.7	2.9	2.2	5.5	3.3	3.1
313	migraine	i	1.3	1.4	1.3	0.6	1.0	1.2	1.4
		p	5.0	4.3	3.8	2.6	4.9	4.1	4.3
314	facial paralysis, Bell's palsy	i	0.1	0.1	0.1	0.2	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.3	0.1	0.1	0.2
315	trigeminal neuralgia	i	0.1	0.1	0.2	0.1	0.1	0.1	0.1
		p	0.2	0.2	0.4	0.2	0.3	0.3	0.3
316	carpal tunnel syndrome	i	0.2	0.3	0.2	0.3	0.3	0.3	0.3
		p	0.4	0.5	0.4	0.5	0.6	0.4	0.6
317	cluster headache	i	0.0	0.0	0.0	0.1	0.1	0.0	0.0
		p	0.1	0.1	0.0	0.1	0.1	0.1	0.0
318	dis. of periph. nerv. syst. NEC	i	2.3	2.3	2.2	1.7	2.6	2.2	1.9
		p	4.3	3.8	4.0	2.8	4.6	3.9	3.5
319	dis. of central nerv. syst. NEC	i	0.2	0.2	0.3	0.4	0.2	0.2	0.3
		p	0.8	0.8	1.6	0.7	1.3	0.9	1.2

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
312	i	0.3	0.2	0.4	0.3	0.3	0.2
	p	3.0	3.7	3.2	4.0	3.9	2.4
313	i	1.1	1.4	1.5	0.9	1.4	1.0
	p	3.9	5.3	4.3	3.5	4.5	3.7
314	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.1	0.1	0.2	0.1	0.1	0.1
315	i	0.2	0.2	0.1	0.0	0.2	0.0
	p	0.4	0.3	0.3	0.1	0.3	0.2
316	i	0.1	0.2	0.5	0.3	0.3	0.2
	p	0.3	0.3	0.8	0.4	0.5	0.3
<hr/>							
317	i	0.0	0.0	0.0	0.1	0.0	0.0
	p	0.1	0.0	0.1	0.1	0.1	0.0
318	i	1.7	1.9	2.2	2.9	2.3	1.9
	p	2.9	3.8	4.0	4.8	4.3	3.1
319	i	0.2	0.4	0.3	0.2	0.3	0.3
	p	0.8	0.9	1.0	1.2	1.1	0.7

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
320	feeling anxious/nervous/tense	i	2.9	4.5	4.6	3.7	2.8	3.8	4.5
		p	22.9	21.5	20.2	15.4	21.7	21.0	20.4
321	acute stress (transient) reaction	i	1.1	1.5	2.8	2.7	1.4	2.0	1.5
		p	3.0	3.6	5.1	11.1	4.0	5.0	3.8
322	feeling depressed	i	0.2	0.4	0.4	0.6	0.4	0.4	0.4
		p	2.4	1.3	0.9	2.1	2.1	1.7	1.2
323	feeling/behaving irritable	i	0.1	0.2	0.3	0.2	0.2	0.2	0.2
		p	1.2	0.8	1.1	0.8	0.9	0.7	1.7
324	suicide attempt	i	0.1	0.1	0.1	0.1	0.2	0.1	0.2
		p	0.2	0.1	0.1	0.2	0.2	0.1	0.2
325	disturbances of sleep	i	0.8	1.3	1.1	0.9	0.8	1.1	1.0
		p	22.7	18.2	19.4	16.9	25.0	20.8	14.6
326	sexual problems	i	0.2	0.4	0.4	0.4	0.1	0.3	0.3
		p	0.3	0.5	0.6	0.7	0.3	0.5	0.5
327	hypochondr. dis./probl. with being ill	i	0.1	0.2	0.2	0.2	0.1	0.2	0.2
		p	0.5	0.8	1.3	1.1	0.7	0.7	1.2
328	tension headache	i	1.4	2.7	2.3	2.3	1.7	2.1	2.5
		p	2.3	4.1	3.5	2.7	2.6	3.0	4.0
329	hyperventilation	i	1.5	2.3	2.1	2.8	1.8	2.2	1.7
		p	2.3	3.6	3.4	3.8	2.8	3.4	2.9
330	enuresis	i	0.2	0.3	0.3	0.2	0.2	0.3	0.3
		p	1.3	0.7	0.9	0.6	1.4	0.9	0.9
331	anorexia nervosa/boulimia	i	0.1	0.1	0.2	0.2	0.0	0.1	0.1
		p	0.3	0.2	0.4	0.4	0.2	0.3	0.3
332	eating/feeding probl. NEC	i	0.0	0.1	0.0	0.1	0.0	0.0	0.0
		p	0.0	0.1	0.3	0.1	0.1	0.1	0.1
333	chronic alcohol abuse	i	0.2	0.4	0.4	0.6	0.4	0.4	0.3
		p	1.3	1.6	2.1	3.0	1.3	1.9	1.6
334	acute alcohol abuse	i	0.1	0.0	0.0	0.1	0.1	0.0	0.1
		p	0.1	0.1	0.1	0.1	0.2	0.1	0.1
335	tobacco abuse	i	0.5	0.4	0.6	0.5	0.4	0.4	0.6
		p	1.8	1.6	3.3	2.3	1.7	2.0	2.5
336	drug abuse	i	0.0	0.0	0.1	0.1	0.0	0.1	0.0
		p	0.2	0.1	0.6	1.1	0.2	0.5	0.2
337	medicinal abuse	i	0.0	0.0	.	0.1	0.0	0.0	0.0
		p	0.2	0.5	0.4	0.7	0.5	0.4	0.5
338	abuse substances NEC	i	.	0.0	0.0
		p	0.1	0.0	0.0	0.1	.	0.0	0.1
339	dist. memory/concentration	i	0.1	0.1	0.1	0.3	0.2	0.1	0.2
		p	0.4	0.6	0.5	0.6	0.7	0.5	0.4
340	disturbance behaviour child	i	0.2	0.1	0.2	0.1	0.2	0.2	0.2
		p	0.2	0.3	0.3	0.2	0.4	0.2	0.3
341	sympt./complt. behaviour adolescent	i	0.0	0.1	0.2	0.1	0.1	0.0	0.2
		p	0.1	0.1	0.3	0.4	0.1	0.2	0.3
342	sympt./complt. adult	i	0.1	0.1	0.1	0.4	0.1	0.2	0.2
		p	0.5	0.7	0.6	0.9	0.6	0.6	0.6
343	phase of live problem adult	i	0.0	0.1	0.1	0.0	0.1	0.0	0.0
		p	0.1	0.3	0.3	0.3	0.5	0.2	0.3
344	dist. learning/mental development	i	0.0	0.1	0.1	.	0.0	0.0	0.1
		p	0.2	0.3	0.2	0.2	0.2	0.2	0.2

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
320	i	3.7	3.9	4.0	3.9	4.3	3.2
	p	19.3	22.3	19.9	22.4	23.8	15.7
321	i	1.7	1.3	1.7	2.6	1.9	1.4
	p	3.9	2.9	3.8	7.7	5.0	3.5
322	i	0.6	0.3	0.3	0.4	0.4	0.3
	p	1.8	1.8	1.7	1.4	1.8	1.4
323	i	0.1	0.2	0.3	0.2	0.2	0.2
	p	0.7	1.6	1.0	0.8	1.1	0.8
324	i	0.2	0.0	0.2	0.1	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.1
325	i	1.0	1.4	1.0	0.8	1.1	1.0
	p	17.0	23.5	19.5	19.0	22.3	15.4
326	i	0.3	0.2	0.4	0.3	0.3	0.3
	p	0.5	0.3	0.6	0.5	0.5	0.5
327	i	0.2	0.1	0.1	0.3	0.2	0.1
	p	1.2	0.4	0.7	1.1	1.0	0.6
328	i	2.1	1.7	2.7	2.1	2.6	1.4
	p	3.3	2.3	4.0	3.2	3.7	2.2
329	i	2.0	1.6	2.7	2.0	2.3	1.4
	p	3.3	2.3	3.9	3.2	3.6	2.1
330	i	0.3	0.2	0.2	0.3	0.3	0.2
	p	1.1	0.7	0.8	1.1	1.1	0.6
331	i	0.1	0.2	0.1	0.2	0.1	0.2
	p	0.3	0.3	0.3	0.3	0.3	0.3
332	i	0.1	0.1	0.0	0.0	0.1	0.0
	p	0.2	0.1	0.1	0.1	0.1	0.1
333	i	0.4	0.3	0.5	0.3	0.4	0.3
	p	1.6	1.6	2.2	1.7	2.2	0.9
334	i	0.0	0.1	0.1	0.1	0.1	0.0
	p	0.0	0.1	0.1	0.1	0.1	0.0
335	i	0.4	0.4	0.5	0.6	0.6	0.3
	p	1.4	1.4	2.9	2.9	2.7	1.2
336	i	0.0	0.0	0.1	0.1	0.1	0.0
	p	0.4	0.3	0.5	0.2	0.4	0.1
337	i	0.0	0.0	0.0	0.1	0.0	0.0
	p	0.5	0.3	0.6	0.4	0.5	0.3
338	i	.	.	0.0	.	0.0	.
	p	0.0	0.1	0.0	0.0	0.1	.
339	i	0.1	0.1	0.1	0.2	0.1	0.0
	p	0.3	0.6	0.5	0.6	0.5	0.4
340	i	0.1	0.1	0.2	0.2	0.2	0.1
	p	0.2	0.2	0.3	0.3	0.3	0.2
341	i	0.0	0.0	0.2	0.1	0.1	0.1
	p	0.0	0.2	0.3	0.3	0.2	0.2
342	i	0.2	0.1	0.1	0.2	0.2	0.1
	p	0.5	0.6	0.7	0.6	0.7	0.4
343	i	0.0	0.0	0.1	0.0	0.0	0.1
	p	0.1	0.3	0.3	0.3	0.3	0.2
344	i	0.1	.	0.0	0.0	0.0	0.1
	p	0.2	0.1	0.3	0.3	0.2	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
345	fear of mental disorder	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.1	0.0	.	0.0	0.0	0.0
346	psychic disability/impairment	i	.	0.0	0.0
		p	0.0	0.0	0.1	.	0.0	0.0	0.1
347	psychic sympt./compit. NEC	i	0.4	0.4	1.0	0.2	1.0	0.4	0.5
		p	1.4	1.0	2.1	3.3	1.9	1.6	1.7
348	dementia (pre)senile	i	0.2	0.2	0.3	0.2	0.1	0.2	0.3
		p	0.9	0.9	1.4	1.5	1.2	1.0	1.2
349	organic psychosis	i	0.0	0.0	0.1	.	0.1	0.1	0.0
		p	0.2	0.3	0.3	0.1	0.2	0.2	0.1
350	schizophrenia, all types	i	0.0	0.1	0.0	0.3	0.1	0.1	0.0
		p	0.5	0.5	0.7	0.9	0.6	0.6	0.5
351	affective psychosis	i	0.2	0.1	0.2	0.4	0.2	0.2	0.2
		p	1.1	0.6	0.9	1.2	1.0	1.0	0.8
352	unspecified psychosis NEC	i	0.1	0.1	0.2	0.2	0.2	0.1	0.1
		p	1.2	0.9	1.2	1.7	1.5	1.1	1.1
353	anxiety disorder/anxiety state	i	0.1	0.2	0.3	0.4	0.1	0.3	0.2
		p	0.6	0.7	0.9	0.8	1.0	0.7	0.7
354	hysterical/hypochondriacal dis.	i	0.2	0.3	0.4	0.6	0.4	0.3	0.2
		p	1.2	1.5	1.5	2.0	1.4	1.4	1.5
355	depressive disorder	i	2.2	2.5	2.1	1.9	1.7	2.4	2.0
		p	9.7	8.2	7.4	6.5	8.0	8.6	7.6
356	mental retardation	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.7	0.9	0.7	0.4	1.1	0.7	0.8
357	neurasthenia, surmenage	i	2.5	3.8	4.6	3.4	4.1	3.5	3.1
		p	4.7	6.4	8.4	5.5	7.3	6.1	5.9
358	personality disorder	i	0.0	0.0	0.1	0.1	.	0.0	0.1
		p	0.3	0.2	0.2	0.2	0.2	0.2	0.3
359	phobia/neurotic disorder NEC	i	0.3	0.4	0.7	0.4	0.3	0.4	0.5
		p	1.1	1.7	3.4	1.7	1.9	2.0	1.7
360	mental/psychological dis. NEC	i	0.2	0.7	0.4	0.7	0.7	0.6	0.3
		p	1.2	2.5	2.2	2.7	2.4	2.2	1.4

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
345	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
346	i	.	.	.	0.0	.	0.0
	p	0.0	0.0	0.1	0.0	0.1	0.0
347	i	0.4	0.4	0.3	0.8	0.5	0.4
	p	1.4	1.3	2.2	1.7	1.9	1.2
348	i	0.2	0.2	0.3	0.1	0.2	0.2
	p	0.8	1.3	1.5	0.8	0.9	1.1
349	i	0.0	0.1	0.0	0.0	0.1	0.1
	p	0.2	0.3	0.1	0.3	0.3	0.1
350	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.6	0.5	0.8	0.5	0.7	0.3
351	i	0.2	0.4	0.2	0.1	0.2	0.2
	p	1.0	1.1	0.9	0.6	0.9	0.9
352	i	0.1	0.2	0.2	0.1	0.2	0.1
	p	0.7	1.5	1.2	1.2	1.3	0.7
353	i	0.4	0.3	0.3	0.1	0.3	0.2
	p	0.9	0.7	0.9	0.5	0.9	0.5
354	i	0.3	0.2	0.3	0.3	0.4	0.2
	p	1.5	1.5	1.3	1.4	1.7	1.0
355	i	1.7	2.5	2.7	2.1	2.5	1.8
	p	7.6	9.3	8.9	7.5	9.1	6.9
356	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.7	0.5	1.0	0.9	1.0	0.3
357	i	2.8	3.6	4.7	2.9	3.7	3.3
	p	5.0	6.4	8.1	5.3	6.5	5.9
358	i	.	0.0	0.1	0.1	0.0	0.0
	p	0.0	0.2	0.5	0.2	0.3	0.1
359	i	0.2	0.4	0.4	0.7	0.4	0.4
	p	0.6	1.6	2.5	2.9	2.0	1.7
360	i	0.3	0.2	0.2	0.4	0.5	0.5
	p	2.4	1.9	1.8	1.1	2.2	1.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
361	pain attr. to respiratory system	i	.	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.1	0.0	.	.	0.0	0.0
362	shortness of breath, dyspnoe	i	0.4	0.2	0.2	0.3	0.2	0.3	0.3
		p	1.4	0.6	1.0	0.8	0.9	0.9	1.1
363	wheezing	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.0	0.1	0.1	0.0	0.0	0.0
364	breathing problems NEC	i	0.0	0.1	0.3	0.2	0.0	0.1	0.1
		p	0.1	0.1	0.4	0.2	0.1	0.2	0.1
365	cough	i	9.5	10.2	9.0	6.2	10.3	9.1	9.0
		p	9.5	10.3	9.2	6.5	10.6	9.2	9.4
366	epistaxis/nose bleed	i	0.5	0.6	0.5	0.1	0.4	0.5	0.5
		p	0.6	0.8	0.6	0.1	0.5	0.6	0.6
367	sneezing/nasal congestion	i	0.3	0.2	0.5	0.3	0.2	0.3	0.3
		p	1.0	0.4	1.0	0.5	0.6	0.8	0.7
368	sympt./complt. nose NEC	i	0.2	0.2	0.3	0.3	0.3	0.2	0.3
		p	0.4	0.4	0.4	0.6	0.5	0.4	0.5
369	sympt./complt. sinus (incl. pain)	i	0.1	0.2	0.1	0.0	0.1	0.1	0.2
		p	0.1	0.3	0.1	0.0	0.2	0.1	0.2
370	sympt./complt. throat	i	0.9	1.5	1.5	1.8	1.1	1.2	1.7
		p	2.5	2.5	2.9	2.7	2.1	2.5	3.0
371	sympt./complt. tonsils	i	0.0	0.0	0.1	0.1	0.1	0.0	0.1
		p	0.1	0.1	0.1	0.1	0.2	0.0	0.1
372	sympt./complt. voice/vocal cords	i	0.3	0.3	0.2	0.4	0.2	0.3	0.4
		p	0.5	0.5	0.2	0.4	0.3	0.4	0.5
373	haemoptysis	i	0.0	0.1	0.1	.	0.0	0.0	0.1
		p	0.0	0.1	0.2	.	0.0	0.1	0.1
374	abnormal sputum/phlegm NEC	i	0.1	0.2	0.4	0.2	0.2	0.2	0.3
		p	0.5	0.6	1.1	0.2	0.5	0.5	1.1
375	fear of respiratory disease	i	0.1	0.1	0.3	0.1	0.0	0.1	0.3
		p	0.1	0.3	0.4	0.2	0.1	0.2	0.5
376	disability/impairment resp. syst.	i	0.0	.	.	.	0.0	0.0	.
		p	0.0	0.0	.	.	0.0	0.0	0.0
377	sympt./complt. resp. syst. NEC	i	0.3	0.1	0.6	0.0	0.3	0.2	0.3
		p	0.6	0.2	1.1	0.2	0.9	0.5	0.5
378	tuberculosis resp. (excl. A70)	i	.	0.0	0.1	.	0.0	0.0	0.0
		p	0.2	0.1	0.3	0.2	0.3	0.1	0.2
379	whooping cough	i	0.6	0.2	0.1	0.1	0.3	0.3	0.3
		p	0.9	0.3	0.2	0.2	0.5	0.5	0.4
380	scarlet fever	i	0.2	0.3	0.2	0.3	0.2	0.3	0.1
		p	0.2	0.4	0.2	0.3	0.2	0.4	0.1
381	streptoc. inf. throat NEC	i	0.0	0.1	0.1	.	0.0	0.0	0.1
		p	0.1	0.1	0.1	0.0	0.0	0.1	0.1
382	boil/abscess nose	i	0.0	0.0	0.1	0.1	0.0	0.1	0.1
		p	0.1	0.0	0.1	0.1	0.0	0.1	0.1
383	URI (head cold)	i	27.3	32.4	32.6	32.4	22.9	28.6	40.0
		p	30.3	34.6	35.1	36.3	25.2	31.5	42.5
384	sinusitis acute/chronic	i	8.5	8.5	10.1	7.3	9.0	8.3	9.5
		p	10.4	10.3	12.4	9.5	10.5	10.4	11.7
385	acute tonsillitis/adenoiditis	i	7.4	4.9	5.8	5.7	8.8	6.1	4.6
		p	7.5	5.1	6.0	6.1	8.9	6.3	4.8

R RESPIRATORY

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
361	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.1	0.0	0.0
362	i	0.2	0.4	0.3	0.2	0.3	0.2
	p	0.8	1.5	0.9	0.6	1.1	0.6
363	i	0.0	.	0.0	.	0.0	.
	p	0.1	0.0	0.1	0.0	0.0	0.0
364	i	0.2	0.0	0.1	0.1	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.2	0.1
365	i	8.7	8.0	9.6	10.1	10.3	6.7
	p	8.9	8.3	10.0	10.2	10.7	6.9
366	i	0.5	0.3	0.5	0.5	0.5	0.4
	p	0.6	0.5	0.6	0.6	0.6	0.6
367	i	0.1	0.2	0.4	0.5	0.4	0.2
	p	0.4	0.7	0.8	1.1	0.8	0.6
368	i	0.2	0.2	0.3	0.4	0.3	0.2
	p	0.3	0.3	0.4	0.6	0.4	0.3
369	i	0.2	0.1	0.1	0.1	0.1	0.2
	p	0.3	0.1	0.1	0.2	0.2	0.2
370	i	1.1	1.3	1.6	1.3	1.5	1.1
	p	2.3	2.3	2.9	2.8	3.0	2.0
371	i	0.2	0.0	0.0	0.0	0.1	0.0
	p	0.2	0.0	0.0	0.1	0.1	0.1
372	i	0.1	0.3	0.2	0.5	0.4	0.2
	p	0.3	0.4	0.3	0.7	0.5	0.3
373	i	0.1	0.0	0.1	0.0	0.1	0.0
	p	0.1	0.0	0.1	0.0	0.1	0.1
374	i	0.2	0.1	0.2	0.2	0.2	0.1
	p	0.8	0.7	0.3	0.8	0.7	0.4
375	i	0.2	0.1	0.2	0.1	0.1	0.1
	p	0.2	0.1	0.3	0.3	0.3	0.2
376	i	.	0.0	0.0	.	.	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
377	i	0.1	0.2	0.5	0.3	0.3	0.2
	p	0.2	0.4	1.0	0.6	0.6	0.5
378	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.3	0.2	0.2	0.1
379	i	0.5	0.3	0.3	0.1	0.2	0.5
	p	0.9	0.4	0.4	0.2	0.4	0.6
380	i	0.6	0.0	0.2	0.2	0.2	0.3
	p	0.7	0.0	0.3	0.2	0.2	0.4
381	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	0.1	0.1	0.1	0.0	0.1	0.1
382	i	0.1	0.1	0.0	0.0	0.1	0.0
	p	0.1	0.1	0.0	0.0	0.1	0.0
383	i	31.4	19.7	38.1	34.1	34.7	22.9
	p	34.3	21.4	40.8	37.5	37.8	24.8
384	i	7.1	5.0	10.2	12.4	8.8	8.6
	p	9.2	6.6	12.0	15.2	11.0	10.1
385	i	3.8	5.4	6.7	8.0	5.8	5.7
	p	4.0	5.5	7.0	8.3	6.1	5.9

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
386	acute laryngitis/tracheitis/croup	i	2.0	3.7	5.7	3.2	1.9	3.6	4.2
		p	2.0	3.9	6.1	3.3	2.0	3.7	4.6
387	acute bronchi(ol)itis	i	9.5	9.1	10.9	9.4	12.0	9.0	10.5
		p	13.0	12.0	14.5	13.4	14.4	12.5	13.9
388	influenza w/o pneumonia	i	5.4	5.7	6.8	8.2	2.9	5.7	8.6
		p	5.5	5.9	7.0	8.4	3.1	5.9	8.7
389	pneumonia	i	1.9	1.6	3.4	1.8	1.1	2.1	2.9
		p	2.0	1.6	3.5	2.0	1.2	2.1	3.0
390	pleurisy all (excl. R70)	i	0.0	0.0	0.1	0.1	0.0	0.1	0.0
		p	0.1	0.1	0.2	0.1	0.1	0.1	0.1
391	infection resp. syst. NEC	i	0.6	0.8	1.0	0.4	1.2	0.5	1.1
		p	1.1	1.5	1.5	0.4	2.0	0.9	2.0
392	mal. neopl. bronchus/lung	i	0.1	0.1	0.1	0.2	0.1	0.1	0.1
		p	1.0	0.8	1.4	1.2	1.1	1.0	1.0
393	mal. neopl. resp. syst. NEC	i	0.0	0.0	0.1	0.1	0.0	0.0	0.0
		p	0.2	0.2	0.3	0.6	0.3	0.3	0.2
394	ben. neopl. resp. syst.	i	.	0.1	0.0	.	0.0	0.0	0.0
		p	0.0	0.1	0.1	.	0.0	0.1	0.1
395	unspecified neopl. resp. syst.	i	0.0	0.0	0.1	.	0.0	0.0	0.0
		p	0.1	0.2	0.2	0.1	0.1	0.2	0.1
396	foreign body resp. syst.	i	0.1	0.0	0.0	.	0.1	0.0	0.1
		p	0.1	0.0	0.0	.	0.2	0.0	0.1
397	injuries resp. syst. NEC	i	0.1	0.1	0.0	.	0.0	0.0	0.0
		p	0.1	0.1	0.0	.	0.0	0.0	0.1
398	cong. anom. resp. syst.	i	.	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.0	0.1	.	.	0.0	0.1
399	hypertrophy/chron. inf. tonsills/aden.	i	0.5	0.5	0.7	1.2	0.6	0.7	0.6
		p	0.7	0.7	1.2	2.0	0.9	1.1	0.8
400	chronic bronchitis	i	1.2	0.8	1.0	0.8	1.3	1.0	0.8
		p	14.4	8.2	8.6	9.7	18.0	10.0	8.4
401	pleural effusion NOS	i	0.0	0.1	0.2	0.1	0.0	0.1	0.0
		p	0.0	0.1	0.2	0.1	0.1	0.1	0.1
402	emfysema	i	0.1	0.2	0.3	0.2	0.1	0.2	0.3
		p	3.1	3.0	3.7	3.0	3.2	2.8	4.0
403	asthma	i	1.8	1.6	2.4	1.5	2.1	1.7	2.1
		p	9.5	10.3	11.5	9.1	10.9	9.2	12.3
404	allergic rhinitis/hay fever	i	1.7	2.5	2.5	1.7	1.8	2.1	2.5
		p	6.7	7.3	6.3	4.6	6.8	6.4	6.8
405	diseases resp. syst. NEC	i	0.7	0.6	0.5	0.4	0.7	0.6	0.5
		p	1.8	1.4	1.5	0.7	1.3	1.4	1.9

R RESPIRATORY

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
386	i	4.4	1.3	4.2	4.3	3.6	3.3
	p	4.6	1.3	4.5	4.6	3.8	3.5
387	i	8.6	7.2	11.9	11.0	10.1	8.9
	p	11.7	9.6	14.8	16.1	13.7	11.4
388	i	5.3	3.5	8.3	7.7	7.2	4.1
	p	5.3	3.6	8.5	7.8	7.4	4.2
389	i	1.3	2.3	2.7	2.1	1.9	2.2
	p	1.3	2.4	2.8	2.3	2.0	2.3
390	i	0.0	0.0	0.1	0.1	0.0	0.0
	p	0.0	0.0	0.2	0.2	0.1	0.1
391	i	0.2	0.5	1.6	0.6	0.8	0.7
	p	0.7	0.9	2.2	1.3	1.4	1.0
392	i	0.1	0.1	0.1	0.0	0.2	0.1
	p	1.0	1.1	1.0	0.9	1.0	0.6
393	i	0.1	0.0	0.1	0.0	0.1	0.0
	p	0.3	0.2	0.4	0.1	0.3	0.1
394	i	0.1	0.0	.	0.0	0.0	0.0
	p	0.1	0.0	0.1	0.1	0.1	0.0
395	i	0.0	0.1	0.0	.	0.0	0.0
	p	0.3	0.1	0.1	0.1	0.2	0.1
396	i	0.1	0.1	0.0	0.1	0.1	0.0
	p	0.1	0.1	0.0	0.1	0.1	0.1
397	i	0.0	0.1	0.1	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.0	0.0
398	i	.	.	0.0	0.0	0.0	0.0
	p	0.1	.	0.0	0.0	0.0	0.0
399	i	1.2	0.6	0.3	0.5	0.5	0.8
	p	1.7	0.9	0.6	0.8	0.9	1.2
400	i	1.1	0.9	1.0	0.9	1.2	0.6
	p	8.8	10.2	12.0	10.9	12.5	6.7
401	i	0.1	0.1	0.1	0.0	0.1	0.0
	p	0.2	0.1	0.1	0.1	0.1	0.0
402	i	0.2	0.2	0.2	0.2	0.3	0.1
	p	2.1	2.7	4.5	3.3	3.7	2.0
403	i	1.5	1.8	2.3	1.6	1.9	1.6
	p	7.4	8.7	13.8	10.8	11.3	8.2
404	i	3.6	2.1	1.7	1.2	2.2	2.1
	p	11.4	5.4	4.1	5.4	6.8	6.1
405	i	0.7	0.5	0.7	0.4	0.6	0.4
	p	1.6	1.7	1.2	1.4	1.6	1.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
406	pain/tenderness of skin	i	0.0	0.1	.	0.1	0.0	0.1	0.0
		p	0.1	0.1	0.0	0.2	0.1	0.1	0.1
407	pruritus, skin itching	i	0.6	0.9	1.2	0.7	0.7	0.8	1.1
		p	1.8	1.9	2.5	1.3	1.9	1.7	2.3
408	warts	i	4.4	5.0	7.0	3.1	5.9	4.8	5.4
		p	5.8	7.6	8.3	4.2	7.3	6.7	6.9
409	condyloma acuminata	i	0.2	0.2	0.6	0.8	0.3	0.4	0.3
		p	0.2	0.2	0.6	0.8	0.3	0.4	0.3
410	local swelling/papul/lump	i	0.5	0.5	0.5	0.4	0.8	0.4	0.6
		p	0.6	0.7	0.6	0.8	0.9	0.5	0.8
411	multiple swelling/papul/lump	i	0.1	0.2	0.1	0.1	0.2	0.1	0.2
		p	0.2	0.2	0.1	0.4	0.2	0.2	0.3
412	local redness/erythema	i	0.6	0.7	0.6	0.7	0.6	0.6	0.8
		p	1.1	1.4	1.0	0.8	1.0	0.9	1.9
413	general redness/erythema	i	0.1	0.2	0.3	0.2	0.2	0.1	0.4
		p	0.4	0.5	0.4	0.2	0.4	0.3	0.7
414	changes in skin color NEC	i	0.1	0.0	0.2	0.3	0.0	0.2	0.1
		p	0.2	0.1	0.2	0.5	0.1	0.2	0.1
415	local infection finger/toe	i	1.4	1.9	1.3	1.1	1.7	1.5	1.6
		p	1.5	2.0	1.4	1.1	1.7	1.6	1.7
416	boil/carbuncle/cellulitis	i	1.7	2.4	1.4	2.2	1.6	2.0	1.9
		p	2.1	2.8	1.7	3.1	1.9	2.4	2.4
417	local infection skin NEC	i	1.4	1.2	1.0	1.1	1.4	1.0	1.5
		p	1.9	1.6	1.5	1.2	1.8	1.5	1.9
418	insect bite	i	1.5	1.2	0.8	0.7	0.9	1.1	1.3
		p	1.6	1.3	0.9	0.7	0.9	1.2	1.3
419	animal/human bite	i	0.6	0.6	0.4	0.4	0.7	0.5	0.5
		p	0.7	0.7	0.5	0.5	0.8	0.6	0.6
420	burns/scalds	i	0.9	0.8	0.6	0.8	0.6	0.8	0.8
		p	1.4	1.1	0.8	1.0	0.8	1.1	1.3
421	foreign body in skin	i	0.6	0.5	0.4	0.2	0.4	0.4	0.6
		p	0.7	0.5	0.5	0.2	0.4	0.5	0.6
422	bruise/contusion intact skin	i	2.1	1.9	2.3	1.9	2.0	1.9	2.5
		p	2.4	2.2	2.5	1.9	2.1	2.2	2.8
423	abrasion/scratch/blister	i	1.1	0.8	0.9	1.0	0.8	1.0	0.9
		p	1.2	0.9	0.9	1.0	0.8	1.1	0.9
424	laceration/cut	i	4.5	4.1	2.6	2.1	3.2	3.6	4.1
		p	5.0	4.5	3.0	2.3	3.5	4.0	4.6
425	injury to skin NEC	i	2.5	1.8	1.2	1.9	3.2	1.5	2.2
		p	3.5	2.8	2.0	2.8	4.4	2.4	3.4
426	corns/callosities	i	0.8	0.9	0.7	1.2	0.6	0.8	1.0
		p	1.1	1.1	0.9	1.2	0.7	1.1	1.2
427	sympt./complt. of skin texture	i	0.4	0.7	0.4	0.5	0.4	0.6	0.6
		p	1.1	1.7	1.1	1.4	1.0	1.5	1.0
428	sympt./complt. nails	i	0.2	0.1	0.2	.	0.2	0.1	0.2
		p	0.3	0.3	0.3	0.1	0.4	0.2	0.3
429	losing hair/baldness	i	0.1	0.2	0.3	0.5	0.2	0.2	0.2
		p	0.2	0.5	0.7	1.1	0.3	0.6	0.5
430	sympt./complt. hair NEC	i	0.1	0.1	0.3	0.2	0.1	0.1	0.2
		p	0.4	0.2	0.4	0.5	0.4	0.3	0.4

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
406	i	0.1	0.0	0.0	0.0	0.0	0.1
	p	0.3	0.1	0.0	0.1	0.1	0.1
407	i	0.9	0.5	0.9	1.0	1.0	0.7
	p	2.1	1.5	1.9	2.2	2.2	1.3
408	i	5.5	3.8	6.2	4.7	4.9	4.9
	p	7.9	5.8	7.6	6.0	6.5	6.9
409	i	0.3	0.1	0.5	0.5	0.4	0.2
	p	0.3	0.1	0.5	0.5	0.4	0.2
410	i	0.4	0.5	0.4	0.7	0.5	0.4
	p	0.6	0.6	0.6	0.8	0.7	0.5
411	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.3	0.1	0.2	0.2	0.2
412	i	0.9	0.5	0.8	0.4	0.7	0.5
	p	1.5	1.0	1.0	1.0	1.3	0.9
413	i	0.3	0.2	0.1	0.2	0.2	0.1
	p	0.5	0.5	0.3	0.3	0.5	0.3
414	i	0.1	0.2	0.2	0.1	0.2	0.1
	p	0.1	0.2	0.2	0.2	0.2	0.1
415	i	1.8	1.0	1.8	1.6	1.5	1.3
	p	1.9	1.1	1.9	1.6	1.6	1.4
416	i	1.6	1.6	2.4	2.1	2.1	1.6
	p	2.0	1.9	2.9	2.7	2.6	1.8
417	i	0.9	1.1	1.3	1.5	1.3	0.9
	p	1.3	1.4	1.9	2.0	1.8	1.3
418	i	0.9	2.8	0.7	0.1	1.1	1.0
	p	0.9	3.0	0.8	0.2	1.2	1.1
419	i	0.6	0.7	0.5	0.4	0.6	0.4
	p	0.7	0.7	0.6	0.5	0.6	0.5
420	i	1.0	1.1	0.5	0.6	0.9	0.6
	p	1.4	1.4	0.8	0.8	1.3	0.8
421	i	0.6	0.5	0.3	0.5	0.5	0.4
	p	0.7	0.5	0.4	0.6	0.6	0.5
422	i	1.6	2.0	2.6	2.0	2.4	1.4
	p	1.8	2.3	3.0	2.2	2.7	1.6
423	i	1.1	1.2	0.7	0.8	1.1	0.6
	p	1.2	1.4	0.7	0.8	1.2	0.7
424	i	4.9	3.9	3.1	2.9	3.6	3.5
	p	5.4	4.3	3.5	3.1	4.0	3.8
425	i	1.3	2.7	1.8	1.7	2.0	1.6
	p	2.2	3.7	2.9	2.6	3.1	2.3
426	i	1.1	0.8	0.7	0.7	1.0	0.7
	p	1.4	1.0	0.9	0.9	1.2	0.9
427	i	0.4	0.5	0.8	0.5	0.6	0.4
	p	1.0	1.3	1.7	1.3	1.5	1.0
428	i	0.2	0.2	0.1	0.1	0.2	0.1
	p	0.3	0.3	0.2	0.2	0.3	0.2
429	i	0.2	0.2	0.3	0.1	0.2	0.2
	p	0.5	0.7	0.6	0.3	0.6	0.5
430	i	0.1	0.1	0.2	0.1	0.2	0.0
	p	0.5	0.4	0.3	0.3	0.5	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
431	fear of skin disease	i	0.0	0.0	0.0	0.1	0.1	0.0	0.0
		p	0.0	0.0	0.2	0.1	0.1	0.1	0.0
432	disability/impairment of skin	i
		p	.	0.0	0.0	.	0.0	0.0	.
433	sympt./complt. of skin NEC	i	0.3	0.3	0.2	0.4	0.2	0.3	0.2
		p	0.8	0.5	0.6	0.7	0.6	0.7	0.6
434	herpes zoster	i	0.8	0.9	1.1	1.7	0.9	1.0	1.0
		p	1.3	1.1	1.2	2.0	1.1	1.3	1.2
435	herpes simplex lip/mouth	i	0.6	0.7	0.9	0.7	0.5	0.7	0.9
		p	1.3	1.2	1.4	1.8	1.0	1.4	1.3
436	herpes simplex genitalis	i	0.0	0.1	0.1	0.5	0.0	0.2	0.1
		p	0.1	0.2	0.1	0.7	0.1	0.2	0.1
437	herpes simplex NEC	i	0.2	0.2	0.3	0.2	0.1	0.2	0.2
		p	0.3	0.2	0.5	0.5	0.2	0.4	0.3
438	scabies/other ascariasis	i	0.1	0.1	0.2	0.5	0.2	0.2	0.0
		p	0.1	0.1	0.2	0.7	0.2	0.3	0.1
439	pediculosis/skininfestation	i	0.2	0.4	0.5	0.4	0.3	0.3	0.4
		p	0.4	0.5	0.6	0.5	0.5	0.4	0.5
440	dermatophytosis/dermatomycosis	i	5.1	6.0	6.0	6.5	4.2	6.0	5.8
		p	8.1	8.2	9.0	8.8	6.3	8.6	8.9
441	moniliasis/candidiasis (excl. GU)	i	0.8	0.7	0.9	1.4	1.1	0.7	1.1
		p	1.4	1.1	1.2	1.7	2.0	1.1	1.5
442	erysipelas	i	1.0	0.5	0.7	1.0	0.7	0.7	0.9
		p	1.4	0.6	0.8	1.3	0.9	0.9	1.3
443	mollusca contagiosa	i	0.5	1.0	1.0	0.3	0.3	0.8	0.6
		p	0.6	1.1	1.0	0.3	0.3	0.9	0.8
444	infection skin NEC	i	0.7	0.6	0.6	0.4	0.9	0.6	0.6
		p	0.9	0.9	0.7	0.6	1.1	0.8	0.8
445	malign. neoplasm skin	i	0.3	0.3	0.1	0.3	0.2	0.2	0.3
		p	0.7	0.6	0.4	0.7	0.6	0.6	0.6
446	lipoma	i	0.6	0.9	0.8	1.4	0.7	1.0	0.5
		p	0.7	1.0	1.0	1.6	0.9	1.1	0.6
447	benign. neoplasm skin NEC	i	1.1	1.9	1.6	0.8	1.6	1.3	1.6
		p	1.4	2.1	1.8	1.3	1.9	1.6	1.9
448	unspecified neoplasm skin	i	0.1	0.2	0.1	0.0	0.1	0.1	0.1
		p	0.1	0.2	0.1	0.0	0.1	0.1	0.2
449	hemangioma/lymfangioma	i	0.1	0.2	0.0	0.5	0.1	0.2	0.1
		p	0.1	0.2	0.0	0.5	0.2	0.2	0.1
450	nevus/mole	i	0.7	0.8	0.9	0.8	1.3	0.8	0.6
		p	0.9	1.0	1.3	1.1	1.8	1.1	0.7
451	cong. anom. of skin NEC	i	.	0.0	.	0.1	.	0.0	0.0
		p	0.0	0.0	.	0.1	.	0.0	0.0
452	impetigo	i	2.0	1.3	1.2	1.0	1.4	1.4	1.8
		p	2.1	1.4	1.3	0.9	1.4	1.4	1.9
453	pilonidal cyst/fistula	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.2	0.2	0.1	0.1	0.2	0.2	0.1
454	seborrhœic dermatitis	i	0.9	1.3	0.9	1.2	0.9	1.0	1.1
		p	1.5	1.8	1.4	1.8	1.7	1.5	1.8
455	atopic dermatitis/eczema	i	0.8	1.4	1.0	1.7	1.3	1.1	1.2
		p	1.5	2.6	2.1	2.8	2.3	2.2	2.1

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
431	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.1
432	i
	p	0.0	0.0	.	0.0	0.0	.
433	i	0.4	0.3	0.3	0.2	0.3	0.2
	p	0.7	0.5	0.7	0.7	0.7	0.5
434	i	1.0	0.8	1.1	1.0	1.0	1.0
	p	1.4	1.2	1.4	1.2	1.3	1.3
435	i	0.6	0.8	0.8	0.7	0.8	0.5
	p	1.2	1.5	1.1	1.6	1.5	1.1
436	i	0.1	0.2	0.1	0.1	0.1	0.1
	p	0.2	0.2	0.1	0.1	0.1	0.2
437	i	0.2	0.2	0.2	0.1	0.2	0.2
	p	0.5	0.4	0.3	0.1	0.4	0.2
438	i	0.2	0.0	0.4	0.0	0.2	0.1
	p	0.2	0.1	0.5	0.1	0.2	0.2
439	i	0.3	0.3	0.4	0.4	0.4	0.2
	p	0.4	0.4	0.5	0.5	0.5	0.3
440	i	5.8	6.6	5.8	4.9	6.0	5.5
	p	8.4	9.3	8.9	7.0	8.8	7.8
441	i	0.7	0.9	1.2	0.8	0.9	0.8
	p	0.8	1.2	1.7	1.5	1.4	1.1
442	i	0.6	1.0	0.7	0.8	0.9	0.4
	p	0.8	1.2	0.9	1.2	1.2	0.6
443	i	1.0	0.5	0.4	1.0	0.5	1.1
	p	1.2	0.5	0.5	1.1	0.6	1.2
444	i	0.7	0.9	0.4	0.4	0.6	0.6
	p	1.0	1.0	0.7	0.5	0.9	0.7
445	i	0.2	0.3	0.2	0.2	0.3	0.2
	p	0.5	0.8	0.6	0.5	0.6	0.5
446	i	0.6	1.0	1.2	0.6	0.9	0.7
	p	0.8	1.1	1.3	0.8	1.0	0.8
447	i	1.8	0.9	1.5	1.6	1.4	1.6
	p	2.1	1.2	1.7	1.9	1.6	1.8
448	i	0.2	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.2
449	i	0.2	0.2	0.1	0.1	0.1	0.2
	p	0.2	0.2	0.1	0.2	0.1	0.2
450	i	0.9	0.7	0.8	0.9	0.8	0.8
	p	1.1	0.9	0.9	1.2	1.0	1.0
451	i	0.0	0.0	.	0.0	0.0	0.0
	p	0.1	0.1	.	0.0	0.0	0.1
452	i	0.9	1.9	2.0	1.2	1.6	1.3
	p	0.9	1.9	2.0	1.3	1.6	1.4
453	i	0.1	0.1	0.1	0.1	0.1	0.0
	p	0.1	0.2	0.2	0.1	0.2	0.1
454	i	1.3	0.9	1.0	1.0	1.0	1.1
	p	2.0	1.3	1.5	1.7	1.6	1.7
455	i	1.3	0.9	1.3	1.1	1.1	1.2
	p	2.9	1.6	2.0	2.2	2.1	2.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
456	contact dermatitis	i	1.3	1.2	1.6	1.0	1.2	1.1	1.6
		p	1.8	1.9	2.2	1.3	1.8	1.7	2.4
457	eczema NEC	i	6.4	7.8	9.0	6.1	5.6	7.4	8.3
		p	16.7	15.2	17.1	12.7	15.1	15.7	16.5
458	diaper rash	i	0.5	0.5	0.5	0.5	0.6	0.4	0.6
		p	0.7	0.7	0.7	0.9	0.9	0.7	0.8
459	pityriasis rosea	i	0.3	0.4	0.6	0.3	0.4	0.3	0.5
		p	0.4	0.5	0.7	0.4	0.6	0.5	0.5
460	psoriasis w/wo arthropathy	i	0.5	0.5	0.5	0.2	0.7	0.4	0.5
		p	2.1	1.9	2.0	1.1	2.3	1.5	2.7
461	pompholyx/dis. sweat glands	i	0.3	0.5	0.6	0.1	0.3	0.5	0.3
		p	0.4	0.6	0.7	0.2	0.4	0.5	0.5
462	sebaceous cyst	i	2.0	2.3	2.9	2.9	2.1	2.3	2.8
		p	2.5	2.7	3.4	2.9	2.6	2.6	3.4
463	ingrown toenail/dis. of nail NEC	i	0.9	1.2	1.0	0.6	0.9	0.9	1.2
		p	1.2	1.5	1.3	0.8	1.1	1.2	1.5
464	alopecia/folliculitis/dis. of hair NEC	i	0.9	0.8	1.5	0.3	0.4	0.9	1.2
		p	1.1	1.1	1.8	1.0	0.6	1.2	1.6
465	acne vulgaris	i	1.3	1.9	1.7	1.9	1.4	1.6	1.8
		p	5.9	5.8	6.2	4.2	4.8	5.7	6.2
466	chron. ulcer skin, incl. varicose	i	0.4	0.3	0.3	0.6	0.4	0.3	0.4
		p	1.4	1.1	0.7	1.5	1.4	1.0	1.4
467	urticaria	i	1.4	1.2	1.4	0.5	1.4	1.2	1.3
		p	1.6	1.4	1.6	0.6	1.7	1.3	1.5
468	seborrheic warts	i	0.2	0.3	0.2	0.2	0.3	0.2	0.3
		p	0.3	0.3	0.3	0.3	0.4	0.3	0.3
469	acne rosacea	i	0.2	0.3	0.3	0.2	0.2	0.2	0.3
		p	0.3	0.5	0.5	0.3	0.4	0.4	0.4
470	dis. skin/subcut. tissue NEC	i	1.1	1.6	1.6	1.6	1.3	1.5	1.3
		p	2.3	2.6	2.3	3.2	2.1	2.5	2.7

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
456	i	1.1	1.2	1.8	1.0	1.4	1.0
	p	1.6	2.0	2.3	1.5	2.1	1.4
457	i	7.9	6.1	8.0	7.7	7.7	6.9
	p	15.0	14.1	15.7	18.4	16.9	13.9
458	i	0.4	0.5	0.6	0.5	0.5	0.3
	p	0.7	0.8	0.9	0.6	0.8	0.5
459	i	0.3	0.3	0.5	0.4	0.4	0.3
	p	0.5	0.3	0.6	0.6	0.5	0.4
460	i	0.4	0.6	0.5	0.3	0.4	0.5
	p	2.0	1.7	1.9	1.9	2.1	1.5
461	i	0.3	0.2	0.7	0.4	0.4	0.3
	p	0.5	0.3	0.8	0.4	0.6	0.3
462	i	2.3	2.4	2.6	2.3	2.5	2.0
	p	2.8	2.8	3.1	2.6	3.0	2.3
463	i	1.1	0.7	0.9	1.1	1.0	0.8
	p	1.5	1.0	1.2	1.3	1.3	1.0
464	i	0.9	1.1	0.9	0.8	1.0	0.8
	p	1.2	1.3	1.5	1.0	1.4	1.1
465	i	1.8	1.4	2.0	1.4	1.8	1.3
	p	5.1	5.3	6.0	6.4	6.3	4.6
466	i	0.2	0.5	0.6	0.2	0.4	0.2
	p	0.9	1.4	1.3	1.0	1.3	0.6
467	i	1.2	1.4	1.2	1.1	1.3	0.9
	p	1.5	1.6	1.4	1.3	1.6	1.0
468	i	0.3	0.2	0.3	0.2	0.3	0.2
	p	0.4	0.3	0.3	0.2	0.3	0.2
469	i	0.3	0.2	0.3	0.2	0.2	0.2
	p	0.6	0.3	0.4	0.3	0.4	0.3
470	i	1.6	1.8	1.3	1.0	1.5	1.2
	p	2.7	2.5	2.2	2.5	2.7	2.0

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
471	excessive thirst	i	0.0	0.0	0.1	0.1	.	0.0	0.0
		p	0.0	0.0	0.1	0.1	.	0.1	0.0
472	excessive appetite, excl. bulimia	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.0	0.0	0.0	0.5	0.0	0.1	0.0
473	loss of appetite, excl. anorexia nerv.	i	0.0	0.1	0.1	0.2	0.1	0.1	0.1
		p	0.1	0.1	0.1	0.3	0.1	0.2	0.1
474	feeding problem infant, excl. P11	i	0.1	0.2	0.1	0.3	0.1	0.1	0.2
		p	0.2	0.2	0.1	0.3	0.2	0.2	0.2
475	feeding problem adult, excl. T06	i	0.1	0.1	.	0.3	0.1	0.1	0.1
		p	0.3	0.3	0.1	1.4	0.2	0.4	0.2
476	feeding problem NEC	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.1	0.1	0.1	0.0	0.1	0.1
477	weight gain, excl. T82	i	0.0	0.1	0.0	0.2	0.0	0.1	0.1
		p	0.0	0.1	0.0	0.6	0.1	0.2	0.1
478	weight loss, excl. T06	i	0.1	0.1	0.1	0.2	0.2	0.1	0.1
		p	0.2	0.3	0.2	0.5	0.3	0.2	0.3
479	weight problem NEC	i	0.1	0.2	0.1	0.2	0.0	0.2	0.1
		p	0.2	0.5	0.4	0.6	0.1	0.4	0.5
480	failure to thrive	i	0.2	0.1	0.2	0.2	0.0	0.1	0.1
		p	0.3	0.2	0.3	0.4	0.2	0.3	0.3
481	dehydration	i	0.0	0.0	0.0	.	0.1	0.0	0.0
		p	0.1	0.1	0.1	.	0.1	0.1	0.0
482	thyroid lump/mass	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.0	0.0	.	.	0.0	0.0
483	fear of endocr. metab. nutr. dis.	i	0.1	0.1	0.1	0.1	0.1	0.1	0.2
		p	0.1	0.1	0.2	0.1	0.1	0.1	0.2
484	disability/impairment end. gl.	i
		p	0.0	0.0
485	sympt./complt. end./met./nutr. NEC	i	0.2	0.1	0.2	0.3	0.1	0.2	0.2
		p	0.3	0.1	0.3	0.4	0.2	0.2	0.3
486	infectious dis. endocrine glands	i
		p	0.0	0.0
487	malign. neopl. thyroid	i	.	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.1	0.1	0.2	0.0	0.0
488	benign. neopl. thyroid	i	.	0.0	.	.	0.0	.	.
		p	0.0	0.0	0.0	.	0.0	.	0.0
489	unspecified neopl. endocr. NEC	i	0.0	0.0	0.0	.	0.0	0.0	.
		p	0.0	0.1	0.1	.	0.1	0.1	0.0
490	injury thyroid	i
		p	.	0.0	.	.	0.0	0.0	.
491	thyroglossal duct(cyst)	i	0.0	0.0	0.0	0.1	0.0	0.0	.
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
492	cong. anom. endocr. NEC	i	.	.	0.0	.	.	.	0.0
		p	0.0	0.0	0.1	0.1	0.0	0.0	0.0
493	goiter/module w/o thyrotox	i	0.0	0.1	0.0	.	0.1	0.0	0.1
		p	0.3	0.2	0.3	.	0.5	0.2	0.2
494	obesity/overweight	i	1.1	1.5	1.6	1.4	1.0	1.3	1.7
		p	4.9	5.6	7.7	5.0	6.2	5.3	6.8
495	hyperthyroidism/thyrotoxicosis	i	0.2	0.1	0.1	0.2	0.2	0.2	0.1
		p	1.4	1.2	0.9	1.8	1.4	1.3	1.3

T ENDOCRINE/METABOLIC/NUTRITIONAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
471	i	0.1	.	0.1	0.0	0.1	0.0
	p	0.1	.	0.1	0.0	0.1	0.0
472	i	.	0.0	0.1	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.1	0.1	0.1
473	i	0.1	0.1	0.0	0.0	0.1	0.0
	p	0.2	0.1	0.1	0.1	0.1	0.1
474	i	0.1	0.1	0.2	0.1	0.2	0.1
	p	0.3	0.1	0.3	0.1	0.2	0.2
475	i	0.1	0.2	0.1	0.2	0.1	0.1
	p	0.2	0.4	0.2	0.7	0.5	0.1
476	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.0	0.1	0.2	0.1	0.1
477	i	0.2	0.0	0.1	0.0	0.1	0.1
	p	0.2	0.1	0.1	0.2	0.2	0.1
478	i	0.1	0.1	0.1	0.2	0.1	0.1
	p	0.1	0.1	0.4	0.4	0.3	0.1
479	i	0.2	0.0	0.1	0.2	0.1	0.1
	p	0.6	0.3	0.2	0.4	0.5	0.3
480	i	0.3	0.1	0.1	0.0	0.1	0.2
	p	0.4	0.3	0.3	0.2	0.2	0.3
481	i	.	0.1	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.0	0.1	0.1	0.0
482	i	.	0.0	0.1	0.0	0.0	0.0
	p	.	0.0	0.1	0.0	0.0	0.0
483	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.1	0.1	0.1	0.2	0.1	0.1
484	i
	p	.	0.0	.	.	.	0.0
485	i	0.0	0.3	0.2	0.2	0.2	0.1
	p	0.0	0.4	0.3	0.3	0.3	0.1
486	i
	p	0.0	.	.	.	0.0	.
487	i	.	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.0	0.1	0.1	0.0
488	i	0.0	.	.	.	0.0	.
	p	0.0	0.0	.	0.0	0.0	0.0
489	i	0.1	0.0	0.0	.	0.0	0.1
	p	0.1	0.0	0.1	0.0	0.0	0.1
490	i
	p	0.0	0.0	.	.	0.0	.
491	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
492	i	.	.	0.0	.	0.0	.
	p	0.0	0.0	0.0	0.0	0.0	.
493	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.2	0.2	0.2	0.3	0.3	0.2
494	i	1.2	1.6	1.1	1.6	1.5	1.1
	p	5.2	5.9	5.5	6.5	6.6	4.2
495	i	0.1	0.1	0.2	0.2	0.2	0.1
	p	1.0	1.2	1.7	1.2	1.6	0.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
496	hypothyroidism/myxedema	i	0.1	0.1	0.1	0.0	0.1	0.1	0.0
		p	1.9	1.4	1.3	1.0	2.4	1.6	0.9
497	hypoglycemia	i	0.1	0.1	0.2	0.3	0.1	0.2	0.1
		p	0.2	0.2	0.2	0.3	0.1	0.3	0.2
498	renal glucosuria	i	0.0	0.0	.	.	.	0.0	0.0
		p	0.0	0.0	.	0.0	0.0	0.0	0.0
499	diabetes mellitus	i	0.9	0.7	1.2	0.7	0.8	0.9	0.9
		p	14.6	10.1	11.7	10.9	14.9	12.0	10.8
500	vitamin defic./nutr. dis. NEC	i	0.1	0.1	0.1	0.3	0.1	0.1	0.1
		p	2.1	0.8	0.9	2.3	1.8	1.4	1.0
501	gout	i	0.4	0.4	0.4	0.7	0.4	0.5	0.4
		p	1.3	1.1	1.0	1.8	1.3	1.2	1.1
502	lipid metabolism dis.	i	0.2	0.1	0.2	0.2	0.2	0.2	0.1
		p	1.6	1.4	1.9	1.3	2.6	1.5	1.4
503	dis. thyroid NEC	i	0.0	0.1	0.0	0.1	0.1	0.1	0.0
		p	0.4	0.2	0.6	0.4	0.3	0.5	0.3
504	dis. endocr. gland NEC	i	0.1	0.0	0.2	0.1	0.1	0.1	0.1
		p	0.5	0.4	0.9	0.3	0.7	0.5	0.5
505	metabolic disease NEC	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	1.3	0.9	0.4	0.6	1.8	0.8	0.7

T ENDOCRINE/METABOLIC/NUTRITIONAL

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
496	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	1.0	1.6	1.6	1.7	1.7	1.3
497	i	0.0	0.1	0.1	0.2	0.1	0.1
	p	0.0	0.3	0.2	0.3	0.3	0.2
498	i	.	0.0	0.0	.	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
499	i	0.7	1.1	0.9	0.8	1.0	0.6
	p	10.9	13.1	11.6	12.4	14.1	7.7
500	i	0.1	0.2	0.1	0.1	0.1	0.1
	p	0.7	1.3	1.3	2.2	1.6	0.9
<hr/>							
501	i	0.4	0.4	0.6	0.4	0.5	0.5
	p	1.1	1.2	1.2	1.3	1.3	1.1
502	i	0.1	0.1	0.2	0.2	0.1	0.2
	p	0.9	1.2	1.9	2.3	1.6	1.5
503	i	0.0	0.1	0.1	0.0	0.0	0.1
	p	0.2	0.8	0.3	0.3	0.5	0.3
504	i	0.0	0.2	0.1	0.1	0.1	0.1
	p	0.3	0.6	0.6	0.6	0.6	0.5
505	i	0.1	0.2	0.1	0.0	0.1	0.1
	p	0.7	1.1	0.6	1.1	1.0	0.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
506	complaints urination	i	0.4	0.5	0.3	0.7	0.4	0.4	0.5
		p	0.7	0.8	0.5	0.9	0.7	0.6	0.9
507	incontinence (excl. P12)	i	0.2	0.3	0.5	0.2	0.2	0.3	0.4
		p	4.5	3.9	2.8	2.9	5.0	3.3	4.2
508	anuria/oliguria	i	0.0	0.0	.	.	.	0.0	0.0
		p	0.0	0.0	.	.	.	0.0	0.0
509	complaints of urine	i	0.1	0.2	0.3	0.1	0.2	0.2	0.2
		p	0.2	0.3	0.5	0.2	0.4	0.3	0.3
510	sympt./complt. bladder NEC	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.2	0.1	0.5	0.1	0.4	0.3	0.1
511	sympt./complt. kidney	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.1	0.0	0.0	.	0.0	0.1	0.0
512	fear of urinary diseases	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.0	0.0	0.1	0.2	0.0	0.1	0.0
513	disability/impairment urinary syst.	i	.	0.0	.	.	0.0	.	.
		p	.	0.0	.	.	0.0	.	.
514	sympt./complt. urin. syst. NEC	i	0.0	0.0	.	0.0	0.0	0.0	0.0
		p	0.1	0.0	0.1	0.0	0.2	0.1	0.1
515	acute pyelonefritis/pyelitis	i	0.2	0.2	0.6	0.3	0.3	0.3	0.2
		p	0.3	0.3	0.6	0.3	0.4	0.4	0.3
516	cystitis/urinary inf. NEC	i	11.0	11.1	13.2	11.7	12.0	11.6	11.4
		p	12.8	12.2	14.6	12.3	14.0	12.9	12.5
517	non-specific urethritis	i	0.2	0.3	0.4	0.6	0.3	0.3	0.3
		p	0.2	0.4	0.5	0.6	0.3	0.4	0.4
518	malign. neoplasm kidney	i	.	0.0	0.0	.	0.0	.	0.0
		p	0.1	0.1	0.3	0.2	0.2	0.1	0.3
519	malign. neoplasm bladder	i	0.1	0.0	0.1	0.1	0.0	0.1	0.0
		p	0.3	0.2	0.2	0.3	0.4	0.3	0.2
520	malign. neoplasm urinary tract NEC	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.1	0.0	0.1	0.0	0.1	0.0
521	benign. neoplasm urinary tract	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.0	.	0.1	0.0	0.0
522	unspecified neopl. urin. tract	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.1	0.1	0.1	0.0	0.1
523	injury urinary tract	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.0	.	.	0.0	0.0	0.0
524	cong. anom. urinary tract	i	0.0	0.0	0.0	.	0.0	.	0.0
		p	0.1	0.1	0.0	0.1	0.1	0.1	0.1
525	glomerulonephritis/nephrosis	i	0.0	0.1	0.0	0.1	0.0	0.0	0.0
		p	0.3	0.3	0.2	0.3	0.3	0.3	0.2
526	orthostatic albuminuria	i	0.0	0.0	.	.	0.0	0.0	.
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
527	urin. calculus, all types/sites	i	0.4	0.6	0.7	0.7	0.5	0.5	0.7
		p	0.9	1.2	1.4	1.3	1.0	1.1	1.3
528	abnormal urine test/findings NOS	i	0.1	0.1	0.0	.	0.1	0.1	0.1
		p	0.2	0.3	0.2	0.2	0.2	0.3	0.2
529	dis. urinary system NEC	i	0.3	0.2	0.5	0.6	0.4	0.4	0.2
		p	2.0	1.4	1.5	1.8	2.1	1.7	1.3

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
506	i	0.4	0.3	0.4	0.6	0.5	0.3
	p	0.6	0.6	0.8	0.9	0.8	0.4
507	i	0.3	0.2	0.5	0.3	0.4	0.2
	p	2.8	3.8	4.0	4.3	4.7	1.8
508	i	.	0.0	0.0	.	0.0	.
	p	.	0.0	0.0	.	0.0	.
509	i	0.2	0.0	0.2	0.3	0.2	0.2
	p	0.3	0.1	0.4	0.4	0.3	0.3
510	i	0.0	0.0	0.2	0.0	0.1	0.1
	p	0.1	0.2	0.4	0.2	0.2	0.2
511	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.0	0.1	0.1	.	0.1	0.0
512	i	0.0	0.0	0.1	0.0	0.0	0.0
	p	0.0	0.0	0.1	0.0	0.1	0.0
513	i	.	0.0	.	.	0.0	.
	p	.	0.0	.	0.0	0.0	.
514	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.1
515	i	0.2	0.3	0.4	0.2	0.3	0.2
	p	0.3	0.4	0.5	0.2	0.4	0.3
516	i	10.6	11.0	13.1	11.8	12.5	9.5
	p	11.4	12.7	14.4	13.3	13.9	10.6
517	i	0.4	0.3	0.4	0.3	0.4	0.3
	p	0.4	0.3	0.4	0.4	0.4	0.4
518	i	.	.	0.0	0.0	0.0	.
	p	0.1	0.2	0.3	0.2	0.2	0.1
519	i	0.0	0.1	0.0	0.1	0.1	0.0
	p	0.1	0.4	0.2	0.4	0.3	0.2
520	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.1	0.1	0.0	0.1
521	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.0	0.0	0.0	0.1
522	i	.	0.0	0.1	.	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.0
523	i	.	0.0	.	0.0	0.0	0.0
	p	0.0	0.0	.	0.0	0.0	0.0
524	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.2	0.0	0.0	0.1	0.1	0.0
525	i	0.1	0.0	0.1	0.0	0.0	0.0
	p	0.2	0.3	0.3	0.2	0.3	0.1
526	i	.	0.0	.	0.0	.	0.0
	p	.	0.0	0.0	0.0	0.0	0.0
527	i	0.6	0.5	0.7	0.5	0.6	0.6
	p	1.2	1.1	1.2	1.1	1.3	0.9
528	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	0.2	0.2	0.3	0.1	0.2	0.3
529	i	0.3	0.4	0.5	0.2	0.4	0.3
	p	1.2	2.2	1.8	1.5	1.8	1.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
530	question of pregnancy (excl. W02)	i	0.2	0.1	0.3	0.4	0.3	0.2	0.2
		p	0.3	0.1	0.4	0.5	0.4	0.3	0.2
531	fear of being pregnant	i	0.1	0.1	0.2	0.1	0.2	0.1	0.2
		p	0.2	0.2	0.3	0.1	0.2	0.2	0.3
532	ante-partum bleeding	i	0.1	0.1	0.2	0.0	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.0	0.1	0.1	0.1
533	vaginal discharge in pregn.	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
534	vomiting/nausea of pregn.	i	0.1	0.0	0.0	0.0	0.0	0.1	0.0
		p	0.1	0.1	0.0	0.2	0.1	0.1	0.1
535	wanted pregnancy	i	0.2	0.2	0.4	0.4	0.2	0.2	0.4
		p	0.3	0.5	0.8	0.6	0.4	0.5	0.6
536	morning-after pill	i	0.2	0.1	0.1	0.1	0.0	0.2	0.1
		p	0.2	0.2	0.3	0.3	0.1	0.3	0.2
537	fam. plan./oral contraceptive	i	2.9	2.7	3.5	2.5	2.9	3.1	2.4
		p	35.7	34.5	37.1	32.0	32.4	34.9	37.0
538	fam. plan./IUD	i	0.4	0.2	0.5	0.7	0.3	0.5	0.4
		p	1.5	1.2	1.7	2.1	1.2	1.6	1.4
539	fam. plan./sterilization ♀	i	0.2	0.3	0.2	0.2	0.2	0.2	0.3
		p	0.3	0.5	0.3	0.3	0.3	0.4	0.4
540	fam. plan./ NEC ♀	i	0.1	0.2	0.1	0.2	0.2	0.1	0.1
		p	0.6	1.0	0.4	0.9	0.8	0.7	0.7
541	sub-/infertility ♀	i	0.3	0.3	0.2	0.3	0.2	0.3	0.3
		p	0.9	0.8	0.6	0.8	0.4	0.8	0.9
542	heavy post-partum bleeding	i	0.0	0.0	.	0.1	0.0	0.0	0.0
		p	0.1	0.0	.	0.1	0.0	0.1	0.0
543	complt. post partum NEC	i	0.1	0.0	0.1	0.1	0.0	0.1	0.0
		p	0.1	0.1	0.1	0.2	0.1	0.1	0.1
544	sympt./complt. lactation	i	0.1	0.0	0.0	0.0	0.0	0.1	0.1
		p	0.2	0.1	0.0	0.3	0.1	0.1	0.2
545	sympt./complt. breast in pregn. NEC	i	0.1	0.0	0.1	.	0.0	0.0	0.1
		p	0.1	0.1	0.1	.	0.1	0.1	0.1
546	starting delivery	i	0.1	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.1	.	0.1	0.1	0.0
547	fear of complic of pregn./deliv.	i	0.0	0.1	0.1	.	0.1	0.1	0.1
		p	0.1	0.1	0.2	.	0.1	0.1	0.1
548	disability/impairment Pr/D/Pu/Fp	i
		p	.	0.0	.	.	0.0	.	.
549	sympt./complt. Pr/D/Pu/Fp NEC	i	0.1	0.1	0.1	.	0.1	0.1	0.1
		p	0.2	0.2	0.2	0.0	0.2	0.1	0.2
550	infect. dis. urinary tr. pregn./puerp.	i	0.3	0.2	0.0	0.2	0.1	0.2	0.1
		p	0.3	0.2	0.1	0.2	0.2	0.3	0.1
551	infect. dis. pregn. NEC	i	0.2	0.1	0.2	0.1	0.1	0.1	0.2
		p	0.2	0.1	0.2	0.2	0.1	0.2	0.2
552	malign. neoplasm pregn.	i
		p	.	0.0	.	.	0.0	.	.
553	benign. neoplasm pregn.	i	.	0.0	.	.	.	0.0	.
		p	.	0.0	.	.	.	0.0	.
554	injuries complicating pregn.	i
		p

W PREGNANCY, CHILD BEARING, FAMILY PLANNING

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
530	i	0.3	0.2	0.1	0.3	0.2	0.2
	p	0.3	0.3	0.2	0.3	0.3	0.2
531	i	0.2	0.1	0.2	0.1	0.2	0.1
	p	0.3	0.2	0.2	0.2	0.3	0.1
532	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.1	0.1	0.1	0.1	0.1	0.1
533	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.1	0.0	0.1	0.1	0.1	0.1
534	i	0.0	0.1	0.0	0.1	0.1	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.0
<hr/>							
535	i	0.3	0.1	0.3	0.4	0.4	0.1
	p	0.6	0.3	0.5	0.6	0.6	0.2
536	i	.	0.1	0.3	0.2	0.1	0.2
	p	0.0	0.1	0.5	0.2	0.2	0.2
537	i	3.1	2.9	2.6	3.0	3.3	1.9
	p	37.5	31.7	34.4	37.1	39.9	24.9
538	i	0.4	0.3	0.4	0.5	0.4	0.4
	p	1.7	1.1	1.7	1.6	1.5	1.4
539	i	0.3	0.1	0.2	0.3	0.2	0.2
	p	0.5	0.2	0.4	0.4	0.4	0.3
<hr/>							
540	i	0.1	0.1	0.2	0.2	0.2	0.1
	p	0.6	0.5	0.8	1.0	0.9	0.4
541	i	0.4	0.2	0.3	0.2	0.3	0.1
	p	0.9	0.6	0.8	0.9	0.9	0.4
542	i	0.0	0.0	0.0	0.1	0.0	0.0
	p	0.0	0.0	0.1	0.1	0.0	0.0
543	i	0.0	0.1	0.0	0.0	0.0	0.1
	p	0.1	0.2	0.0	0.0	0.1	0.1
544	i	0.0	0.1	0.0	0.0	0.1	0.0
	p	0.1	0.3	0.2	0.1	0.2	0.1
<hr/>							
545	i	0.0	0.1	0.1	0.0	0.0	0.1
	p	0.1	0.1	0.1	0.1	0.0	0.1
546	i	.	0.1	0.0	0.0	0.0	0.0
	p	.	0.1	0.1	0.0	0.0	0.1
547	i	0.1	0.0	0.1	0.0	0.1	0.0
	p	0.2	0.0	0.1	0.0	0.1	0.0
548	i
	p	.	0.0	.	.	0.0	.
549	i	0.1	0.1	0.2	0.0	0.1	0.0
	p	0.2	0.1	0.3	0.1	0.2	0.1
<hr/>							
550	i	0.2	0.2	0.3	0.2	0.2	0.1
	p	0.2	0.2	0.4	0.2	0.3	0.1
551	i	0.0	0.1	0.2	0.3	0.2	0.1
	p	0.0	0.1	0.3	0.3	0.2	0.1
552	i
	p	0.0	.	.	.	0.0	.
553	i	.	.	.	0.0	0.0	.
	p	.	.	.	0.0	0.0	.
554	i
	p

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
555	cong. anom. of mother/complic. pregn.	i	0.0	.	0.0	.	0.0	0.0	.
		p	0.0	0.0	0.1	.	0.0	0.0	.
556	non-obst. cond. complic. pregn. NEC	i	0.2	0.1	0.2	0.1	0.1	0.1	0.2
		p	0.3	0.1	0.3	0.1	0.2	0.2	0.2
557	pregnancy: confirmed	i	4.2	2.7	2.6	2.7	2.9	3.5	2.5
		p	6.6	3.8	3.9	3.7	4.8	5.0	3.9
558	unwanted pregnancy: confirmed	i	0.1	0.1	0.2	0.7	0.2	0.2	0.0
		p	0.1	0.1	0.2	0.7	0.2	0.3	0.0
559	ectopic pregnancy	i	0.0	0.1	0.1	0.1	0.1	0.1	0.0
		p	0.0	0.1	0.1	0.2	0.1	0.1	0.1
560	toxemia/(pre)eclampsia	i	0.0	0.0	0.1	.	0.1	0.0	0.1
		p	0.1	0.0	0.2	.	0.1	0.1	0.1
561	abortion, imminens/spontan.	i	0.4	0.3	0.4	0.3	0.4	0.4	0.4
		p	0.6	0.5	0.5	0.4	0.5	0.5	0.6
562	abortion, (il)legally induced	i	0.0	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.1	0.2	0.2	0.1	0.1	0.2	0.2
563	pregnancy, high risk	i	0.5	0.3	0.4	0.0	0.3	0.3	0.4
		p	1.3	0.6	0.8	0.3	0.8	0.8	1.0
564	normal delivery live born(s)	i	1.0	0.4	0.5	0.4	0.6	0.7	0.6
		p
565	normal delivery dead born(s)	i	0.0	0.0	0.0	.	.	.	0.0
		p
566	complicated delivery live born	i	0.5	0.2	0.2	0.0	0.3	0.3	0.2
		p
567	complicated delivery dead born	i	0.0	0.0	.	0.1	0.0	0.0	0.0
		p
568	normal puerperium	i	2.1	1.1	1.0	0.5	1.8	1.3	1.2
		p	2.8	1.5	1.4	0.6	2.6	1.8	1.5
569	mastitis puerperalis	i	0.3	0.1	0.2	0.1	0.2	0.2	0.1
		p	0.3	0.2	0.2	0.1	0.2	0.2	0.1
570	dis. of breast puerperium NEC	i	0.1	0.1	0.1	0.2	0.1	0.1	0.1
		p	0.2	0.1	0.1	0.2	0.2	0.1	0.2
571	complic. puerperium NEC	i	0.4	0.1	0.2	0.0	0.1	0.2	0.2
		p	0.5	0.2	0.3	0.1	0.2	0.3	0.3
572	diseases Pr/D/Pu NEC	i	0.2	0.1	0.2	0.1	0.1	0.2	0.1
		p	0.3	0.2	0.4	0.2	0.2	0.2	0.2

W PREGNANCY, CHILD BEARING, FAMILY PLANNING

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
555	i	.	.	0.0	0.0	0.0	0.0
	p	.	0.0	0.0	0.1	0.0	0.0
556	i	0.2	0.2	0.1	0.1	0.2	0.1
	p	0.2	0.3	0.2	0.2	0.3	0.2
557	i	2.8	4.1	2.7	3.0	3.2	2.6
	p	4.3	5.8	4.3	4.6	4.7	4.1
558	i	0.1	0.1	0.2	0.2	0.1	0.1
	p	0.2	0.1	0.3	0.2	0.2	0.2
559	i	0.1	0.1	0.1	0.0	0.1	0.0
	p	0.1	0.1	0.1	0.0	0.1	0.1
<hr/>							
560	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.1
561	i	0.2	0.3	0.4	0.5	0.4	0.3
	p	0.4	0.4	0.7	0.6	0.5	0.5
562	i	0.1	0.0	0.2	0.1	0.1	0.0
	p	0.3	0.0	0.2	0.1	0.2	0.1
563	i	0.4	0.5	0.3	0.2	0.4	0.2
	p	0.8	0.9	0.7	1.0	0.9	0.7
564	i	0.5	1.0	0.5	0.5	0.7	0.5
	p
<hr/>							
565	i	.	0.0	0.0	0.0	0.0	.
	p
566	i	0.1	0.4	0.3	0.3	0.3	0.2
	p
567	i	0.0	.	0.0	0.0	0.0	0.0
	p
568	i	1.1	1.4	1.4	1.4	1.4	1.2
	p	1.6	2.1	1.8	1.8	1.8	1.7
569	i	0.2	0.3	0.1	0.1	0.2	0.2
	p	0.2	0.3	0.1	0.1	0.2	0.2
<hr/>							
570	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.1	0.2	0.1	0.1	0.1
571	i	0.3	0.2	0.1	0.2	0.3	0.1
	p	0.3	0.3	0.3	0.3	0.3	0.2
572	i	0.2	0.1	0.2	0.1	0.2	0.1
	p	0.2	0.2	0.3	0.2	0.3	0.2

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
573	genital pain ♀	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
574	menstruation absent/scanty	i	0.4	0.5	0.4	0.2	0.2	0.4	0.5
		p	0.6	0.8	0.8	0.4	0.4	0.7	0.8
575	menstruation excessive	i	0.9	1.0	1.1	0.8	0.9	1.0	1.0
		p	1.5	1.6	2.0	1.1	1.6	1.6	1.7
576	menstrual pain	i	0.9	0.7	1.1	0.4	1.1	0.8	0.8
		p	2.3	1.6	2.0	1.1	2.3	1.9	1.6
577	sympt./complt. menstruation NEC	i	0.9	1.1	1.4	0.8	1.5	1.0	1.1
		p	1.5	1.6	2.0	1.1	2.2	1.5	1.7
578	premenstrual symptoms	i	0.1	0.1	0.2	0.2	0.2	0.1	0.1
		p	0.2	0.2	0.3	0.5	0.3	0.3	0.2
579	sympt./complt. menopausal	i	0.8	1.0	1.1	1.1	1.1	1.0	0.9
		p	2.8	2.7	3.9	3.7	2.9	3.4	2.5
580	postmenopausal bleeding	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.2	0.1	0.1	0.1
581	painful intercourse ♀	i	0.2	0.2	0.1	0.1	0.2	0.2	0.1
		p	0.3	0.3	0.3	0.1	0.2	0.3	0.3
582	postcoital bleeding	i	0.0	0.2	0.1	0.1	0.0	0.1	0.1
		p	0.0	0.2	0.1	0.1	0.0	0.1	0.1
583	vaginal discharge	i	0.6	1.3	0.5	0.5	0.6	0.9	0.7
		p	0.8	1.6	0.7	0.7	0.9	1.1	0.9
584	sympt./complt. vagina NEC	i	0.2	0.2	0.3	0.3	0.4	0.2	0.3
		p	0.4	0.4	0.5	0.4	0.7	0.4	0.5
585	sympt./complt. vulva	i	0.2	0.2	0.2	0.2	0.1	0.2	0.3
		p	0.3	0.2	0.3	0.2	0.3	0.2	0.4
586	sympt./complt. pelvis ♀	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.1	0.0	.	0.0	0.0	0.1
587	lump/mass mamma ♀	i	0.1	0.2	0.1	0.1	0.1	0.2	0.1
		p	0.2	0.3	0.1	0.1	0.1	0.2	0.2
588	pain mamma ♀	i	0.3	0.5	0.3	0.0	0.1	0.3	0.6
		p	0.4	0.6	0.4	0.1	0.2	0.4	0.7
589	sympt./complt. mamma NEC ♀	i	0.8	1.0	1.2	1.4	0.7	1.0	1.1
		p	1.1	1.2	1.5	1.9	1.0	1.3	1.6
590	postponement of menstruation	i	1.2	1.1	0.8	0.5	0.9	1.0	1.0
		p	1.4	1.3	0.9	0.5	1.1	1.2	1.2
591	fear genital/mamma dis. ♀	i	0.2	0.2	0.3	0.2	0.2	0.2	0.2
		p	0.4	0.4	0.7	0.7	0.4	0.5	0.4
592	disab./impairm. gen. syst. ♀	i	0.0	0.0
		p	0.0	0.0
593	sympt./complt. gen. syst. ♀	i	0.0	0.1	0.0	0.1	0.0	0.1	0.0
		p	0.1	0.1	0.1	0.2	0.0	0.1	0.1
594	syphilis, (genit. and NOS) ♀	i	0.0	.	0.0	.	.	0.0	0.0
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
595	gonorrhoea, (genit. and NOS) ♀	i	0.0	0.0	0.0	.	0.0	.	0.0
		p	0.0	0.0	0.1	0.0	0.0	0.0	0.0
596	venereal dis. NEC ♀	i	0.0	0.2	0.2	0.4	0.1	0.2	0.0
		p	0.0	0.2	0.3	0.5	0.1	0.3	0.1
597	urogenital candidiasis ♀	i	2.1	2.6	2.9	2.4	2.7	2.4	2.8
		p	3.0	3.3	3.4	3.2	3.4	3.2	3.2

X FEMALE GENITAL SYSTEM (INCLUDING BREAST)

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
573	i	0.0	0.0	0.0	.	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
574	i	0.4	0.4	0.4	0.4	0.5	0.2
	p	0.7	0.6	0.7	0.7	0.9	0.2
575	i	1.2	0.8	0.9	1.0	1.1	0.6
	p	1.9	1.3	1.5	1.8	1.9	1.2
576	i	0.9	0.7	0.8	0.9	0.9	0.6
	p	2.1	1.7	1.6	2.0	1.9	1.5
577	i	1.0	1.0	1.2	1.0	1.1	0.9
	p	1.5	1.4	1.8	1.5	1.8	1.2
578	i	0.1	0.0	0.2	0.2	0.1	0.2
	p	0.1	0.1	0.3	0.5	0.3	0.3
579	i	0.8	1.4	0.9	0.9	1.1	0.9
	p	2.8	3.5	3.1	3.0	3.2	2.9
580	i	0.1	0.1	0.1	0.0	0.1	0.1
	p	0.2	0.1	0.1	0.1	0.1	0.1
581	i	0.1	0.1	0.2	0.3	0.2	0.1
	p	0.2	0.1	0.4	0.4	0.3	0.2
582	i	0.2	0.0	0.0	0.1	0.1	0.1
	p	0.2	0.0	0.1	0.1	0.1	0.1
583	i	0.6	1.0	0.8	0.9	0.8	0.7
	p	0.8	1.3	1.0	1.1	1.1	1.0
584	i	0.2	0.1	0.3	0.3	0.3	0.2
	p	0.4	0.3	0.5	0.5	0.5	0.4
585	i	0.1	0.1	0.3	0.2	0.2	0.1
	p	0.3	0.2	0.3	0.2	0.3	0.2
586	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
587	i	0.2	0.1	0.1	0.2	0.1	0.1
	p	0.3	0.1	0.2	0.2	0.2	0.2
588	i	0.4	0.2	0.4	0.3	0.4	0.3
	p	0.7	0.3	0.5	0.3	0.5	0.3
589	i	0.7	0.8	1.4	1.1	1.1	0.8
	p	0.9	0.9	1.8	1.6	1.4	1.0
590	i	1.5	1.4	0.3	0.7	0.9	1.1
	p	1.8	1.6	0.5	0.8	1.1	1.3
591	i	0.2	0.2	0.4	0.2	0.2	0.3
	p	0.4	0.5	0.6	0.4	0.5	0.4
592	i	.	0.0	.	.	0.0	.
	p	.	0.0	.	.	0.0	.
593	i	0.0	0.1	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.0	0.1	0.1	0.1
594	i	0.0	0.0	.	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
595	i	0.0	.	0.0	0.0	0.0	.
	p	0.0	0.0	0.0	0.0	0.0	0.0
596	i	0.3	0.1	0.1	0.1	0.2	0.1
	p	0.3	0.1	0.2	0.2	0.2	0.2
597	i	2.2	2.2	3.3	2.3	2.7	2.0
	p	2.8	3.1	3.9	3.0	3.4	2.8

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
598	urogenital trichomoniasis ♀	i	0.2	0.2	0.5	0.3	0.4	0.3	0.2
		p	0.2	0.3	0.5	0.4	0.5	0.3	0.2
599	pelvic inflammatory disease	i	0.3	0.2	0.3	0.4	0.2	0.3	0.3
		p	0.5	0.4	0.4	0.7	0.4	0.5	0.4
600	carcinoma in situ cervix	i	0.0	0.0	.	.	0.0	0.0	.
		p	0.1	0.1	0.0	0.1	0.1	0.1	0.1
601	cervix carcinoma NEC	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.1	0.2	0.2	0.2	0.1	0.1
602	mal. neopl. mamma ♀	i	0.3	0.2	0.2	0.1	0.2	0.2	0.2
		p	1.6	1.4	2.1	1.1	1.7	1.7	1.4
603	mal. neopl. genit. syst. ♀	i	0.0	0.1	0.0	0.1	0.1	0.1	0.0
		p	0.4	0.5	0.3	0.5	0.5	0.4	0.4
604	fibroid/myoma (uterus/cervix)	i	0.2	0.3	0.3	0.3	0.2	0.3	0.2
		p	0.6	0.8	0.5	1.0	0.7	0.7	0.6
605	ben. neopl. mamma ♀	i	0.3	0.2	0.1	0.1	0.3	0.2	0.2
		p	0.3	0.4	0.2	0.2	0.3	0.3	0.3
606	ben. neopl. genit. syst. ♀	i	0.1	0.2	0.2	0.1	0.2	0.2	0.1
		p	0.3	0.3	0.4	0.4	0.4	0.3	0.2
607	unspec. neopl. genit. syst. ♀	i	0.1	0.1	0.1	.	0.0	0.1	0.0
		p	0.2	0.1	0.2	0.1	0.1	0.2	0.1
608	injury genit. syst./mamma ♀	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.1	0.0	0.2	0.1	0.0	0.1	0.0
609	cong. anom. genit. syst./mamma ♀	i	0.0	0.0	.	.	0.0	.	.
		p	0.1	0.0	0.0	.	0.0	0.0	0.0
610	vaginitis/vulvitis NOS	i	0.6	0.8	1.3	1.3	0.9	0.9	1.0
		p	1.1	1.2	2.0	2.0	1.4	1.4	1.4
611	cervicitis/cerv. dis. NEC	i	0.1	0.2	0.4	0.1	0.1	0.2	0.3
		p	0.2	0.4	0.7	0.1	0.3	0.4	0.4
612	abnormal PAP-smear	i	0.0	0.0	0.1	0.1	.	0.1	0.0
		p	0.2	0.2	0.4	0.5	0.1	0.3	0.2
613	uterovaginal prolapse	i	0.6	0.4	0.6	0.4	0.6	0.6	0.4
		p	2.5	2.0	2.2	2.1	2.6	2.2	1.9
614	cystic breast (chronic)	i	0.3	0.3	0.4	0.4	0.3	0.3	0.3
		p	0.4	0.4	0.4	0.6	0.5	0.4	0.5
615	premenstrual tension syndrome	i	0.1	0.2	0.2	0.2	0.2	0.1	0.3
		p	0.3	0.3	0.3	0.3	0.4	0.2	0.4
616	gardnerella infection	i	0.1	0.0	0.2	0.9	0.0	0.2	0.1
		p	0.1	0.1	0.3	1.4	0.1	0.4	0.1
617	infection/abscess gl. Bartholini	i	0.1	0.1	0.1	0.1	0.0	0.1	0.1
		p	0.1	0.2	0.1	0.2	0.0	0.1	0.2
618	endometriosis	i	0.1	0.1	0.0	0.1	0.1	0.1	0.0
		p	0.3	0.2	0.2	0.2	0.3	0.2	0.2
619	atrophial vaginae/vulvae	i	0.1	0.2	0.2	0.0	0.2	0.2	0.2
		p	0.3	0.7	0.6	0.3	0.6	0.5	0.5
620	dis. genit. syst./mamma NEC ♀	i	0.4	0.3	0.5	0.5	0.5	0.4	0.4
		p	0.6	0.6	1.0	0.9	0.8	0.7	0.7

X FEMALE GENITAL SYSTEM (INCLUDING BREAST)

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
598	i	0.2	0.3	0.4	0.1	0.3	0.2
	p	0.3	0.4	0.4	0.1	0.3	0.3
599	i	0.3	0.4	0.3	0.2	0.3	0.2
	p	0.5	0.5	0.5	0.3	0.5	0.3
600	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.0	0.1	0.0
601	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.2	0.1
602	i	0.1	0.2	0.2	0.1	0.2	0.1
	p	1.4	1.9	1.8	1.3	1.7	1.3
<hr/>							
603	i	0.1	0.0	0.1	0.1	0.1	0.0
	p	0.4	0.4	0.5	0.5	0.4	0.3
604	i	0.3	0.2	0.2	0.3	0.3	0.2
	p	0.9	0.6	0.5	0.8	0.7	0.6
605	i	0.3	0.2	0.2	0.2	0.2	0.1
	p	0.5	0.3	0.3	0.3	0.3	0.2
606	i	0.1	0.1	0.2	0.2	0.2	0.2
	p	0.2	0.3	0.4	0.3	0.3	0.3
607	i	0.1	0.0	0.0	0.1	0.1	0.1
	p	0.2	0.1	0.1	0.2	0.2	0.1
<hr/>							
608	i	0.0	0.0	0.0	0.0	0.0	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.1
609	i	0.0	.	.	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.1	0.1	0.0
610	i	1.1	0.9	1.1	0.6	1.0	0.7
	p	1.5	1.4	1.6	1.1	1.5	1.2
611	i	0.1	0.1	0.2	0.3	0.2	0.2
	p	0.3	0.3	0.4	0.5	0.4	0.3
612	i	0.0	0.1	0.1	0.0	0.0	0.0
	p	0.2	0.2	0.4	0.3	0.3	0.2
<hr/>							
613	i	0.5	0.6	0.4	0.6	0.5	0.6
	p	1.9	2.8	1.9	2.1	2.1	2.3
614	i	0.4	0.4	0.3	0.2	0.3	0.2
	p	0.5	0.6	0.4	0.3	0.5	0.4
615	i	0.2	0.1	0.2	0.2	0.2	0.2
	p	0.2	0.2	0.4	0.4	0.3	0.2
616	i	0.1	0.1	0.4	0.2	0.2	0.2
	p	0.1	0.2	0.4	0.4	0.3	0.3
617	i	0.1	0.2	0.1	0.0	0.1	0.1
	p	0.2	0.2	0.1	0.1	0.2	0.1
<hr/>							
618	i	0.1	0.1	0.0	0.1	0.1	0.1
	p	0.2	0.3	0.2	0.2	0.2	0.2
619	i	0.2	0.1	0.3	0.1	0.2	0.2
	p	0.3	0.4	0.8	0.5	0.5	0.6
620	i	0.4	0.5	0.4	0.3	0.5	0.3
	p	0.7	0.7	0.9	0.6	0.8	0.5

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
621	sympt./complt. gen. syst. ♂	i	0.2	0.3	0.3	0.1	0.2	0.2	0.2
		p	0.3	0.4	0.3	0.3	0.3	0.3	0.3
622	discharge penis/urethra	i	0.0	0.0	.	0.0	.	0.0	0.0
		p	0.0	0.0	.	0.0	.	0.0	0.0
623	sympt./complt. prostate	i	0.0	0.0	0.1	0.1	0.1	0.0	0.0
		p	0.1	0.1	0.1	0.1	0.1	0.1	0.1
624	sympt./complt. potency	i	0.1	0.0	0.1	0.1	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.2	0.1	0.1	0.2
625	sympt./complt. sexual dysf. ♂	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.1	0.1	0.0	0.1	0.0	0.1	0.0
626	sub-/infertility ♂	i	0.0	0.1	0.1	.	0.0	0.1	0.1
		p	0.2	0.1	0.2	.	0.1	0.2	0.2
627	fam. plan. ♂	i	0.3	0.4	0.3	0.3	0.5	0.3	0.4
		p	0.4	0.4	0.4	0.5	0.5	0.4	0.5
628	sympt./complt. mamma ♂	i	0.1	0.0	0.0	0.2	0.1	0.1	0.0
		p	0.1	0.0	0.0	0.2	0.1	0.1	0.0
629	fear dis. gen. syst./mamma ♂	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.1	0.1	0.2	0.1	0.1	0.1	0.1
630	disab./impairm. gen. syst./mamma ♂	i
		p	.	0.0	.	.	.	0.0	.
631	sympt./complt. gen. syst./mam.NEC ♂	i	0.0	0.0	0.1	0.1	0.0	0.0	0.1
		p	0.0	0.0	0.2	0.1	0.0	0.1	0.1
632	syphilis ♂	i	.	0.0	.	0.1	0.0	0.0	.
		p	.	0.0	0.0	0.1	0.0	0.0	.
633	gonorrhoea ♂	i	0.0	0.0	0.2	0.5	0.0	0.1	0.1
		p	0.0	0.0	0.2	0.4	0.0	0.1	0.1
634	venereal disease NEC ♂	i	0.0	0.0	0.0	0.2	0.0	0.1	0.0
		p	0.0	0.0	0.1	0.3	0.1	0.1	0.0
635	urogenital trichomoniasis ♂	i	0.0	0.0	0.0	0.1	0.1	0.0	0.0
		p	0.1	0.0	0.0	0.1	0.1	0.1	0.0
636	prostatitis/seminal vesiculitis	i	0.3	0.2	0.3	0.3	0.5	0.3	0.3
		p	0.5	0.4	0.7	0.5	0.7	0.5	0.5
637	orchitis/epididymitis	i	0.3	0.2	0.3	0.7	0.4	0.4	0.3
		p	0.4	0.3	0.4	0.7	0.4	0.4	0.3
638	infection gen. syst. NEC ♂	i	0.4	0.7	0.9	0.7	0.5	0.6	0.6
		p	0.5	0.8	1.0	0.9	0.7	0.8	0.7
639	malign. neopl. prostate	i	0.0	0.1	0.1	0.1	0.0	0.1	0.1
		p	0.4	0.5	0.4	0.7	0.5	0.4	0.4
640	malign. neopl. testis	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.1	0.1	0.0	.	0.1	0.1	0.1
641	malign. neopl. gen. syst. NEC ♂	i	.	0.0	.	.	0.0	.	.
		p	0.0	0.0	.	.	0.0	0.0	0.1
642	benign. neopl. gen. syst. ♂	i	0.0	0.0	0.0	.	0.0	0.0	0.0
		p	0.0	0.0	0.0	.	0.0	0.0	0.0
643	unspecif. neopl. gen. syst. ♂	i	0.0	0.0	0.0	0.1	0.0	0.0	0.0
		p	0.0	0.0	0.0	0.1	0.0	0.0	0.0
644	unspecif. neopl. mamma ♂	i	0.0	0.1	0.0	.	0.0	0.0	0.0
		p	0.0	0.1	0.0	.	0.0	0.0	0.0
645	injury gen. syst./mamma ♂	i	0.0	0.1	0.1	0.1	0.0	0.1	0.1
		p	0.0	0.1	0.1	0.1	0.0	0.1	0.1

Y MALE GENITAL SYSTEM (INCLUDING BREAST)

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
621	i	0.3	0.2	0.2	0.2	0.2	0.2
	p	0.4	0.3	0.3	0.3	0.3	0.3
622	i	.	0.0	0.0	.	0.0	0.0
	p	.	0.0	0.0	.	0.0	0.0
623	i	0.0	0.1	0.0	0.0	0.0	0.1
	p	0.0	0.1	0.0	0.2	0.1	0.1
624	i	0.1	0.1	0.1	0.0	0.1	0.0
	p	0.1	0.1	0.2	0.1	0.2	0.0
625	i	0.1	0.0	0.0	0.0	0.0	0.0
	p	0.1	0.1	0.1	0.1	0.1	0.0
<hr/>							
626	i	0.1	0.0	0.1	0.0	0.1	0.1
	p	0.2	0.2	0.2	0.0	0.2	0.1
627	i	0.3	0.3	0.4	0.4	0.2	0.5
	p	0.3	0.4	0.5	0.4	0.3	0.6
628	i	0.1	0.1	0.0	0.1	0.0	0.1
	p	0.1	0.1	0.0	0.1	0.0	0.1
629	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.2	0.1	0.1	0.1	0.1	0.1
630	i
	p	.	.	0.0	.	0.0	.
<hr/>							
631	i	.	0.1	0.1	0.0	0.1	0.0
	p	0.0	0.1	0.1	0.0	0.1	0.1
632	i	0.0	0.0	0.0	.	0.0	.
	p	0.0	0.0	0.0	.	0.0	0.0
633	i	0.2	0.0	0.2	0.0	0.1	0.1
	p	0.2	0.0	0.2	0.0	0.1	0.1
634	i	0.1	0.0	0.0	0.0	0.1	0.0
	p	0.1	0.0	0.1	0.0	0.1	0.0
635	i	0.0	0.0	0.0	0.1	0.0	0.0
	p	0.1	0.0	0.1	0.1	0.0	0.1
<hr/>							
636	i	0.2	0.2	0.3	0.4	0.3	0.3
	p	0.6	0.3	0.5	0.6	0.5	0.5
637	i	0.3	0.4	0.3	0.4	0.3	0.4
	p	0.4	0.5	0.3	0.4	0.3	0.4
638	i	0.7	0.4	0.7	0.7	0.7	0.5
	p	0.8	0.7	0.8	0.7	0.8	0.6
639	i	0.0	0.1	0.1	0.0	0.0	0.1
	p	0.3	0.5	0.5	0.4	0.4	0.5
640	i	0.0	0.0	.	0.0	0.0	.
	p	0.1	0.1	0.1	0.0	0.1	0.0
<hr/>							
641	i	.	0.0	.	.	0.0	.
	p	0.0	0.0	.	0.1	0.0	.
642	i	0.0	0.0	.	.	0.0	0.0
	p	0.0	0.0	0.0	.	0.0	0.0
643	i	.	0.0	0.1	.	0.0	0.1
	p	.	0.0	0.1	0.0	0.0	0.1
644	i	0.1	0.0	0.0	0.0	0.0	0.1
	p	0.1	0.0	0.0	0.0	0.0	0.1
645	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.2	0.0	0.1	0.1	0.1	0.1

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
646	phimosi/redund. prepuce	i	0.2	0.2	0.2	0.1	0.2	0.2	0.2
		p	0.2	0.4	0.3	0.1	0.2	0.2	0.3
647	hypospadi	i
		p	0.0	0.0	.	.	.	0.0	0.0
648	undesc. testicle/cryptorchism	i	0.1	0.1	0.1	0.2	0.2	0.1	0.1
		p	0.2	0.1	0.2	0.3	0.3	0.1	0.2
649	cong. an. gen. syst./mamma NEC ♂	i	0.0	.	0.0	.	.	0.0	.
		p	0.0	0.0	0.0	.	0.1	0.0	0.0
650	benign prostatic hypertrophy	i	0.2	0.1	0.1	0.2	0.2	0.1	0.1
		p	0.4	0.4	0.5	0.5	0.6	0.5	0.4
651	hydrocele	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.2	0.1	0.1	0.1	0.2	0.1	0.1
652	dis. gen. syst./mamma NEC ♂	i	0.3	0.3	0.2	0.3	0.3	0.2	0.4
		p	0.5	0.4	0.4	0.3	0.5	0.3	0.6

Y MALE GENITAL SYSTEM (INCLUDING BREAST)

no.		season spring	summer	autumn	winter	type of insurance	
						public	private
646	i	0.3	0.1	0.1	0.2	0.2	0.2
	p	0.4	0.2	0.2	0.3	0.2	0.4
647	i
	p	0.0	0.0	0.0	0.0	0.0	.
648	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.1	0.1	0.2	0.2	0.2	0.1
649	i	0.0	.	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.0	0.0	0.0	0.0
650	i	0.1	0.2	0.2	0.1	0.1	0.2
	p	0.3	0.4	0.7	0.4	0.5	0.4
<hr/>							
651	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.1	0.1	0.1	0.1	0.2	0.1
652	i	0.3	0.2	0.3	0.2	0.3	0.3
	p	0.5	0.3	0.5	0.4	0.4	0.4

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
653	poverty/financial probl.	i	0.1	0.1	0.1	0.4	0.0	0.1	0.1
		p	0.1	0.3	0.3	0.6	0.1	0.3	0.4
654	probl. food/water	i	.	0.0	0.0
		p	.	0.0	.	.	.	0.0	0.0
655	probl. housing/neighborh.	i	0.2	0.3	0.4	0.3	0.2	0.3	0.2
		p	0.4	0.7	1.2	0.9	0.5	0.7	0.9
656	probl. house moving	i	0.1	0.1	0.1	0.1	0.1	0.1	0.1
		p	0.3	0.3	1.2	0.5	0.3	0.6	0.5
657	fear of having a social probl.	i	0.0	0.0	0.0	.	.	0.0	0.0
		p	0.0	0.0	0.1	0.1	0.0	0.0	0.0
658	probl. working conditions	i	0.1	0.3	0.1	0.3	0.1	0.2	0.1
		p	0.3	0.7	0.4	0.9	0.4	0.6	0.6
659	probl. job/employment NEC	i	0.5	0.6	0.8	0.8	0.6	0.6	0.6
		p	1.0	1.3	1.8	2.0	1.1	1.4	1.5
660	probl. w. being unemployed	i	0.0	0.0	0.1	.	0.1	0.0	0.1
		p	0.1	0.2	0.2	0.1	0.2	0.1	0.2
661	probl. w. education	i	0.2	0.1	0.4	0.2	0.3	0.2	0.3
		p	0.6	0.4	0.7	0.3	0.5	0.4	0.6
662	probl. soc. insurance/welfare	i	0.1	0.1	0.1	0.2	0.2	0.1	0.1
		p	0.3	0.3	0.4	0.8	0.4	0.3	0.5
663	probl. legal/police	i	0.0	0.0	.	.	0.0	0.0	0.0
		p	0.0	0.1	0.1	.	0.1	0.0	0.1
664	probl. health care syst./access	i	0.1	0.2	0.2	0.3	0.1	0.2	0.2
		p	0.2	0.5	0.5	0.6	0.3	0.4	0.5
665	probl. soc./cultural syst.	i	0.1	0.1	0.2	0.1	0.2	0.1	0.1
		p	0.2	0.4	0.3	1.4	0.4	0.4	0.4
666	relation problems partners	i	0.7	0.9	1.0	1.4	0.9	0.9	1.1
		p	1.5	2.1	1.9	3.0	1.6	2.0	2.0
667	problems w. behaving partner	i	0.1	0.1	0.0	0.1	0.1	0.1	0.0
		p	0.2	0.2	0.2	0.2	0.3	0.2	0.3
668	probl. w. partner being ill	i	0.3	0.6	0.4	0.5	0.5	0.4	0.6
		p	1.0	1.7	2.0	1.5	1.4	1.4	1.9
669	probl. loss of partner	i	0.4	0.6	0.8	1.3	0.6	0.7	0.6
		p	2.0	2.6	3.4	3.6	2.3	2.7	2.8
670	relation probl. with child	i	0.6	0.5	0.8	0.7	0.4	0.6	0.8
		p	1.1	1.1	1.3	1.0	0.7	1.0	1.8
671	probl. behaving child	i	0.1	0.2	0.1	0.1	0.0	0.2	0.1
		p	0.1	0.4	0.2	0.2	0.1	0.2	0.3
672	probl. with child being ill	i	0.1	0.1	0.1	.	0.2	0.1	0.1
		p	0.5	0.5	0.4	0.1	0.6	0.4	0.5
673	loss or death of child	i	0.1	0.1	0.2	0.1	0.1	0.1	0.2
		p	0.2	0.3	0.4	0.3	0.3	0.2	0.5
674	relation probl. parent/other family	i	0.7	0.6	0.8	0.3	0.6	0.6	0.7
		p	1.4	1.3	1.9	1.1	1.2	1.3	1.9
675	probl. w. behaving parent/other family	i	0.1	0.0	.	.	0.0	0.1	0.0
		p	0.1	0.1	0.0	.	0.0	0.1	0.1
676	probl. w. parent/other family being ill	i	0.2	0.2	0.2	0.1	0.3	0.2	0.3
		p	0.5	0.6	0.9	0.5	0.8	0.5	0.8
677	loss parent/family	i	0.4	0.3	0.5	0.4	0.4	0.3	0.4
		p	0.7	0.9	1.4	0.9	1.0	0.9	1.2

Z SOCIAL PROBLEMS

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
653	i	0.2	0.1	0.1	0.1	0.1	0.1
	p	0.4	0.2	0.3	0.2	0.3	0.2
654	i	.	.	.	0.0	0.0	.
	p	.	.	.	0.0	0.0	.
655	i	0.3	0.2	0.3	0.2	0.3	0.1
	p	0.7	0.5	1.1	0.6	0.9	0.4
656	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.4	0.5	0.7	0.6	0.6	0.3
657	i	.	.	0.0	0.0	0.0	0.0
	p	0.0	0.0	0.1	0.0	0.0	0.0
658	i	0.3	0.1	0.1	0.2	0.2	0.2
	p	0.6	0.4	0.5	0.6	0.5	0.6
659	i	0.6	0.4	0.7	0.8	0.7	0.5
	p	1.3	0.8	1.7	1.7	1.5	1.1
660	i	0.0	0.0	0.1	0.1	0.1	0.0
	p	0.1	0.1	0.2	0.2	0.2	0.1
661	i	0.3	0.1	0.2	0.4	0.2	0.3
	p	0.6	0.3	0.5	0.6	0.5	0.5
662	i	0.1	0.1	0.1	0.1	0.1	0.1
	p	0.3	0.3	0.5	0.5	0.4	0.2
663	i	0.0	.	.	0.0	0.0	.
	p	0.0	0.0	0.1	0.1	0.1	0.0
664	i	0.1	0.2	0.2	0.3	0.2	0.2
	p	0.2	0.3	0.5	0.7	0.4	0.3
665	i	0.1	0.0	0.1	0.2	0.1	0.0
	p	0.3	0.5	0.4	0.5	0.6	0.1
666	i	1.0	0.6	1.0	1.1	0.9	0.9
	p	2.4	1.4	1.9	2.1	2.0	1.6
667	i	0.0	0.1	0.1	0.0	0.1	0.1
	p	0.2	0.2	0.2	0.2	0.2	0.1
668	i	0.3	0.4	0.5	0.6	0.4	0.6
	p	1.3	1.4	1.6	1.8	1.4	1.7
669	i	0.6	0.4	1.0	0.7	0.7	0.5
	p	3.0	2.1	3.1	2.6	2.9	1.9
670	i	0.5	0.8	0.5	0.7	0.8	0.4
	p	1.1	1.1	1.1	1.3	1.4	0.7
671	i	0.2	0.0	0.1	0.1	0.1	0.1
	p	0.3	0.1	0.2	0.2	0.2	0.2
672	i	0.1	0.0	0.1	0.1	0.1	0.1
	p	0.4	0.3	0.5	0.5	0.4	0.3
673	i	0.1	0.1	0.2	0.1	0.1	0.1
	p	0.2	0.2	0.4	0.3	0.3	0.3
674	i	0.4	0.6	0.6	0.9	0.7	0.4
	p	1.2	1.1	1.5	2.0	1.6	0.9
675	i	0.0	0.1	0.0	0.0	0.1	0.0
	p	0.1	0.1	0.0	0.1	0.1	0.0
676	i	0.1	0.2	0.3	0.2	0.2	0.2
	p	0.4	0.5	0.8	0.8	0.7	0.5
677	i	0.3	0.3	0.6	0.3	0.4	0.3
	p	0.6	0.8	1.4	1.1	1.1	0.7

**Incidence and prevalence of morbidity in general practice in a 3 months period:
according to degree of urbanization, region, season and type of insurance**

no.	episode		degree of urbanization				region		
			rural	suburban	urban	city	north	centre	south
678	loss or death of friend	i	0.0	0.0	0.0	0.1	0.1	0.0	0.0
		p	0.0	0.1	0.1	0.2	0.1	0.1	0.1
679	relation probl. w. friend	i	0.1	0.1	0.1	.	0.0	0.1	0.0
		p	0.1	0.1	0.1	.	0.0	0.1	0.1
680	probl. assault/harmfull events	i	0.1	0.1	0.2	0.2	0.0	0.1	0.2
		p	0.2	0.2	0.6	0.4	0.2	0.3	0.4
681	probl. with being ill	i	0.0	0.0	0.0	0.1	0.1	0.0	0.1
		p	0.1	0.1	0.5	0.2	0.1	0.2	0.2
682	social handicap	i	0.1	0.0	0.0	0.1	0.1	0.0	0.1
		p	0.2	0.2	0.2	0.1	0.2	0.2	0.2
683	social isolation	i	0.1	0.3	0.1	0.4	0.1	0.2	0.1
		p	0.7	1.1	0.8	1.9	0.7	1.1	0.7
684	social problems	i	0.2	0.2	0.2	0.8	0.2	0.3	0.2
		p	0.5	0.9	0.5	3.0	0.7	1.0	0.7

Z SOCIAL PROBLEMS

no.		season				type of insurance	
		spring	summer	autumn	winter	public	private
678	i	.	0.0	0.0	0.1	0.0	0.0
	p	.	0.1	0.1	0.2	0.1	0.1
679	i	0.0	0.0	0.0	0.1	0.1	0.0
	p	0.0	0.1	0.1	0.2	0.1	0.0
680	i	0.1	0.1	0.2	0.1	0.2	0.0
	p	0.3	0.1	0.5	0.2	0.4	0.2
681	i	.	0.0	0.1	0.1	0.0	0.0
	p	.	0.1	0.2	0.5	0.2	0.2
682	i	0.0	0.1	0.0	0.1	0.0	0.1
	p	0.1	0.2	0.2	0.1	0.2	0.1
683	i	0.1	0.2	0.1	0.2	0.2	0.2
	p	0.9	1.4	0.8	0.9	1.0	0.8
684	i	0.3	0.1	0.4	0.2	0.3	0.1
	p	1.0	0.6	1.3	0.7	1.1	0.5

APPENDIX B Diagnosis(-numbers) used in tables related to the ICPC-codes from the NIVEL-edition (1989) and the Oxford-edition (1987)

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
A	GENERAL AND UNSPECIFIED		
1	pain: generalized/unspec.	A010	A01
2	chills/fever		
	cold (chills)	A020	A02
	fever	A030	A03
3	general weak-/tiredness	A040	A04
4	fainting (syncope)	A060	A06
5	symptoms infant		
	infantile colic	A111	A14
	excess. crying infant	A112	A15
	irrit. infant/sympt. NEC	A119	A16/A17
6	allergy NEC	A120	A12
7	feeling/behaving old/senile	A140	P05
8	fear of disease/death		
	fear of medical treatment	A240	-
	fear of death	A250	A25
	fear of cancer NEC	A260	A26
	fear of disease NEC	A270	A27
9	disability/impairment NOS	A28	A28
10	general sympt./complt. NEC		
	general deterioration	A050	A05
	coma	A070	A07
	swelling NEC	A080	A08
	sweating problems	A090	A09
	bleeding, site NOS	A100	A10
	concern about drug reaction	A130	A13
	euthanasia by request	A150	A20
	concern about health family	A230	-65
	general sympt./complt. NEC	A290	A29
11	tuberculosis, general, excl. lung	A700	A70
12	measles	A710	A71
13	chickenpox	A720	A72
14	malaria	A730	A73
15	rubella	A740	A74
16	infectious mononucleosis	A750	A75
17	viral dis. with exanthems NEC	A760	A76
18	viral dis. NOS/NEC	A770	A77
19	infectious dis. NOS/NEC	A780	A78
20	carcinomatosis (prim. site unkn)	A790	A79
21	accident/injury NOS	A800	A80
22	mult. trauma/internal injury	A810	A81
23	late effects of trauma	A820	A82
24	poisoning by/abuse of med. agents	A830	A84/P18
25	poisoning by benzodiazepines	A841	A84
26	poisoning by med. agents NOS	A849	A84
27	adv. eff. med. agent proper dose: allergy	A851	A85/S88
28	adv. eff. OAC	A852	A85
29	adv. eff. med. agent NEC	A859	A85
30	toxic eff. other substances	A860	A86
31	complic. surg./med. treatm.: hematoma	A871	A87
32	complic. surg./med. treatm.: wounds	A872	A87
33	complic. surg./med. treatm.: NEC	A879	A87

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
34	adv. eff. physical factors NEC		
	motion sickness	A881	A88
	adv. eff. physical factors NEC	A889	A88
35	adv. eff. prosthetic device	A890	A89
36	mult.congenital syndromes	A900	A90
37	abnormal results investigation NEC	A910	A91
38	toxoplasmosis	A920	A92
39	premat./immature liveborn infant	A930	A93
40	all perinatal morbidity		
	icterus neonatorum	A941	A94
	aphyxia neonatorum/RDS	A942	A94
	dyspepsia/diarrhoea neonatorum	A943	A94
	conjunctivitis neonatorum	A944	A94
	anomalies perinatal period NEC	A949	A94
41	perinatal mortality	A950	A95
42	death (excl. perinatal)	A960	A96
43	no disease	A970	A97
44	disease NEC	A990	A99
B	BLOOD, BLOODFORM. ORG.,LYMPHSYSTEM AND SPLEEN		
45	enlarged lymph gland(s)	B020	B02
46	sympt. lymph glands(s) NEC	B030	B03
47	fear of blood/lymph disease		
	fear of AIDS	B250	B25
	fear of cancer blood/blood org.	B260	B26
	fear of other blood/lymph dis.	B270	B27
48	disability/impairment blood	B280	B28
49	sympt./complt. blood NOS/NEC		
	sympt. blood/bloodform. organs	B040	B04
	sympt./complt. blood/lymph/spleen	B290	B29
50	acute lymphadenitis/-angitis	B700	B70
51	chron. aspec. lymphadenitis	B710	B71
52	malignant lymphoma	B721	B72
53	leukemia	B722	B73
54	malignant neopl. blood NEC	B730	B74
55	benigne neopl. blood/bloodform. org.	B740	B75
56	ruptured spleen	B760	B76
57	injuries blood/bloodform. org. NEC	B770	B77
58	hered. haemolytic anaemias	B780	B78
59	cong. anom. blood/bloodform. org. NEC	B790	B79
60	iron deficiency anemia	B800	B80
61	pernic./other deficiency anemia		
	pernic. anemia	B811	B81
	folate deficiency anemia	B812	B81
	deficiency anemia NEC	B819	B82
62	anemia NOS/NEC	B820	B82
63	hemophilia/coagulation disorders		
	hemophilia	B831	B83
	purpura/abn. plate/coag. dis. NOS	B839	B83
64	abnormal white cells	B840	B84
65	unexplained abn. blood test	B850	B85
66	abn. blood test NOS	B860	B86
67	splenomegaly	B870	B87
68	HIV-infection (AIDS/ARS)	B900	B90
69	diseases blood/bloodform. org. NOS/NEC	B990	B99

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
D	DIGESTIVE SYSTEM		
70	generalized abd. pain/cramps	D010	D01
71	stomach pain	D020	D02
72	heartburn	D030	D03
73	peri-anal itching	D050	D05
74	localised abd. pain	D060	D06
75	flatulence/belching	D080	D08
76	nausea	D090	D09
77	vomiting	D100	D10
78	diarrhoea	D110	D11
79	constipation	D120	D12
80	jaundice	D130	D13
81	blood vomiting	D140	D14
82	melena	D150	D15
83	rectal bleeding	D160	D16
84	fecal incontinence	D170	D17
85	defecation anomalies NEC	D180	D18
86	sympt./complt. of teath, gums	D190	D19
87	sympt./complt. mouth, tongue, lip	D200	D20
88	swallowing problems	D210	D21
89	fear of digestive diseases		
	fear of cancer of dig. system	D260	D26
	fear of other digestive diseases	D270	D27
90	digestive disability/impairment	D280	D28
91	sympt./complt. digestive NEC		
	perianal itching	D040	D04
	indigestion	D070	D29/T29
	localised abd. mass NOS	D240	D24
	ascites	D251	D25
	change in abd. size/distension NEC	D259	D25
	sympt./complt. digest. NEC	D290	D29
92	infectious diarrhoea/dysentery	D700	D70
93	mumps		D710D71
94	infectious hepatitis		
	hepatitis A	D721	D72
	hepatitis B	D722	D72
	hepatitis non A non B	D723	D72
95	parasites/worms	D730	D22
96	other presumed infections	D740	D73
97	malign. neopl. stomach	D751	D74
98	malign. neopl. colon, rectum	D752	D75
99	malign. neopl. colon, rectum	D753	D75
100	malign. neopl. pancreas	D761	D76
101	malign. neopl. dig. syst. NEC		
	malign. neopl.stomach/intestine NEC	D759	D77
	malign. neopl. liver/bileduct NEC	D769	D77
102	benign. neoplasms dig. syst. NEC	D770	D78
103	neoplasms dig. syst. NEC	D780	D77
104	foreign body through orifice	D790	D79
105	injuries dig. syst. NEC	D800	D80
106	cong. anom. digest. system	D810	D81
107	disease of teeth/gums	D820	D82
108	disease of mouth/tongue/lips	D830	D83
109	disease of esophagus	D840	D84
110	duodenal ulcer	D850	D85
111	gastric ulcers NEC	D861	D86
112	peptic ulcers NEC	D869	D86

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
113	disord. stomach function	D870	D87
114	appendicitis	D880	D88
115	inguinal hernia	D890	D89
116	hiatus (diaphr.) hernia	D900	D90
117	abdominal hernia NEC	D910	D91
118	diverticular dis. intestines	D920	D92
119	irritable bowel syndrome	D930	D93
120	chron. enteritis/colitis ulc.	D940	D94
121	anal fissure/perianal abscess		
	anal fissure	D951	D95
	anal fist.	D952	D95
	perianal abscess	D953	D95
122	hepatomegaly	D960	D96
123	cirrhosis/liver disease NEC		
	livercirrhosis	D971	D97
	chron. hepatitis NEC	D979	D97
124	diseases of gall-bladder/biliary duct		
	cholecystitis	D981	D98
	cholelithiasis	D982	D98
	diseases bile duct/-bladder NEC	D989	D99
125	diseases digestive system NEC		
	anomalies pancreas NEC	D991	D99
	adhesions	D992	D99
	(sub)acute diseases of bowel	D993	D99
	diseases digestive system NEC	D999	D99
F	EYE		
126	complaints of vision	F050	F05
127	fear of eye disease	F270	F27
128	disability/impairment eye	F280	F28
129	sympt./complt. eye NEC		
	eye pain	F010	F01
	red eye	F020	F02
	discharge from eye	F030	F03
	floaters/spots	F040	F04
	abn. sensations of eye	F130	F13
	abn. eye movements	F140	F14
	abn. appearance of eyes	F150	F15
	symptoms of eyelids	F160	F16
	sympt./complt. glasses	F170	F17
	sympt./complt. contact lens	F180	F18
	sympt./complt. eye NEC	F290	F29
130	infectious conjunctivitis		
	inf. conjunctivitis: viral NOS	F701	F70
	inf. conjunctivitis: bact. NOS	F702	F70
	inf. conjunctivitis NEC/Trachoma	F709	F70/F86
131	allergic conjunctivitis	F710	F71
132	blepharitis/stye/chalazion	F720	F72
133	infection of eye NEC		
	iritidocyclitis	F731	F73
	keratitis herpetica	F732	F85
	infection of eye NEC	F739	F73
134	corneal ulcer	F733	F85
135	neoplasms of eye/adnexa	F740	F74
136	contusion/abrasion/black eye	F750	F75
137	foreign body in eye	F760	F76
138	injuries eye NEC	F790	F79

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
139	blocked lacrimal duct of infants	F800	F80
140	cong. anom. of eye NEC	F810	F81
141	refractive errors	F910	F91
142	cataract	F920	F92
143	glaucoma	F930	F93
144	blindness, all types	F940	F94
145	strabismus	F991	F95
146	retinopathy	F992	F82/F83/F84
147	diseases of eye NEC		
	ectropion/entropion	F993	F99
	diseases of eye NEC	F999	F99
H	EAR		
148	earache	H010	H01
149	hearing complt.	H020	H02
150	ringing/buzzing/tinnitus	H030	H03
151	blood/discharge from ear		
	discharge from ear	H040	H04
	blood in/from ear	H050	H05
152	fear of ear disease	H270	H27
153	disability/impairment of ear	H280	H28
154	symp./complt. of ear NEC		
	plugged feeling	H130	H13
	concern with appear. of ears	H150	H15
	symp./complt. of ear NEC	H290	H29
155	otitis externa	H700	H70
156	acute otitis media/myringitis	H710	H71
157	serous otitis media/glue ear	H720	H72
158	eustachian tube dysfunction	H730	H73
159	infections of ear NEC		
	cholesteatoma	H741	H74
	chronic supp. otitis media	H742	H74
	mastoiditis	H743	H74
	infections of ear NEC	H749	H74
160	neoplasm of ear	H750	H75
161	foreign body in ear	H760	H76
162	perforation tymp. membrane	H770	H77
163	superficial injury of ear	H780	H78
164	injuries of ear NEC	H790	H79
165	cong. anom. of ear	H800	H80
166	ear wax (excessive)	H810	H81
167	vertiginous syndromes		
	M. Ménière	H821	H82
	otogenous vert. synndr. NEC	H829	H82
168	otosclerosis	H830	H83
169	presbycusis	H841	H84
170	acoustic trauma	H842	H85
171	deafness NEC	H849	H86
172	disease of ear/mastoid NEC	H990	H99
K	CIRCULATORY SYSTEM		
173	complaints of heart		
	pain attributed to heart	K010	K01
	pressure tightness	K020	K02
	palpitations	K040	K04
	abnorm./irreg. heartbeat/pulse	K050	K05

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
174	edema of extremities	K070	K07
175	fear of dis. circ. system		
	fear of heart attack	K240	K24
	fear of hypertension	K250	K25
	fear of dis. heart/circ. syst. NEC	K270	K27
176	disability/impairment circ. syst. NEC	K280	K28
177	sympt./complt. circ. syst. NEC		
	pain attributed to bloodvessels	K030	K03
	prominent veins	K060	K06
	sympt./complt. circ. syst. NEC	K290	K29
178	infections dis. circ. syst.	K700	K70
179	neoplasm circ. syst.	K710	K72
180	injuries heart/circ. syst.	K720	K99/K84
181	cong. anom. heart/circ. syst.	K730	K73
182	rheumatic fever/heart dis.	K740	K71
183	acute myocardial infarction	K751	K75
184	angina pectoris		
	imminent heart attack	K752	K75
	angina pectoris	K761	K74
185	healed myocardial infarction	K762	K76
186	chron. ischemic heart dis. NEC		
	coronary sclerosis	K763	K76
	chron. isch. heart dis. NEC	K769	K76
187	heart failure	K770	K77
188	atrial fibrillation/flutter	K780	K78
189	paroxysmal tachycardia	K790	K79
190	ectopic beats, all types	K800	K80
191	heart murmur NOS	K810	K81
192	pulmonary heart dis.	K820	K82
193	heart valve dis. NOS, non-rheum.	K830	K83
194	heart disease NEC		
	atrio ventr./bundle branch block	K841	K84
	cardiomyopathy	K842	K84
	dis. of heart rhythm NEC	K843	K84
	heart diseases NEC	K849	K84
195	elevated blood pressure w.o. hypertension	K850	K85
196	uncomplicated hypertension	K860	K86
197	hypertension with involv.targ. organs	K870	K87
198	(postural) hypotension	K880	K88
199	transient cerebral ischemia	K890	K89
200	cerebrovascular diseases		
	cerebrovasculair accident	K901	K90
	subarachnoid hemorrhage	K902	K90
	cerebrovascular dis. NEC	K909	K92
201	atherosclerosis, excl. heart/brain	K910	K91
202	arterial vascular diseases NEC		
	intermittent claudication	K921	K92
	art. obstr./periph. art. vasc. dis. NEC	K929	K92
203	pulmonary embolism	K930	K93
204	(thrombo) phlebitis	K941	K94
205	venous thrombosis	K942	K94
206	various veins of legs	K951	K95
207	various ulcer of skin	K952	S97
208	hemorrhoids	K960	K96
209	dis. circulatory system NEC		
	(postmastectomy) lumphedema	K991	K99
	dis. circulatory system NEC	K999	K99

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
L	MUSCULOSKELETAL SYSTEM		
210	neck symp./complt., excl. headache	L010	L01
211	back symp./complt.	L020	L02
212	low back symp./complt. w/o radiation	L030	L03
213	chest sympt./complt.	L040	L04
214	flank sympt./complt.	L050	L05
215	axilla sympt./complt.	L060	L06
216	jaw sympt./complt.	L070	L07
217	shoulder sympt./complt.	L080	L08
218	arm sympt./complt.	L090	L09
219	elbow sympt./complt.	L100	L10
220	wrist sympt./complt.	L110	L11
221	hand/finger sympt./complt.	L120	L12
222	hip sympt./complt.	L130	L13
223	leg/thigh sympt./complt.	L140	L14
224	knee sympt./complt.	L150	L15
225	ankle sympt./complt.	L160	L16
226	foot/toe sympt./complt.	L170	L17
227	muscle pain/fibrositis	L180	L18
228	(mult.) muscle sympt./complt. NEC	L190	L19
229	(mult.) joints sympt./complt. NEC	L200	L20
230	limbs sympt./complt. NOS	L250	L29
231	fear of dis. musculoskeletal syst.		
	fear of cancer musc. skeletal syst.	L260	L26
	fear of musculoskeletal dis. NEC	L270	L27
232	disability/impairment musc. skel. syst.	L280	L28
233	symp./complt. musc. skel. syst.	L290	L29
234	infections musc. skel. syst.		
	Bornholm disease	L701	L70
	osteomyelitis/osteitis pyog./abscess	L702	L70
	infections musc. skel. syst. NEC	L703	L70
235	neoplasms musc. skel. syst.		
	benign. neopl. musc. skel. syst.	L711	L71
	malign. neopl. musc. skel. syst.	L712	L71
	unspecified neopl. musc. skel. syst.	L713	L71
236	fracture: radius/ulna	L720	L72
237	fracture: tibia/fibula	L730	L73
238	fracture: hand/foot bones	L740	L74
239	fracture: collum femoris	L751	L75
240	fracture: femur NEC	L759	L75
241	fracture: skull	L761	L76
242	fracture: facial bones	L762	L76
243	fracture: spine	L763	L76
244	fracture: rib	L764	L76
245	fracture: clavicula	L765	L76
246	fracture: humerus	L766	L76
247	fracture: pelvis	L767	L76
248	fracture: NEC	L769	L76
249	sprains/strains ankle(s)	L770	L77
250	sprains/strains knee(s)	L780	L78
251	sprains/strains shoulder/arm	L791	L79
252	sprains/strains wrist/hand	L792	L79
253	sprains/strains foot/toe	L793	L79
254	sprains/strains spine, excl. cerv. spine	L794	L79
255	sprains/strains cerv. spine	L795	L79
256	whiplash calf	L796	L81
257	sprains/strains musc. skel. syst. NEC	L799	L79/L81

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
258	(sub)luxation, dislocation		
	(sub)luxation/dislocation shoulder	L801	L80
	(sub)luxation/dislocation jaw	L802	L80
	(sub)luxations/dislocation NEC	L809	L80
259	cong. luxation of hip	L821	L82
260	cong. anom. musc. skel. syst. NEC	L829	L82
261	syndromes cervical spine	L830	L83
262	osteoarthritis of cervical spine	L841	L84
263	osteoarthritis of spine NEC	L842	L84
264	acquired deformities of spine	L850	L85
265	back pain with radiating sympt.	L861	L86
266	lumbago	L862	L03
267	sciatica	L863	L86
268	slipped lumbar disc	L864	L86
269	ganglion joint/tendon	L870	L87
270	rheumatoid arthritis	L881	L88
271	spondylitis (ankylosing)	L882	L88
272	osteoarthritis hip	L891	L89
273	osteoarthritis knee	L892	L90
274	osteoarthritis NEC	L899	L91
275	arthritis in allied conditions NEC	L910	L99
276	shoulder syndrome	L920	L92
277	dis. bursa, excl. shoulder	L931	L99
278	epicondylitis lateral	L932	L93
279	tendinitis/synovitis	L939	L99
280	osteochondrosis	L940	L94
281	osteoporosis	L950	L95
282	ac. traum. meniscus/lig. knee	L960	L96
283	retropatellar chondropathy	L971	L97
284	chron. internal knee derangement	L979	L97
285	acquired deformities of limbs		
	pes planus	L981	L98
	hallux valgus	L982	L98
	genua valga/vara	L983	L98
	acquired deformities limbs NEC	L989	L98
286	dis. musc. skel. system NEC	L990	L99
N	NEUROLOGICAL SYSTEM		
287	headache/facial pain		
	headache	N010	N01
	pain facial	N020	N03
288	sensibility disturbance		
	tingling fingers/feet/toes	N030	N05
	sens. dist./abn. invol. movements	N050	N06
289	restless legs syndrome	N040	N04
290	convulsions	N070	N07
291	symptoms sense/smell/taste	N160	N16
292	vertigo	N170	N17
293	paralysis	N180	N18
294	sympt./complt. speech	N190	N19
295	fear of neurological disease		
	fear of cancer neurol. syst.	N260	N26
	fear of other neurological disease	N270	N27
296	disability/impairment neurol. syst.	N280	N28
297	sympt./complt. neurol. syst. NEC	N290	N29
298	poliomyelitis	N700	N70

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
299	meningitis/encefalitis	N710	N71
300	tetanus	N720	N72
301	infections neur. syst. NEC	N730	N73
302	malignant neoplasm neur. syst. mal. neopl. brains	N741	N74
	mal. neopl. neur. syst. NEC	N749	N74
303	benigne neopl. neur. syst.	N750	N75
304	unspecified neopl. neur. syst.	N760	N76
305	concussion: commotio cer.	N791	N79
306	concussion: contusio cer.	N792	N79
307	head injury NOS/NEC, w/o skull fract.	N800	N80
308	injuries neur. syst. NEC	N810	N81
309	cong. anom. neur. syst.	N850	N85
310	multiple sclerosis	N860	N86
311	Parkinsonism	N870	N87
312	epilepsy, all types	N880	N88
313	migraine	N890	N89
314	facial paralysis, Bell's palsy	N991	N91
315	trigeminal neuralgia	N992	N92
316	carpal tunnel syndrome	N993	N93
317	cluster headache	N994	N90
318	dis. of periph. nerv. syst. NEC	N998	N94
319	dis. of central nerv. syst. NEC	N999	N99
P	PSYCHOLOGICAL SYSTEM		
320	feeling anxious/nervous/tense	P010	P01
321	acute stress (transient) reaction	P020	P02
322	feeling depressed	P030	P03
323	feeling/behaving irritable irritated/angry feeling/behaving feeling restless/agitated	P040 P050	P04 P29
324	suicide attempt	P060	P77
325	disturbances of sleep	P070	P06
326	sexual problems	P080	P07/P08
327	hypochondr. dis./probl. with being ill	P090	P75/Z11
328	tension headache	P100	N02
329	hyperventilation	P110	R98
330	enuresis	P120	P12
331	anorexia nervosa/boulimia	P131	T06
332	eating/feeding probl. NEC	P139	P11/T04/T05/T29
333	chronic alcohol abuse	P150	P15
334	acute alcohol abuse	P160	P16
335	tobacco abuse	P170	P17
336	drug abuse	P181	P19
337	medicinal abuse	P189	P18
338	abuse substances NEC	P190	P19
339	dist. memory/concentration	P200	P20
340	disturbance behaviour child encopresis overactive child, hyperkin. syndr. Minimal Brain Dysfunction sympt./complt. behaviour child	P221 P222 P223 P229	P13 P21 P22 P22
341	sympt./complt. behaviour adolescent	P260	P23
342	sympt./complt. adult	P230	P80
343	phase of live problem adult	P240	P25
344	dist. learning/mental development	P250	P24
345	fear of mental disorder	P270	P27

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
346	psychic disability/impairment	P280	P28
347	psychic sympt./complt. NEC	P290	P29
348	dementia (pre)senile		
	presenile dementia Alzheimer	N995	P70
	dementia senile	P701	P70
349	organic psychosis		
	puerperal psychosis	P702	P71
	organic psychosis NEC	P709	P71
350	schizophrenia, all types	P710	P72
351	affective psychosis		
	endogenous unipolar depression	P721	P73
	manic depressive psychosis	P722	P73
	affective psychosis NEC	P729	P73
352	unspecified psychosis NEC	P730	P98
353	anxiety disorder/anxiety state	P740	P74
354	hysterical/hypochondriacal dis.		
	hysteria	P751	P75
	hypochondriasis	P752	P75
355	depressive disorder	P760	P76
356	mental retardation	P770	P85
357	neurasthenia, surmenage	P780	P78
358	personality disorder	P800	P80
359	phobia/neurotic disorder NEC		
	phobia	P991	P79
	neurotic disorders NEC	P998	P79
360	mental/psychological dis. NEC	P999	P99
R	RESPIRATORY SYSTEM		
361	pain attr. to respiratory system	R010	R01
362	shortness of breath, dyspnoe	R020	R02
363	wheezing	R030	R03
364	breathing problems NEC	R040	R04
365	cough	R050	R05
366	epistaxis/nose bleed	R060	R06
367	sneezing/nasal congestion	R070	R07
368	sympt./complt. nose NEC	R080	R08
369	sympt./complt. sinus (incl. pain)	R090	R09
370	sympt./complt. throat	R210	R21
371	sympt./complt. tonsils	R220	R22
372	sympt./complt. voice/vocal cords	R230	R23
373	haemoptysis	R240	R24
374	abnormal sputum/phlegm NEC	R250	R25
375	fear of respiratory disease		
	fear of cancer of resp. syst.	R260	R26
	fear of respiratory dis. NEC	R270	R27
376	disability/impairment resp. syst.	R280	R28
377	sympt./complt. resp. syst. NEC		
	cyanosis	R291	R29
	hiccup	R292	R29
	sympt./complt. resp. syst. NEC	R299	R29
378	tuberculosis resp. (excl. A70)	R700	R70
379	whooping cough	R710	R71
380	scarlet fever	R721	R72
381	streptoc. inf. throat NEC	R729	R72
382	boil/abscess nose	R730	R73
383	URI (head cold)	R740	R74
384	sinusitis acute/chronic	R750	R75

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
385	acute tonsillitis/adenoiditis	R760	R76
386	acute laryngitis/tracheitis/croup	R770	R77
387	acute bronchi(oli)tis	R780	R78
388	influenza w/o pneumonia	R800	R80
389	pneumonia	R810	R81
390	pleurisy all (excl. R70)	R820	R82
391	infection resp. syst. NEC	R830	R83
392	mal. neopl. bronchus/lung	R841	R84
393	mal. neopl. resp. syst. NEC	R849	R85
394	ben. neopl. resp. syst.	R850	R86
395	unspecified neopl. resp. syst.	R860	R85
396	foreign body resp. syst.	R870	R87
397	injuries resp. syst. NEC	R880	R88
398	cong. anom. resp. syst.	R890	R89
399	hypertrophy/chron. inf. tonsills/aden.	R900	R90
400	chronic bronchitis		
	chronic bronchitis/bronchiectasis	R911	R91
	COPD	R912	R95
401	pleural effusion NOS	R930	R93
402	emfysema	R950	R95
403	asthma	R960	R96
404	allergic rhinitis/hay fever	R970	R97
405	diseases resp. syst. NEC		
	deviation nasal septum	R991	R99
	polyp of nose	R992	R99
	vocal cord nodes	R993	R85/R86
	disorder resp. syst. NEC	R999	R99
S SKIN/SUBCUTANEOUS TISSUE			
406	pain/tenderness of skin	S010	S01
407	pruritus, skin itching	S020	S02
408	warts	S031	S03
409	condyloma acuminata	S032	X91/Y76
410	local swelling/papul/lump	S040	S04
411	multiple swelling/papul/lump	S050	S05
412	local redness/erythema	S060	S06
413	general redness/erythema	S070	S07
414	changes in skin color NEC	S080	S08
415	local infection finger/toe	S090	S09
416	boil/carbuncle/cellulitis	S100	S10
417	local infection skin NEC	S110	S11
418	insect bite	S120	S12
419	animal/human bite	S130	S13
420	burns/scalds	S140	S14
421	foreign body in skin	S150	S15
422	bruise/contusion intact skin	S160	S16
423	abrasion/scratch/blister	S170	S17
424	laceration/cut	S180	S18
425	injury to skin NEC	S190	S19
426	corns/callosities	S200	S20
427	sympt./complt. of skin texture	S210	S21
428	sympt./complt. nails	S220	S22
429	losing hair/baldness	S231	S23
430	sympt./complt. hair NEC		
	hirsutismus	S232	S24
	sympt./complt. hair NEC	S239	S24

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
431	fear of skin disease		
	fear of cancer of skin	S260	S26
	fear of diseases of skin NEC	S270	S27
432	disability/impairment of skin	S280	S28
433	sympt./complt. of skin NEC	S290	S29
434	herpes zoster	S700	S70
435	herpes simplex lip/mouth	S711	S71
436	herpes simplex genitalis	S712	X90/Y72
437	herpes simplex NEC	S719	S71
438	scabies/other ascariasis	S720	S72
439	pediculosis/skininfestation	S730	S73
440	dermatophytosis/dermatomycosis		
	athlete's foot	S741	S74
	pityriasis versicolor	S742	S74
	onychomycosis	S743	S74
	dermatophytosis/-mycosis NEC	S749	S74
441	moniliasis/candidiasis (excl. GU)	S750	S74
442	erysipelas	S761	S76
443	mollusca contagiosa	S762	S95
444	infection skin NEC	S769	S76
445	malign. neoplasm skin	S770	S77
446	lipoma	S780	S78
447	benign. neoplasm skin NEC	S790	S79
448	unspecified neoplasm skin	S800	S80
449	hemangioma/lymfangioma	S810	S81
450	nevus/mole	S820	S82
451	cong. anom. of skin NEC	S830	S83
452	impetigo	S840	S84
453	pilonidal cyst/fistula	S850	S85
454	seborrheic dermatitis	S860	S86
455	atopic dermatitis/eczema	S870	S87
456	contact dermatitis	S881	S88
457	eczema NEC	S882	S88
458	diaper rash	S890	S89
459	pityriasis rosea	S900	S90
460	psoriasis w/wo arthropathy	S910	S91
461	pompholyx/dis. sweat glands	S920	S92
462	sebaceous cyst	S930	S93
463	ingrown toenail/dis. of nail NEC	S940	S94
464	alopecia/folliculitis/dis. of hair NEC	S950	S23/S11/S24
465	acne vulgaris	S960	S96
466	chron. ulcer skin, incl. varicose	S970	S97 (incl. ulcus cruris)
467	urticaria	S980	S98
468	seborrheic warts	S991	S99
469	acne rosacea	S992	S99
470	dis. skin/subcut. tissue NEC	S999	S99
T	ENDOCRINE/METABOLIC/NUTRITIONAL SYSTEM		
471	excessive thirst	T010	T01
472	excessive appetite, excl. boulimia	T020	T02
473	loss of appetite, excl. anorexia nerv.	T030	T03
474	feeding problem infant, excl. P11	T040	T04
475	feeding problem adult, excl. T06	T050	T05
476	feeding problem NEC	T060	T06
477	weight gain, excl. T82	T070	T07
478	weight loss, excl. T06	T080	T08
479	weight problem NEC	T090	T07/T08

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
480	failure to thrive	T100	T10
481	dehydration	T110	T11
482	thyroid lump/mass	T120	T15
483	fear of endocr. metab. nutr. dis.		
	fear of cancer of endocr. syst.	T260	T26
	fear of end. met. nutr. dis. NEC	T270	T27
484	disability/impairment end. gl.	T280	T28
485	sympt./complt. end./met./nutr. NEC	T290	T29
486	infectious dis. endocrine glands	T700	T70
487	malign. neopl. thyroid	T710	T71
488	benign. neopl. thyroid	T720	T72
489	unspecified neopl. endocr. NEC	T730	T73
490	injury thyroid	T740	T74
491	thyroglossal duct(cyst)	T780	T78
492	cong. anom. endocr. NEC	T800	T80
493	goiter/module w/o thyrotox	T810	T81
494	obesity/overweight	T820	T82/T83
495	hyperthyroidism/thyrototoxicosis	T850	T85
496	hypothyroidism/myxedema	T860	T86
497	hypoglycemia	T870	T87
498	renal glucosuria	T880	T88
499	diabetes mellitus	T900	T90
500	vitamin defic./nutr. dis. NEC	T910	T91
501	gout	T920	T92
502	lipid metabolism dis.	T930	T93
503	dis. thyroid NEC	T991	T99
504	dis. endocr. gland NEC	T992	T99
505	metabolic disease NEC	T993	T99
U	URINARY SYSTEM		
506	complaints urination		
	painful urination, dysuria	U010	U01
	frequent/urgent urination	U020	U02
	retention of urine	U051	U05
	urination problems NEC	U059	U05
507	incontinence (excl. P12)	U040	U04
508	anuria/oliguria	U052	U05
509	complaints of urine		
	blood in urine	U060	U06
	complaints urine NEC	U070	U07
510	sympt./complt. bladder NEC	U130	U13
511	sympt./complt. kidney	U140	U14
512	fear of urinary diseases		
	fear of cancer of urinary syst.	U260	U26
	fear of urinary dis. NEC	U270	U27
513	disability/impairment urinary syst.	U280	U28
514	sympt./complt. urin. syst. NEC	U290	U29
515	acute pyelonefritis/pyelitis	U700	U70
516	cystitis/urinary inf. NEC	U710	U71
517	non-specific urethritis	U720	U72
518	malign. neoplasm kidney	U750	U75
519	malign. neoplasm bladder	U760	U76
520	malign. neoplasm urinary tract NEC	U770	U77
521	benign. neoplasm urinary tract	U780	U78
522	unspecified neopl. urin. tract	U790	U79
523	injury urinary tract	U800	U80
524	cong. anom. urinary tract	U850	U85

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
525	glomerulonephritis/nephrosis	U880	U88
526	orthostatic albuminuria	U900	U90
527	urin, calculus, all types/sites	U950	U95
528	abnormal urine test/findings NOS		
	glucosuria	U981	U98
	albuminuria	U982	U98
	abnorm. urine testfinding NEC	U989	U98
529	dis. urinary system NEC		
	chron. pyelitis/pyelonephritis	U991	U99
	urethral stricture	U992	U99
	uremia	U993	U99
	dis. urinary system NEC	U999	U99
W	PREGNANCY, FAMILY PLANNING (Pr/D/Pu/Fp)		
530	question of pregnancy (excl. W02)	W010	W01
531	fear of being pregnant	W020	W02
532	ante-partum bleeding		
	ante-partum bleeding < 16 wks	W031	W03
	ante-partum bleeding > 16 wks	W032	W03
533	vaginal discharge in pregn.	W040	W29
534	vomiting/nausea of pregn.	W050	W05
535	wanted pregnancy	W060	W01
536	morning-after pill	W100	W10
537	fam. plan./oral contraceptive	W110	W11
538	fam. plan./IUD	W120	W12
539	fam. plan./sterilization ♀	W130	W13
540	fam. plan./ NEC ♀	W140	W14
541	sub-/infertility ♀	W150	W15
542	heavy post-partum bleeding	W170	W17
543	complt. post partum NEC	W180	W18
544	sympt./complt. lactation	W190	W19
545	sympt./complt. breast in pregn. NEC	W200	W20
546	starting delivery	W210	W29
547	fear of complic of pregn./deliv.	W270	W27
548	disability/impairment Pr/D/Pu/Fp	W280	W28
549	sympt./complt. Pr/D/Pu/Fp NEC		
	albuminuria in pregn.	W291	W29
	excessive "weight" in pregn.	W292	W29
	sympt./complt. Pr/D/Pu/Fp NEC	W299	W29
550	infect. dis. urinary tr. pregn./puerp.	W700	U71
551	infect. dis. pregn. NEC	W710	W71
552	malign. neoplasm pregn.	W720	W72
553	benign. neoplasm pregn.	W730	W73
554	injuries complicating pregn.	W750	W75
555	cong. anom. of mother/complic. pregn.	W760	W76
556	non-obst. cond. complic. pregn. NEC	W770	W77
557	pregnancy: confirmed	W780	W78
558	unwanted pregnancy: confirmed	W790	W79
559	ectopic pregnancy	W800	W80
560	toxemia/(pre)eclampsia	W810	W81
561	abortion, imminens/spontan.		
	abortion imminens	W821	W03
	abortion spontaneous NOS	W822	W82
562	abortion, (il)legally induced	W830	W83

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
563	pregnancy, high risk		
	hyperemesis gravidarum	W841	W05
	cervical incompetence	W842	W84
	multiple pregnancy	W843	W84
	elderly primipare	W844	W84
	fetal malposition	W845	W84
	premature labour	W846	W84
	post-term pregnancy	W847	W84
	previous obst. cond. in pregn.	W848	W84
	high risk pregnancy NEC	W849	W84
564	normal delivery live born(s)	W900	W90
565	normal delivery dead born(s)	W910	W91
566	complicated delivery live born		
	abn. forces of labour complic. deliv.	W921	W92
	disproportion complic. deliv.	W922	W92
	umbilical cord complic.	W923	W92
	cesarean section live born	W924	W92
	ventouse extr./forceps deliv. live born	W925	W92
	perineal rupture	W926	W92
	episiotomy	W927	W92
	retained placenta	W928	W92
	complic. deliv. live born NEC	W929	W92
567	complicated delivery dead born	W930	W93
568	normal puerperium	W310	-
569	mastitis puerperalis	W940	W94
570	dis. of breast puerperium NEC	W950	W95
571	complic. puerperium NEC		
	fever puerp.	W961	W70
	thrombosis puerp.	W962	W96
	endometritis puerp.	W963	W70
	infection puerp. NEC	W968	W70
	complic. puerp. NEC	W969	W96/W99/W18/W29
572	diseases Pr/D/Pu NEC		
	abnormalities placenta	W991	W99
	dysmaturitas	W992	W92
	premature rupture of membranes	W993	W92
	dis. Pr/D/Pu NEC	W999	W99
X	FEMALE GENITAL SYSTEM and MAMMA		
573	genital pain ♀	X010	X01
574	menstruation absent/scanty		
	primary amenorrhea	X021	X05
	secondary amenorrhea, hypo-/am. NEC	X029	X05
575	menstruation excessive		
	menstr. excess. (regular)	X031	X06
	menstr. excess. (irregular/freq.)	X032	X07
	metrorrhagia	X051	X08
576	menstrual pain		
	dysmenorrhea	X041	X02
	ovulation pain	X042	X03
577	sympt./complt. menstruation NEC	X059	X08
578	premenstrual symptoms	X060	X09
579	sympt./complt. menopausal	X070	X11
580	postmenopausal bleeding	X080	X12
581	painful intercourse ♀	X090	X04
582	postcoital bleeding	X100	X13
583	vaginal discharge	X110	X14

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
584	sympt./complt. vagina NEC	X120	X15
585	sympt./complt. vulva	X130	X16
586	sympt./complt. pelvis ♀	X140	X17
587	lump/mass mamma ♀	X150	X19
588	pain mamma ♀	X160	X18
589	sympt./complt. mamma NEC ♀	X170	X20/X21
590	postponement of menstruation	X180	X10
591	fear genital/mamma dis. ♀		
	fear venereal dis. ♀	X230	X23
	fear of sexual dysfunction ♀	X240	X24
	fear genital cancer ♀	X250	X25
	fear mamma cancer ♀	X260	X26
	fear genital/mamma dis. ♀ NEC	X270	X27
592	disab./impairm. gen. syst. ♀	X280	X28
593	sympt./complt. gen. syst. ♀	X290	X29
594	syphilis, (genit. and NOS) ♀	X700	X70
595	gonorrhoea, (genit. and NOS) ♀	X711	X71
596	venereal dis. NEC ♀	X712	X90/X91/X99
597	urogenital candidiasis ♀	X720	X72
598	urogenital trichomoniasis ♀	X730	X73
599	pelvic inflammatory disease	X740	X74
600	carcinoma in situ cervix	X751	X75
601	cervix carcinoma NEC	X759	X75
602	mal. neopl. mamma ♀	X760	X76
603	mal. neopl. genit. syst. ♀	X770	X77
604	fibroid/myoma (uterus/cervix)	X780	X78
605	ben. neopl. mamma ♀	X790	X79
606	ben. neopl. genit. syst. ♀	X800	X80
607	unspec. neopl. genit. syst. ♀	X810	X81
608	injury genit. syst./mamma ♀	X820	X82
609	cong. anom. genit. syst./mamma ♀	X830	X83
610	vaginitis/vulvitis NOS	X840	X84
611	cervicitis/cerv. dis. NEC	X850	X85
612	abnormal PAP-smear	X860	X86
613	uterovaginal prolapse		
	uterovaginal prolapse	X871	X87
	stress-incontinence female	X872	X87
614	cystic breast (chronic)	X880	X88
615	premenstrual tension syndrome	X890	X89
616	gardnerella infection	X900	X99
617	infection/abscess gl. Bartholini	X991	X99
618	endometriosis	X992	X99
619	atrophial vaginae/vulvae	X993	X99
620	dis. genital syst./mamma NEC ♀		
	menarche sympt.	X994	X08
	dis. gen. syst. NEC ♀	X999	X99
Y	MALE GENITAL SYSTEM and MAMMA		
621	sympt./complt. gen. syst. ♂		
	pain in penis	Y010	Y01
	pain in testis/scrotum	Y020	Y02
	sympt./complt. penis NEC	Y040	Y04
	sympt./complt. testis/scrotum	Y050	Y05
622	discharge penis/urethra	Y030	Y03
623	sympt./complt. prostate	Y060	Y06
624	sympt./complt. potency	Y070	Y07
625	sympt./complt. sexual dysf. ♂	Y080	Y08

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
626	sub-/infertility σ	Y100	Y10
627	fam. plan. σ		
	fam. plan./sterilization σ	Y130	Y13
	fam. plan./NEC σ	Y140	Y14
628	sympt./complt. mamma σ	Y160	Y16
629	fear dis. gen. syst./mamma σ		
	fear sexual dysf. σ	Y240	Y24
	fear venereal dis. σ	Y250	Y25
	fear cancer gen. syst./mamma σ	Y260	Y26
	fear dis. gen. syst./mamma NEC σ	Y270	Y27
630	disab./impairm. gen. syst./mamma σ	Y280	Y28
631	sympt./complt. gen. syst./mamma NEC σ	Y290	Y29
632	syphilis σ	Y700	Y70
633	gonorrhoea σ	Y711	Y71
634	venereal disease NEC σ	Y712	Y72/Y76/Y99
635	urogenital trichomoniasis σ	Y720	Y99
636	prostatitis/seminal vesiculitis	Y730	Y73
637	orchitis/epididymitis	Y740	Y74
638	infection gen. syst. NEC σ	Y750	Y75/Y99
639	malign. neopl. prostate	Y761	Y77
640	malign. neopl. testis	Y762	Y78
641	malign. neopl. gen. syst. NEC σ	Y769	Y78
642	benign. neopl. gen. syst. σ	Y770	Y79
643	unspecif. neopl. gen. syst. σ	Y780	Y78
644	unspecif. neopl. mamma σ	Y790	Y78
645	injury gen. syst./mamma σ	Y800	Y80
646	phimosis/redund. prepuce	Y810	Y81
647	hypospadias	Y820	Y82
648	undesc. testicle/cryptorchism	Y830	Y83
649	cong. anom. gen. syst./mamma NEC σ	Y840	Y84
650	benign prostatic hypertrophy	Y850	Y85
651	hydrocele	Y860	Y86
652	dis. gen. syst./mamma NEC σ	Y990	Y99
Z	SOCIAL PROBLEMS		
653	poverty/financial probl.	Z010	Z01
654	problem food/water	Z020	Z02
655	probl. housing/neighborh.	Z030	Z03
656	probl. house moving	Z040	Z03
657	fear of having a social probl.		
	fear losing job	Z051	Z05
	fear of having soc. probl. NEC	Z270	Z27
658	probl. working conditions	Z052	Z05
659	probl. job/employment NEC	Z059	Z05
660	probl. w. being unemployed	Z060	Z06
661	probl. w. education	Z070	Z07
662	probl. soc. insurance/welfare	Z080	Z08
663	probl. legal/police	Z090	Z09
664	probl. health care syst./access	Z100	Z10
665	probl. soc./cultural syst.		
	discrimination (race/rel./sex/NEC)	Z111	Z04
	extramarital pregn.	Z112	Z04
	illegal stay	Z113	Z04
	probl. w. military service	Z114	Z04
	probl. soc./cult./rel. values NEC	Z119	Z04

nr.	description	ICPC-code NIVEL	ICPC-code OXFORD
666	relation problems partners		
	battering/violence by partner	Z121	Z12
	marital problems	Z122	Z12
	relation probl. partners NEC	Z129	Z12
667	problems w. behaving partner		
	alcohol abuse partner	Z131	Z13
	drug abuse partner	Z132	Z13
	behaving probl. partner NEC	Z139	Z13
668	probl. w. partner being ill	Z140	Z14
669	probl. loss of partner		
	probl. by divorce	Z151	Z15
	probl. by death partner	Z152	Z15
	probl. loss partner NEC	Z159	Z15
670	relation probl. with child		
	battering/neglecting child	Z161	Z16
	relation probl. parent-child	Z162	Z16
	probl. parent-child NEC	Z169	Z16
671	probl. behaving child	Z170	Z16
672	probl. with child being ill	Z180	Z18
673	loss or death of child	Z190	Z19
674	relation probl. parent/other family	Z200	Z20/Z12/Z16
675	probl. w. behaving parent/other family	Z210	Z21
676	probl. w. parent/other family being ill	Z220	Z22
677	loss parent/family		
	loss or death of parents	Z231	Z23
	loss or death of family	Z232	Z23
678	loss or death of friend	Z241	Z24
679	relation probl. w. friend	Z249	Z24
680	probl. assault/harmfull events	Z250	Z25
681	probl. with being ill	Z260	Z11
682	social handicap	Z280	Z28
683	social isolation	Z291	Z29
684	social problems		
	problems environmental pollution	Z292	Z29
	social problems NEC	Z299	Z29